



Original article

Adherence to the Mediterranean diet in children: Is it associated with economic cost?



G. Albuquerque^a, P. Moreira^{a,b,c}, R. Rosário^{d,e}, A. Araújo^a, V.H. Teixeira^{a,b},
O. Lopes^f, A. Moreira^{g,h}, P. Padrão^{a,c,*}

^a Faculty of Nutrition and Food Sciences, University of Porto, Porto, Portugal

^b Research Centre in Physical Activity, Health and Leisure, University of Porto, Porto, Portugal

^c EPI Unit – Institute of Public Health, University of Porto, Porto, Portugal

^d Education School, Child Study Centre, University of Minho, Braga, Portugal

^e Nursing School, University of Minho Braga, Portugal

^f Erdal Association, Guimarães, Portugal

^g Department of Immunology, Faculty of Medicine, University of Porto, Porto, Portugal

^h Department of Immunoallergology, Hospital of São João, Porto, Portugal

ARTICLE INFO

Article history:

Received 17 October 2016

Accepted 23 January 2017

Available online 1 March 2017

Keywords:

Diet cost
Mediterranean diet
Children

ABSTRACT

Objective: To assess how the diet cost is associated with socio-demographic factors and adherence to Mediterranean diet in children.

Methods: Data were obtained from a community-based survey of children selected from public elementary schools in Portugal. Of a total of 586 children attending these schools, 464 (6–12 years), were studied. Dietary intake was assessed by a 24 hour recall and the adherence to Mediterranean diet was evaluated through the KIDMED index. The cost of the diet was calculated based on the collection of food prices of a national leader supermarket, and expressed as Total Daily Cost (TDC) and Total Daily Cost-Adjusted for Energy (TDEC). Anthropometric measures were taken and socio-demographic data were gathered from a questionnaire filled by parents. Logistic regression was used to quantify the association between diet cost, socio-demographics and adherence to Mediterranean diet.

Results: The average TDC was 4.58€ (SD = 1.24). Most children (69.1%) reported medium adherence to Mediterranean diet, and 4.6% rated the higher score. TDC was higher for children with highest adherence to Mediterranean diet, compared to those with lowest adherence [TDC: OR = 5.70 (95% CI 1.53, 21.33), *p* for trend = 0.001; TDEC: OR = 2.83 (95% CI 0.89, 8.96, *p* for trend 0.018)]. No meaningful variation in the diet cost with age and parental education was observed.

Conclusion: Higher adherence to Mediterranean diet was associated with higher diet cost in children.

© 2017 PBJ-Associação Porto Biomedical/Porto Biomedical Society. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The Mediterranean Diet has been object of study since the 1950s¹ and is nowadays recognized for its numerous health benefits, namely protection against weight gain, obesity and diabetes mellitus type 2, cardiovascular diseases, certain tumors and other oxidative stress-related diseases.^{2,3}

However, its original defining-characteristics do not fully correspond to the diet practiced nowadays among populations living

in the Mediterranean region.⁴ This dietary pattern was firstly characterized by a high consumption of fruits, vegetables and grains; moderate consumption of dairy products and wine and low consumption (and amount) of red meat. Olive oil would be the main source of dietary lipids.¹ Regrettably, this dietary pattern is being replaced by unhealthier choices,^{5,6} similarly to what is happening around the world. A low consumption of fruits and vegetables⁷ parallel to a growing consumption of high-density energy foods – such as snacks, sugar-rich food, fast food and soft drinks – is well documented and associated with lifestyle changes.⁸ These dietary modifications contribute to poorer diet quality and have been indicted as a significant explanation for the rising obesity rates, specially concerning in children.⁸ The same phenomenon has been also documented in Portugal.⁹

* Corresponding author.

E-mail address: patriciapadrao@fcna.up.pt (P. Padrão).

Changes in the physical environment and food supply have been pointed recently as major causes of the Obesity epidemic,¹⁰ which is becoming to be perceived from an economic perspective. In fact, substantial research on diet cost has been performed in recent years, in the U.S.A. and some European countries. The results consistently show that the cost of food is a primary determinant of food choice^{11,12} and that higher energy density foods, which are less nutrient-rich, are associated with lower prices. On the other hand, low-energy-dense foods such as fruits and vegetables appear to be more expensive.^{11,13,14} In this context, healthier diets are associated with higher costs.^{14–17}

Taking into account the benefits of the Mediterranean diet, its health promoter potential, as well as the importance of price as food choice determinant, we aimed to estimate the cost of children's diet according to the degree of adherence to Mediterranean diet. To the best of our knowledge, this is one of the first European studies on this topic in children, which may provide new information to health professionals and policy makers so they can better educate and act toward the availability of healthy eating at low cost to the common citizen. The objectives of this study were to estimate the daily cost of diet and to quantify its association with socio-demographic factors and the degree of adherence to the Mediterranean diet in children.

Methods

Participants

The data were derived from a community-based survey of children selected from 7 of the eighty public elementary schools in the city of Guimarães, Portugal, between October 2007 and March 2008. Letters were distributed to all parents or guardians outlining the aims of the study along with a consent form. From the total of 586 children attending these schools, 464 (225 boys and 239 girls) between 6 and 12 years accepted to participate in the study. Anthropometric measurements and dietary data were collected from all consenting children and questionnaires surveying sociodemographics and lifestyle information were distributed among parents or educational guardians, of which 405 have answered (87%).

The study was approved by the University of Porto Ethics Committee, the schools where the study was carried out and the Portuguese Data Protection Authority (CNPD-Comissão Nacional de Protecção de Dados, process number 7613/2008).

Assessments

Height and weight were measured by previously trained health professionals or students, following international standardized procedures.^{18,19} Children wore light indoor clothing and were bare-footed. Weight was measured in an electronic scale, with an error of ± 100 g (Seca®, Model 703, Germany), and height was measured using a stadiometer, with the head in the Frankfort plane. BMI was calculated as weight (kg)/height² (m)²⁰ and children's weight status was categorized using the IOTF criteria and cut-points for BMI, defined specifically for sex and age.²¹ Only three categories were considered in analysis of results: normal, overweight and obesity.

Dietary intake information was assessed by a 24 hour recall, in which children were asked to recall all food and beverages consumed in the previous 24 h. A photographic manual of portion sizes and household measures (Manual of Food Quantification) was used²² as an auxiliary tool to estimate sizes of foods and beverages consumed.

KIDMED index⁵ was applied to verify the adherence level to the Mediterranean diet. This index was created according to the

Table 1

Kidmed index used to access the adherence to the Mediterranean diet.

Scoring	
+1	Eats a fruit or drinks fruit juice
+1	Eats a second fruit
+1	Has vegetables once
+1	Has vegetables a second time
+1	Eats fish
-1	Goes to a fast-food restaurant
+1	Eats pulses
+1	Eats rice and or pasta
+1	Has Breakfast
+1	Has a dairy product for breakfast
+1	Eats bread or cereals at breakfast
+1	Has at least 2 yoghurts or 40 g of cheese
-1	Eats pastries at breakfast
+1	Eats nuts
+1	Diet includes olive oil
-1	Eats sweets and candy

Mediterranean diet principles and provides a score ranging from 0 to 12 according 16 questions. Questions denoting a negative connotation with respect to the Mediterranean diet were assigned a value of -1 and those with a positive aspect, +1. This score was applied according to the food consumption in the previous 24 h, as described in Table 1. In accordance with the sum obtained, 3 classes were created: >8, high adherence to the Mediterranean diet; 4–7, medium adherence to the Mediterranean diet and ≤ 3 , low adherence to the Mediterranean diet (Table 1).

The socioeconomic information and family characteristics were collected from the survey distributed to the parents or educational guardians. It contained questions about gender and age of children and parents' education, recorded in five categories of years: 0, 1–4, 5–9, 10–12, and more than 12 years of formal education. This information was further grouped for analysis into four categories: up to 5 years, between 5 and 9 years, 10–12 years and more than 12 years of education.

Estimation of diet cost

The estimation of diet cost was divided in two tasks. Firstly, the collection of food prices, that took place between March and April of 2011. The source was an online supermarket, belonging to a Portuguese leader supermarket chain. Price data was obtained by gathering mean prices of correspondent food or package size, as well as the price per kilogram. Measurements were taken on regular prices, excluding discounts. In the case of composed dishes, diet costs were calculated using recipes available in Food Processor Plus® database (most of them previously adapted from traditional Portuguese recipes) and from a Portuguese website of traditional recipes.²³ The price of the drinking water was estimated by the median price of the bottled natural mineral water and of the municipal water.

After this procedure, food items were assigned into different groups according to the staple food that was in its origin, and the median price per gram was computed. For example, the price of rice was obtained by calculating the median of the prices of the various brands and types available in the supermarket webpage. The choice for using the median rather than the average price was based on the fact that it better represents the central values, minimizing the effect of the very high and very low prices for each group. Finally, the cost of each meal was calculated according to the contribution of each and every food ingredient taking into account its proportion.

At dietary level, two variables were created: "Total daily cost" (TDC), representing the cost of each individual's diet and obtained by summing the cost of each meal, and "Total daily cost adjusted for energy" (TDEC) which eliminated the possible differences in

Table 2
Participants' characteristics.

	Boys (n = 225)		Girls (n = 239)		Total (n = 464)	
	n	%	N	%	n	%
<i>Age (years)</i>						
6–7	52	23.1	55	23.0	107	23.1
8	59	26.2	68	28.5	127	27.4
9	63	28.0	52	21.8	115	24.8
10–12	51	22.7	64	26.8	115	24.8
<i>Mother educational level (years)*</i>						
<5	41	18.2	44	21.8.4	85	22.1
5–9	77	34.2	83	41.1	160	41.7
10–12	40	17.8	50	24.8	90	23.4
>12	24	10.7	25	12.4	49	12.8
<i>Father educational level (years)*</i>						
<5	40	22.6	52	26.9	92	19.6
5–9	86	48.6	77	39.9	263	56.0
10–12	29	16.4	42	21.8	71	15.1
>12	22	12.4	22	11.4	44	9.4
<i>Kidmed score*</i>						
Low	62	28.6	56	23.8	119	26.3
Medium	149	68.7	164	69.8	313	69.1
High	6	2.7	15	6.4	21	4.6

* For these variables, sample size is lower.

costs associated with different energetic intake between individuals. TDEC was computed dividing TDC by the energy consumed (€/kcal) and expressed as €/1000 kcal, in order to point differences not seen with TDC.

Statistical analysis

The statistical analyses were performed using the software *Statistical Package for the Social Sciences* (SPSS), version 17.0. Descriptive statistics were used to characterize the sample (mean and standard deviations). A total of 342 subjects for whom assessment of overall dietary intake was available were considered for data analysis.

Unconditional binary logistic regression models were fitted to estimate the magnitude of the association between diet cost (considering two categories, using the median value as the cut-off) and socio-demographic characteristics (sex, age and parents' education) and the degree of adherence to the Mediterranean Diet (Kidmed score).⁵ The level of significance was set at 0.05.

Results

Participant characterization

In this sample of 464 children (51.5% girls), the prevalence of overweight and obesity were 23.3% and 7.3%, respectively. Approximately two-thirds of the study population had parents with less than 10 years of formal education. The majority of children reported a medium adherence level to Mediterranean diet (69.1%), and only 4.6% rated a higher adherence score (Table 2).

Diet cost

The average (\pm standard deviation) TDC was 4.58 (\pm 1.24) € and the average TDEC was 2.17 (\pm 0.42) €/1000 kcal (Table 3). No meaningful differences were observed between gender, age or parental education regarding TDC and TDEC. Accordingly, no meaningful or consistent variation in the diet cost with age and parents' education was observed (Table 3). Concerning Kidmed score, it is noticeable an increasing in the cost of the diet with increasing level of adherence to the Mediterranean Diet (TDC of 4.79€ in high adherence

category vs. 4.09€ in low adherence category, $p=0.471$; TDEC of 2.19€/1000 kcal in high adherence category vs. 2.10€/1000 kcal in low adherence category, $p=0.047$). Considering children reporting the higher adherence to Mediterranean diet in comparison with those with the lowest adherence, the odds favoring higher diet cost was 5.70 (95% CI 1.53–21.33, p for trend = 0.001) for TDC and 2.83 (95% CI 0.89–8.96, p for trend = 0.018) for TDEC (Table 3).

Discussion

The present study showed that a higher adherence to the Mediterranean diet was associated with higher diet cost. The average TDC found in this sample of school-aged children was 4.58€. As far as we know, only few studies^{24–26} have focused on the estimation of diet cost among children. In a study conducted within a Spanish sample aged 2–24 years old, researchers reported a mean daily diet cost of 3.16€ (data collected in the year 2000) and, more recently, two studies^{25,26} conducted within the DONALD cohort study, which includes German children, found daily diet costs also close to 3€. Studies among adults in Europe and US, reported average prices higher than 5€.^{12,14,17}

One out every twenty children in this study reported high adherence level to the Mediterranean dietary pattern, results that are in line with previous studies in Mediterranean countries.²⁷ The explored relationship between diet cost and adherence to the Mediterranean diet has brought interesting results, as it was verified an increasing cost with a higher adherence to Mediterranean diet. A similar study in Spanish youth (participants were aged between 2 and 24 years)²⁴ has found similar results. In literature, much has already been discussed on the higher cost of healthy diets,^{13,14,25,29–31} of which Mediterranean diet is a good example.^{12,15,16,28–31} There are a few pointed aspects underlying this phenomenon, which are important to refer. The first is the content in energy dense foods that, apparently cheapen the diet,^{12–14,26,29} association that has already been demonstrated in the current sample of school-aged children.³² In another study, Rydén et al. verified higher cost of diet associated with its healthiness (assessed using the Healthy Eating Index), in which energy-density was low.³¹ Secondly, variety, a characteristic of healthy dietary patterns, is associated with a large number of food groups and foods among groups.³³ Hence, and according to the literature, healthier groups are associated with higher costs, making the diet more expensive. In accordance to our results, food items that play an important role in the Mediterranean diet such as fruits and vegetables, but also fish, were associated to higher costs in different studies.^{12,30,31} Moreover, the contribution of healthier options within the latter group, such as lean meats and low-fat products, was further associated in the current study with an increased cost. Data in the literature relates higher costs and healthier options within the same food group.³¹ However, when analyzing Mediterranean diet in this perspective, Drewnowski et al.²⁸ stated that *not all nutrient-rich foods necessarily cost more* and so, *it should be possible to construct a Mediterranean-style diet using the lower cost options in every category.*

A possible consequence of higher cost of healthy diets, such as the Mediterranean, is the higher prevalence of poor quality meals within low socioeconomic position (SEP) families, who cannot afford to spend much of the family budget on healthy foods.^{7,34} A study conducted in Portugal in 2006 by Moreira et al. showed that a higher education was positively linked to a better dietary quality, represented by a higher frequency of milk, vegetable soup, vegetables, fruit and fish consumption,³⁵ all of which are commonly consumed within the Mediterranean pattern. The three most common SEP indicators are education, occupation and income. However, in the present study, only parental education was evaluated, and no significant association with diet cost was found. A

Table 3

Diet cost and odds ratio for diet cost according to sex, age, weight status, parents' education and Kidmed score.

	Total day cost				Total day energy cost			
	Mean	SD	OR [†]	95% CI	Mean	SD	OR [†]	95% CI
Sex								
Girls	4.42	1.19	1	ref	2.18	0.37	1	ref
Boys	4.73	1.28	1.49	(0.88–2.52)	2.16	0.48	0.93	(0.59–1.48)
	<i>p</i> = 0.475		<i>p</i> = 0.259		<i>p</i> = 0.485		<i>p</i> = 0.872	
Age (years)								
06/jul	4.48	1.33	1	ref	2.15	0.40	1	ref
8	4.92	1.14	2.55	(1.12–5.90)	2.23	0.45	1.32	(0.66–2.63)
9	4.47	1.16	0.79	(0.35–1.78)	2.22	0.41	1.96	(0.98–3.95)
10/dez	4.43	1.30	1.31	(0.59–2.94)	2.08	0.41	1.10	(0.53–2.25)
	<i>p</i> = 0.468		<i>p</i> for trend = 0.868		<i>p</i> = 0.424		<i>p</i> for trend = 0.803	
Mother educational level (years)								
<5	4.34	1.43	1	ref	2.13	0.55	1	ref
5–9	4.62	1.17	1.02	(0.49–2.14)	2.17	0.37	1.08	(0.56–2.08)
10/dez	4.48	1.23	0.55	(0.19–1.54)	2.12	0.33	0.49	(0.20–1.19)
>12	4.55	1.08	0.58	(0.18–1.19)	2.11	0.36	0.88	(0.31–2.49)
	<i>p</i> = 0.464		<i>p</i> for trend = 0.368		<i>p</i> = 0.456		<i>p</i> for trend = 0.295	
Father educational level (years)								
<5	4.29	1.36	1	ref	2.09	0.43	1	ref
5–9	4.54	1.23	1.41	(0.69–2.89)	2.14	0.41	1.19	(0.63–2.26)
10/dez	4.76	1.10	3.51	(1.25–9.85)	2.23	0.41	2.19	(0.93–5.14)
>12	4.41	1.12	1.47	(0.42–5.13)	2.12	0.32	1.29	(0.43–3.86)
	<i>p</i> = 0.464		<i>p</i> for trend = 0.167		<i>p</i> = 0.527		<i>p</i> for trend = 0.279	
Kidmed score								
Low	4.09	1.26	1	ref	2.10	0.34	1	ref
Medium	4.70	1.21	2.69	(1.32–5.48)	2.19	0.45	1.79	(1.00–3.20)
High	4.79	1.10	5.70	(1.53–21.33)	2.19	0.47	2.83	(0.89–8.96)
	<i>p</i> = 0.471		<i>p</i> for trend = 0.001		<i>p</i> = 0.047		<i>p</i> for trend = 0.018	
Total	4.58	1.24			2.17	0.42		

SD: standard deviation; OR: odds ratio; 95% CI: 95% confidence interval.

† Adjusted for sex, age, parental education level and Kidmed score.

subsequent study covering disposable family income could add a vital step to overtake this limitation, as it was described as a better SEP marker with regard to food budget choice by Rydén et al.³¹

Some methodological limitations are worth noting. First, dietary intake and cost estimates were derived from a 24 h recall. The use of this instrument may have compromised the collected information, since it has been recognized that children younger than 8 years-old may not accurately recall foods and estimate portion sizes.³⁶ Nevertheless, only approximately a quarter of our sample was younger than 8 years-old. In addition, single 24 h recall may not represent the usual dietary intake and may fail to include foods and beverages that are either forgotten or consumed infrequently, influencing the KIDMED score. As most participants were not able to detail the ingredients of the recipes (and the fat used for cooking, especially concerning for the item “diet includes olive oil”) some assumptions were made, based on traditional Portuguese recipes. However, this fact may be diluted as a whole, given the sample size.

Second, the food price collection was based only in one source, which may have been minimized by the fact that the supermarket chain where the prices were collected has the largest share of food market in Portugal. Also, food prices were collected three years after the dietary survey was conducted, and some changes might have occurred in that period. However, this fact is mitigated by the expectation that prices have varied in a similar way, since the *Value Added Tax* has not changed during the period elapsed. In addition, the seasonal variability of fresh foods production, namely fruits and vegetables, may have compromised the accuracy of the prices collected, due to the fact that food prices were gathered in a different season of the survey. Nevertheless, it is expected that, in all seasons, the prices of fresh products whose production is seasonally variable, vary so that the rise in some prices is offset by a decline of others. In turn, energy dense foods are the

most resistant to inflation.¹⁴ In some cases, a lack of details about food consumed determined the need to use average values (grouping fresh and tanned foods, for instance). The use of the average price does not capture differences between brand foods and consequently underestimate the variability of food prices and of the costs associated with individual food consumption,³⁷ which may contribute to explain the lack of association between diet cost and parental education. Furthermore, since the current study has a cross-sectional design, we are limited to demonstrate associations and the direction of these associations.

Nevertheless, this was an original study that brought important insight into dietary costs of children and its association with socio-demographics and adherence to Mediterranean diet, a topic becoming significant in terms of Public Health nutrition worldwide, especially given the economic crisis that has settled in Europe and Mediterranean countries. As food price is becoming a primary determinant of food choice, this new information should concern and be considered by nutritional health care providers and public health authorities. Nutritional education and promotion of healthy eating should be provided in a cost effective manner.

Conflict of interests

There were no conflicts of interest declared.

Author's contribution

GA, PM and PP designed the study; GA, PM, RR, AA, VT, RB, OL, AM and PP conducted the study; RR, AA and OL collected the data; GA, PM, VT and PP analyzed the data; GA and PP wrote the manuscript. All authors read and approved the final manuscript.

References

1. Keys A. Mediterranean diet and public health: personal reflections. *Am J Clin Nutr.* 1995;61 6 Suppl.:1321S–3S.
2. Schröder H. Protective mechanisms of the Mediterranean diet in obesity and type 2 diabetes. *J Nutr Biochem.* 2007;18:149–60.
3. Barros R, Moreira A, Fonseca J, Ferraz de Oliveira J, Delgado L, Castel-Branco MG, et al. Adherence to the Mediterranean diet and fresh fruit intake are associated with improved asthma control. *Allergy.* 2008;63:917–23.
4. CIHEAM/FAO. Mediterranean food consumption patterns: diet, environment, society, economy and health. A White Paper Priority 5 of Feeding Knowledge Programme. Expo Milan 2015. Rome: CIHEAM-IAMB, Bari/FAO; 2015.
5. Serra-Majem L, Trichopoulou A, de la Cruz JN, Cervera P, Álvarez AG, La Vecchia C, et al. Does the definition of the Mediterranean diet need to be updated? *Public Health Nutr.* 2004;7:927–9.
6. Bonaccio M, Bes-Rastrollo M, de Gaetano G, Iacoviello L. Challenges to the Mediterranean diet at a time of economic crisis. *Nutrition, Metabolism and Cardiovascular Diseases.*
7. Bihan H, Castetbon K, Mejean C, Peneau S, Pelabon L, Jellouli F, et al. Sociodemographic factors and attitudes toward food affordability and health are associated with fruit and vegetable consumption in a low-income French population. *J Nutr.* 2010;140:823–30.
8. Mwangi AM, den Hartog AP, Mwadime RK, van Staveren WA, Foeken DW. Do street food vendors sell a sufficient variety of foods for a healthful diet? The case of Nairobi. *Food Nutr Bull.* 2002;23:48–56.
9. Moreira P, Santos S, Padrão P, Cordeiro T, Bessa M, Valente H, et al. Food patterns according to Sociodemographics, physical activity, sleeping and obesity in Portuguese children. *Int J Environ Res Public Health.* 2010;7:1121–38.
10. James WPT. The fundamental drivers of the obesity epidemic. *Obesity Rev.* 2008;9 Suppl. 1:6–13.
11. Rao M, Afshin A, Singh G, Mozaffarian D. Do healthier foods and diet patterns cost more than less healthy options? A systematic review and meta-analysis. *BMJ Open.* 2013;3. December 1, 2013.
12. Darmon N, Briand A, Drewnowski A. Energy-dense diets are associated with lower diet costs: a community study of French adults. *Public Health Nutr.* 2004;7:21–7.
13. Drewnowski A, Darmon N. Food choices and diet costs: an economic analysis. *J Nutr.* 2005;135:900–4.
14. Rehm CD, Monsivais P, Drewnowski A. Relation between diet cost and Healthy Eating Index 2010 scores among adults in the United States 2007–2010. *Prev Med.* 2015 Apr;73:70–5.
15. Waterlander WE, De Haas WE, Van Amstel I, Schuit AJ, Twisk JW, Visser M, et al. Energy density, energy costs and income – how are they related? *Public Health Nutr.* 2010;13:1599–608.
16. Andrieu EDN, Drewnosky A. Low-cost diets: more energy, fewer nutrients. *Eur J Clin Nutr.* 2006;60:434–6.
17. Morris MA, Hulme C, Clarke GP, Edwards KL, Cade JE. What is the cost of a healthy diet? Using diet data from the UK Women's Cohort Study. *J Epidemiol Commun Health.* 2014, 2014, July 22.
18. Marfell-Jones M, Olds T, Stewart A, Carter JL. International Standards for Anthropometric Assessment; 2006.
19. Rudolf MCJ, Walker J, Cole TJ. What is the best way to measure waist circumference? *Int J Pediatric Obesity.* 2007;2:58–61, 2007/01/01.
20. World Health Organization. The International Classification of adult underweight, overweight and obesity according to BMI; 2008.
21. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *Br Med J.* 2000;320:1240–3.
22. Marques M, Pinho O, De Almeida MDV. Manual de Quantificação de alimentos; 1996.
23. *Gastronomias. Roteiro Gastronómico de Portugal; 1997 [Internet]. 1997;http://gastronomias.com/. Accessed 11 June 2011.*
24. Schroder H, Gomez SF, Ribas-Barba L, Perez-Rodrigo C, Bawaked RA, Fito M, et al. Monetary diet cost, diet quality, and parental socioeconomic status in Spanish Youth. *PLoS One.* 2016;11:e0161422.
25. Alexy U, Schwager V, Kersting M. Diet quality and diet costs in German children and adolescents. *Eur J Clin Nutr.* 2014;68:1175–6.
26. Alexy U, Bolzenius K, Kopper A, Clausen K, Kersting M. Diet costs and energy density in the diet of German children and adolescents. *Eur J Clin Nutr.* Dec 2012;66:1362–3.
27. Lazarou C, Panagiotakos DB, Matalas A-L. Level of adherence to the Mediterranean diet among children from Cyprus: the CYKIDS study. *Public Health Nutr.* 2009;12:991–1000.
28. Drewnowski A, Eichelsdoerfer P. The Mediterranean diet: does it have to cost more? *Public Health Nutr.* 2009;12(9 SPEC. ISSUE 9A):1621–8.
29. Maillot M, Darmon N, Vieux F, Drewnowski A. Low energy density and high nutritional quality are each associated with higher diet costs in French adults. *Am J Clin Nutr.* 2007;86:690–6. September 1, 2007.
30. Lopez CN, Martinez-Gonzalez MA, Sanchez-Villegas A, Alonso A, Pimenta AM, Bes-Rastrollo M. Costs of Mediterranean and western dietary patterns in a Spanish cohort and their relationship with prospective weight change. *J Epidemiol Commun Health.* 2009;63:920–7. November 1, 2009.
31. Rydén PJ, Hagfors L. Diet cost, diet quality and socio-economic position: how are they related and what contributes to differences in diet costs? *Public Health Nutr.* 2011:1–13.
32. Faria AP, Albuquerque G, Moreira P, Rosário R, Araújo A, Teixeira V, et al. Association between energy density and diet cost in children. *Porto Biomed J.* 2016;1:106–11, 7//.
33. Drescher LS, Thiele S, Mensink GBM. A New index to measure healthy food diversity better reflects a healthy diet than traditional measures. *J Nutr.* 2007;137:647–51. March 1, 2007.
34. Darmon N, Drewnowski A. Does social class predict diet quality? *Am J Clin Nutr.* 2008;87:1107–17. May 1, 2008.
35. Moreira P, Padrão P. Educational, economic and dietary determinants of obesity in Portuguese adults: a cross-sectional study. *Eating Behav.* 2006;7:220–8.
36. Livingstone MBE, Robson PJ. Measurement of dietary intake in children. *Proc Nutr Soc.* 2000;59:279–93, 2000/005/001.
37. Darmon N, Drewnowski A. Contribution of food prices and diet cost to socioeconomic disparities in diet quality and health: a systematic review and analysis. *Nutr Rev.* 2015;73:643–60.