



## Abstracts 12th YES Meeting



Internal Medicine Paralell Oral Session

Friday, September 15th, 14h00

**PS004**

### Why novel methods are not always the best? – Multifactorial analysis of hyperandrogenism in women



Sylwia Gajda\*, Damian Sieńko,  
Urszula Ambroziak

*Ist Endocrinology Clinic of Hospital affiliated to the  
Medical University of Warsaw, Poland*

*E-mail address: [sylviagajda@gmail.com](mailto:sylviagajda@gmail.com)  
(S. Gajda).*

**Aim:** The aim of the work was to compare different methods of hormones evaluation, including blood and saliva samples and the reliability of those methods in diagnosing hyperandrogenism among women caused by various reasons.

**Introduction:** Hyperandrogenism among women is a common problem. There are different hormones that can be evaluated with various methods to diagnose and monitor patients. Less invasive and quicker methods of screening, like salivary samples, more and more are used in medicine. However, they may be not as accurate as expected.

**Methods:** 39 women with clinical or biochemical hyperandrogenism and 29 healthy individuals in control group were enrolled. The diagnosis of hyperandrogenic syndrome covered: 13 patients with polycystic ovary syndrome (PCOS), 23 with idiopathic hyperandrogenism, 2 with congenital adrenal hyperplasia and 1 adrenal cortical carcinoma. Assessed hormones included: serum total testosterone (T) measured with liquid chromatography-tandem mass spectrometry LC-MS and immunoassay, salivary T by Salimetrics test; DHEA-S, androstendione and 17-OH Progesterone by LC-MS and immunoassay.

**Results:** In 9 out of 38 patients' results of salivary testosterone showed normal levels, while with LC-MS method increased levels were depicted in the same women. Similarly, 41% women with hyperandrogenism had elevated testosterone with ELISA method, whilst having Salimetrics test results within normal range. In 28% normal testosterone levels measured by LC-MS method, DHES-S was elevated. All patients with elevated androstendione presented with elevated concentration of either testosterone or DHEA-S. Elevated DHEA-S was observed in 56.5% patients with FHS and 15.4% with PCOS.

**Conclusion:** Salivary testosterone is not a sufficient method in diagnosing biochemical hyperandrogenism. Measurement of serum testosterone by LC-MS itself is not enough to diagnose biochemical hyperandrogenism. DHEA-S should also be evaluated when hyperandrogenism is suspected. Androstendione measurement is not obligatory in diagnosis. This is the first study analyzing numerous hormones with various methods in patients with hyperandrogenism caused by different diseases.<sup>1–4</sup>

### References

1. Weisser JJ, Hansen CH, Poulsen R, Larsen LW, Cornett C, Styrisshave B. Two simple cleanup methods combined with LC-MS/MS for quantification of steroid hormones in in vivo and in vitro assays. *Anal Bioanal Chem.* 2016;408:4883–95.
2. Janssens G, Manginckx S, Courtheyn D, De Kimpe N, Matthijs B, Le Bizec B. Simultaneous detection of androgen and estrogen abuse in breeding animals by gas chromatography-mass spectrometry/combustion/isotope ratio mass spectrometry (GC-MS/C/IRMS) evaluated against alternative methods. *J Agric Food Chem.* 2015;63:7574–81.
3. Leerasing P, Wongwananuruk T, Indhavivadhana S, Techatrasak K, Rattanachaiyanont M, Angsuwathana S. Correlation of clinical and biochemical hyperandrogenism in Thai women with polycystic ovary syndrome. *J Obstet Gynaecol Res.* 2016;42:678–83, <http://dx.doi.org/10.1111/jog.12945> [Epub 2016 February 18].
4. Wang C, Catlin DH, Demers LM, Starcevic B, Swerdloff RS. Measurement of total serum testosterone in adult men: comparison of current laboratory methods versus liquid chromatography-tandem mass spectrometry. *J Clin Endocrinol Metab.* 2004;89:534–43.

<http://dx.doi.org/10.1016/j.pbj.2017.07.005>

**PS151**

### Comparison between effects of antibiotics, NSAIDs and their mixture on the growth of microorganisms



S. Bhattacharya\*, Y. Akula, G.M. Mitongo,  
Q. Khorram

*Lviv National Medical University, Ukraine*

*E-mail address: [shayariq5@gmail.com](mailto:shayariq5@gmail.com)  
(S. Bhattacharya).*

**Aim:** To compare the effects of antibiotics, NSAIDs and their mixture on the growth of microorganisms.

**Introduction:** Commonly, when a patient has an infection, doctors prescribe NSAIDs for pain and inflammation that may be caused by infection as a part of symptomatic treatment. And antibiotics are also prescribed as an etiological treatment. Our experiment that was performed last year came to a conclusion

that NSAIDs, notably: Aspirin, Ibuprofen and Diclofenac could inhibit the growth of some microorganisms including *Staphylococcus aureus*, *Escherichia coli* and *Candida albicans*. These results, although performed in vitro were promising especially with the growing rate of bacterial resistance towards antimicrobial agents.

**Methods:** We used antibiotics: Penicillin, Gentamicin and Ceftriaxone; NSAIDs-non-selective: Aspirin, Diclofenac and Ibuprofen and COX-2 selective: Celecoxib. Samples were taken from the oral cavity of patients with liver diseases. Cultures were made of the samples taken and they were inoculated onto an agar plate. Then three well were made in the agar plate: in the first well we put an NSAIDs, second well with an antibiotic and in the third we put the mixture of both NSAID and antibiotic. The agar plates were placed into an incubator for 24 h at a temperature of 37 °C. The experiment was done twice to get accurate results.

**Results:** The analysis of the obtained results shows that in group 1 (antibiotics) was the highest inhibition  $39.3 \pm 3.6$  mm, in the group 2 in which there were NSAIDs gave the results as shown  $31.7 \pm 4.1$  mm, and last investigative group 3 with mixture was  $27.3 \pm 1.8$ .

**Conclusion:** From the obtained results we can conclude that a mixture of NSAIDs and antibiotics does not improve antibacterial effect of antibiotics. In fact, NSAIDs seem to even lower the efficacy of antimicrobial drugs. Special attention should be paid while administering NSAIDs to patients who are on antibiotic therapy since the combination of these two groups of drugs lower the antimicrobial effect.

**Acknowledgements:** Assistant professor Marta Panas (our scientific advisor).

<http://dx.doi.org/10.1016/j.pbj.2017.07.006>

### PS003

#### Comparison of metabolic syndrome rates in living-donor and deceased-donor kidney recipients – A three-year follow-up

Aleksandra Chabior\*, Jolanta Gozdowska, Ewelina Jędrych, Magdalena Durlik

Department of Transplantation Medicine, Nephrology and Internal Diseases, Medical University of Warsaw, Warsaw, Poland  
E-mail address: [o.chabior@gmail.com](mailto:o.chabior@gmail.com)  
(A. Chabior).

**Aim:** Comparison of MS rates in kidney recipients.

**Introduction:** Metabolic syndrome (MS) is characterized by coexistent pro-atherogenic disorders and insulin resistance. MS also increases cardiovascular risk.

**Methods:** A total of 112 living-donor ( $n=54$ ) and deceased-donor ( $n=58$ ) kidney transplant recipients were evaluated for metabolic syndrome (MS) in months 6, 12, and 36. The National Cholesterol Education Program – Adult Treatment Panel III (NCEP-ATP III) criteria were used. Both groups were compared in terms of MS rates. Moreover, correlations between MS and other parameters (age, gender, dialysis type and duration, donor type, immunosuppressant drugs, acute rejection episodes, smoking, levels of triglycerides, uric acid, creatinine, eGFR, and proteinuria) were evaluated. The chi-square, McNemar's test, Student's *t* test, Welch's *t* test, Mann-Whitney *U* test, Fisher's test, and Shapiro-Wilk test were used in the statistical analysis.

**Results:** MS rates following living-donor (LD) and deceased-donor (DD) kidney transplantation (KTx) in months 6, 12, and 36 were 0.148 vs. 0.276; 0.173 vs. 0.316; 0.235 vs. 0.182, respectively. MS rates in LD KTx recipients were lower than those in DD KTx recipients in months 6 and 12, especially in males (0.14 vs.

0.379;  $p=0.0251$ ), but they increased systematically in subsequent months of follow-up. MS was more commonly diagnosed in older recipients ( $p=0.019$ ), with lower MDRD eGFR values ( $p=0.009$ ), who received more anti-hypertensive drugs ( $p=0.046$ ). The dialysis type, donor type and the number of transplantations had no effect. The logistic regression model indicated that the factors contributing to MS were elevated uric acid levels and proteinuria.<sup>1,2</sup>

#### Conclusion:

1. MS rates in LD KTx recipients in month 6 and 12 following transplantation are lower than those in DD KTx recipients.
2. MS rates in LD KTx recipients tended to progressively increase during follow-up.
3. MS was more common in older patients with poorer kidney function, higher uric acid levels and proteinuria.

#### References

1. Prasad GV, Huang M, Silver SA, Al-Lawati AI, Rapi L, Nash MM, et al. Metabolic syndrome definitions and components in predicting major adverse cardiovascular events after kidney transplantation. *Transpl Int*. 2015;28:79–88.
2. Nagaraja P, Sharif A, Ravindran V, Baboolal K. Long-term progression of abnormal glucose tolerance and its relationship with the metabolic syndrome after kidney transplantation. *Transplantation*. 2014;97:576–81.

<http://dx.doi.org/10.1016/j.pbj.2017.07.007>

### PS073

#### Association between body composition and magnesium level in middle aged women



B. Ilincic, Dragana Oluski\*

Pathophysiology, Faculty of Medicine of the University of Novi Sad, Serbia

E-mail address: [dragana.oluski@gmail.com](mailto:dragana.oluski@gmail.com)

(D. Oluski).

**Aim:** Aim of the study was to compare total magnesium serum concentration between subjects with increased fat mass in the body composition and subjects with normal body composition as to determine the association between total magnesium serum level and parameters of the body composition and glucose metabolism.

**Introduction:** Metabolic disorders and chronic diseases may associate alterations in body composition and could be related with disturbances of the magnesium blood level. Obesity is a chronic disease characterized by disturbances of the body composition, commonly associated with disorders of carbohydrate metabolism.

**Methods:** The study included 40 women with body composition disturbances (increased percentage of the total fat mass) and 20 age matched women with normal percentage of the total fat mass. All subjects underwent analysis of the components of body composition [bioelectrical impedance analysis, fat mass percentage (FAT%), fat free mass percentage (FFM%)], laboratory analysis of blood samples (automated analyzer systems) with determining the parameters of glucose metabolism and total magnesium serum concentration. Insulin resistance index (HOMA-IR) was calculated using equation involving fasting insulin and glucose concentration.

**Results:** Women with increased percentage of the total fat mass had significantly lower total magnesium serum concentration compared to control group ( $0.83 \pm 0.07$  vs.  $0.9 \pm 0.06$  mmol/l,  $p=0.00$ ). Moderate correlation was found between serum concentrations of total magnesium and FAT% ( $r=-0.47$ ,  $p=0.00$ ), FFM% ( $r=0.44$ ,  $p=0.00$ ), fasting insulin levels ( $r=-0.43$ ,  $p=0.00$ ) and HOMA-IR ( $r=-0.44$ ,  $p=0.00$ ).

**Conclusion:** Women with increased total fat mass in the body composition have significantly lower total magnesium serum concentration, compared to women with normal body composition.



Additionally to increased fat mass, insulin resistance is associated with total magnesium level in middle aged women.

<http://dx.doi.org/10.1016/j.pbj.2017.07.008>

### PS137

#### Increased paraoxonase and arylesterase activity in thyroiditis patients compared to healthy individuals



S.S.K. Marasinghe<sup>1,\*</sup>, R. Sivakanesan<sup>2</sup>

<sup>1</sup> Postgraduate Institute of Science, University of Peradeniya, Peradeniya, Sri Lanka

<sup>2</sup> Department of Biochemistry, University of Peradeniya, Peradeniya, Sri Lanka

E-mail address: [sanjila.marasinghe@gmail.com](mailto:sanjila.marasinghe@gmail.com) (S.S.K. Marasinghe).

**Aim:** The aim of this study was to assess whether there is a significant difference in paraoxonase and arylesterase activities and distribution of phenotypes in thyroiditis patients compared to healthy volunteers.

**Introduction:** Human serum paraoxonase 1 (PON1; EC3.1.1.2) is an antioxidant enzyme showing both paraoxonase and arylesterase activities. The PON1-192 polymorphism has two isoforms, namely PON1 Q and PON1 R. PON1 Q contains a glutamine at position 192. PON1 R contains an arginine at position 192. It shows a 6 fold higher activity towards paraoxon hydrolysis compared to Q isoform. Arylesterase activity is similar in both isozymes. The R allozyme shows a greater degree of stimulation of its paraoxon-hydrolyzing activity by 1 M NaCl than does the Q allozyme. The ratio of Salt stimulated PON 1 activity/Arylesterase activity (P/A ratio) is trimodally distributed. The three modes correspond to paraoxonase phenotypes, QQ, QR and RR.

**Methods:** Fifty thyroiditis patients and one hundred and thirty seven apparently healthy individuals were enrolled in this study. Serum samples of both groups were analysed for basal paraoxonase activity, salt stimulated paraoxonase activity (with 1 M NaCl) and arylesterase activity (spectrophotometrically). P/A ratio was used to assess the phenotypes (dual substrate method).

**Results:** Basal PON 1 activity ( $205.27 \pm 115.00$  U/l vs.  $251.1 \pm 129.6$  U/l,  $p=0.002$ ) and arylesterase activity ( $159.53 \pm 37.11$  vs.  $177.59 \pm 46.90$ ,  $p=0.024$ ) was significantly higher in thyroiditis patients compared to healthy volunteers. Percentage of QQ phenotype was significantly higher in thyroiditis patients compared to healthy individuals. Percentage of QR phenotype was significantly lower in thyroiditis patients compared to healthy individuals. There was no difference in percentage of RR phenotype in thyroiditis patients and healthy individuals.

**Conclusion:** Serum PON 1 activity and arylesterase activity was significantly higher in thyroiditis patients compared to healthy individuals. Percentage distribution of phenotypes in thyroiditis patients was significantly different from healthy individuals.

<http://dx.doi.org/10.1016/j.pbj.2017.07.009>

### PS111

#### The relationship between dyslipidemia and disease activity in Iranian population with systemic lupus erythematosus



Sepideh Hajian<sup>1</sup>, Mohammad Ali Hosseini<sup>2,\*</sup>, Sara Khosraviani<sup>2</sup>, Farnaz Tavasoli<sup>2</sup>

<sup>1</sup> Department of Nephrology, Hasheminejad Center, Iran University of Medical Sciences, Tehran, Iran

<sup>2</sup> Student research Committee, School of medicine, Qazvin University of Medical Sciences, Qazvin, Iran  
E-mail address: [smahoseini@gmail.com](mailto:smahoseini@gmail.com) (M.A. Hosseini).

**Aim:** This study was designed for evaluating the relationship between dyslipidemia and diseases activity in systemic lupus erythematosus (SLE) patients.

**Introduction:** In spite of high prevalence of dyslipidemia in SLE patients and its role in patients' cardiovascular events, there was scant study about the relation between dyslipidemia and disease activity in SLE patients in Iran.

**Methods:** This analytical cross-sectional study was conducted during 2014–2016 on SLE patients who referred to the Hasheminejad hospital (Tehran – Iran). The serum levels of triglyceride, cholesterol, LDL, and HDL were measured, then dyslipidemia and correlated factors were evaluated. The activity of disease was determined by SLE disease activity index (SLEDAI).

**Results:** 62 out of 72 patients (87%) were female and the mean age was 34 years. The median disease duration was 1 year and 49% of patients had active disease (SLEDAI  $\geq 6$ ). Proteinuria and nephritis were observed in 18% and 24%, respectively. 62% of patients had at-least one abnormality in their lipid profile. High cholesterol ( $>200$  mg/dL), high triglyceride ( $>150$  mg/dL), high LDL ( $>130$  mg/dL) and low HDL ( $<50$  mg/dL in females and  $<40$  mg/dL in males) levels were observed in 25%, 42%, 20% and 49% of patients, respectively. Patients with active disease had lower age and disease duration in comparison of others ( $P < 0.05$ ), while there were no differences in terms of sex and weight between patients in active and inactive phases ( $P > 0.05$ ). The frequency of proteinuria, nephritis and decreased level of complements were higher in active SLE patients, too. Patients with active disease had also higher levels of serum cholesterol, triglyceride and LDL and lower level of serum HDL. In logistic regression, the odds ratios of patients with high cholesterol, using more than 10 mg/day prednisolone and with low serum HDL level for having active disease were 6.6, 5.6 and 3.4, respectively ( $P < 0.05$ ).

**Conclusion:** Our findings showed that dyslipidemia is prevalent in SLE patients especially in patients with active SLE disease. In addition, patients with high cholesterol, using more than 10 mg/day prednisolone and with low HDL had higher chance for having active disease. Hence, it seems that there is a relation between disease activity and lipid profile abnormalities in SLE patients.

<http://dx.doi.org/10.1016/j.pbj.2017.07.010>

Neurosciences Paralell Oral Session

Friday, September 15th, 14h00

### PS168

#### Astrocytic A2A receptors: Novel targets to manage brain disorders



Vanessa Henriques<sup>1,\*</sup>, Nélío Gonçalves<sup>1</sup>, Paula Agostinho<sup>1,2</sup>, Rodrigo A. Cunha<sup>1,2</sup>

<sup>1</sup> CNC – Center for Neuroscience and Cell Biology, University of Coimbra, Portugal

<sup>2</sup> Faculty of Medicine, University of Coimbra, Portugal

E-mail address: [vanessajhenriques@gmail.com](mailto:vanessajhenriques@gmail.com) (V. Henriques).

**Aim:** To validate astrocytic adenosine A2A receptors (A2AR) as a novel target to prevent abnormal glutamate overexcitation.

**Introduction:** Astrocytes are responsible for clearance of extracellular glutamate, a process controlled by A2AR, extolling

their key role as regulators of synaptic transmission and of the abnormal glutamate overexcitation implicated in both acute and chronic brain diseases. We have previously showed that activation of astrocytic A2AR reduce astrocytic glutamate uptake under physiological and pathological conditions,<sup>1–3</sup> and that A2AR are aberrantly up-regulated upon multiple brain insults.<sup>4–6</sup>

**Methods:** We incorporated EGFP reporter either alone or combined with either a small hairpin to down-regulate A2AR (shA2AR) or a control sequence (shCTR) into Mokola Lyssavirus (Mok-G) and Vesicular Stomatitis Virus (VSV-G) lentivectors and tested whether Mok-G-coated lentivirus selectively and efficiently transduced astrocytes in primary culture or in mouse brain through stereotaxic administration of lentivectors into striatum [STR], hippocampus [HIPPP] and prefrontal cortex [PFC] (compared to neurotropic VSV-G-coated lentivirus as controls). Herein, we evaluated viral spreading and cell-type transduction through immunofluorescent colocalization of EGFP with glial (GFAP and vimentin) and neuronal (NeuN) markers.

**Results:** After 25 days post-infection, Mok-G-EGFP transduced 68% of cultured astrocytes (EGFP- and DAPI-positive,  $n=1$ ); 100% of GFAP-positive cells colocalized with EGFP as well as 86% cells expressing Vimentin only and 47% expressing both Vimentin and GFAP. Mok-G shA2AR lentiviruses robustly reduced A2AR immunoreactivity compared to Mok-G shCTR in cultured astrocytes. At 4 weeks post-brain administration, Mok-G-EGFP was expressed mainly in astrocytes (GFAP-positive cells) in both STR and HIPPP, and to a lower extent in the PFC, whereas VSV-G-coated lentivirus colocalized with NeuN marker and not with GFAP in any tested brain areas.

**Conclusion:** These data supports the ability of Mok-G lentivectors to efficiently transduce astrocytes to control A2AR density, paving the way for their application to control pathophysiological processes involving astrocytes.

**Acknowledgements:** Supported by DARPA grant 09-68-ESR-FP-010, FCT grants PTDC/SAU-NSC/122254/2010, UID/NEU/04539/2013, PTDC/NEU-NMC/4154/2014 and PEst-C/SAU/LA0001/2013-2014, QREN grant CENTRO-07-ST24-FEDER-002006, NARSAD and Santa Casa da Misericórdia.

## References

1. Matos M, Augusto E, dos Santos-Rodrigues A. *Glia*. 2012;60:702–16.
2. Matos M, Augusto E, Machado NJ. *J Alzheimer's Dis*. 2012;30:1–13.
3. Matos M, Shen HY, Augusto E. *Biol Psychiatry*. 2015;78:763–74.
4. Cunha GM, Canas PM, Oliveira CR, Cunha RA. *Neuroscience*. 2006;141:1775–81.
5. Rebola N, Porciúncula LO, Lopes LV. *Epilepsia*. 2005;46 Suppl. 5:159–65.
6. Tomiyama M, Kimura T, Maeda T. *Synapse*. 2004;52:218–22.

<http://dx.doi.org/10.1016/j.pbj.2017.07.011>

## PS077

### Adenosine A1 receptor antagonism prevents DSI in hippocampal CA1 pyramidal cells



J. Freire<sup>1,2,\*</sup>, D.M. Rombo<sup>1,2</sup>, A.M. Sebastião<sup>1,2</sup>

<sup>1</sup> Instituto de Farmacologia e Neurociências, Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal

<sup>2</sup> Instituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal  
E-mail address: [joanamorimf@gmail.com](mailto:joanamorimf@gmail.com)

(J. Freire).

**Aim:** How adenosine interfere with a short-term form of neuronal plasticity dependent on endocannabinoid, the depolarization-induced suppression of inhibition (DSI).

**Introduction:** The widely consumed psychoactive drug cannabis, containing cannabinoid compounds, and/or caffeine, with adenosinergic antagonizing proprieties, exert their central actions by affecting cognitive operations such as learning and memory. Indeed, endogenous adenosine and endocannabinoids (eCB) are known to interfere with physiological synaptic plasticity phenomena that represent the neuronal substrate of memory formation.

**Methods:** Whole-cell voltage-clamp recordings ( $V_h = -70$  mV) were performed on hippocampal CA1 pyramidal cells of 3 to 5 weeks-old C57BL/6 mice. Slices (350  $\mu$ m thick) were perfused with artificial cerebrospinal fluid (aCSF) supplemented with glutamate receptor antagonists (CNQX, 25  $\mu$ M and DL-APV, 50  $\mu$ M) to block glutamatergic transmission and isolate GABA-mediated responses. Inhibitory postsynaptic currents (IPSCs) were evoked every 3 s through a stimulation electrode placed in stratum radiatum. The recording electrode was filled with a CsCl-based intracellular solution and DSI was evoked through a 5 s voltage step of +80 mV. The magnitude of DSI was measured 9 s after the depolarizing step and DSI recovery was evaluated between 30 and 60 s after depolarization.

**Results:** When recording eCB-mediated DSI we observed a decrease in electrical-evoked IPSC amplitudes to  $81.0 \pm 5.4\%$  of baseline ( $p < 0.01$ ,  $n = 14$ ) that fully recovered to  $90.2 \pm 5.4\%$  after 30–60 s. The adenosine A1 receptor antagonist, DPCPX (100 nM), prevented DSI, recordings showing a non-significant change in IPSCs amplitude to  $95.1 \pm 12.0\%$  of baseline ( $p = 0.3473$ ,  $n = 10$ ) that was maintained throughout the recovery period ( $87.1 \pm 12.0\%$ ).

**Conclusion:** These results suggest that tonic adenosine A1 receptor activation is necessary for the occurrence of DSI. The mechanisms involved in this process remain unclear and need further investigation.<sup>1–4</sup>

## References

1. Chevaleyre V, et al. *Neuron*. 2004;43:871–81.
2. Chevaleyre V, et al. *Annu Rev Neurosci*. 2006;29:37–76.
3. Sebastião AM, et al. *Brain Res*. 2014;1621:102–13.
4. Rombo DM, et al. *Cereb Cortex*. 2016;26:1081–95.

<http://dx.doi.org/10.1016/j.pbj.2017.07.012>

## PS087

### High-sucrose diet effects on the dendritic trees of developing neurons of the adolescent rat



R. Rodrigues<sup>1,2,\*</sup>, F. Barreto<sup>1,2</sup>, A. Cardoso<sup>1,2</sup>, J.P. Andrade<sup>1,2</sup>

<sup>1</sup> Department of Biomedicine – Unit of Anatomy, Faculty of Medicine, University of Porto, Alameda Prof. Hernâni Monteiro, 4200-319 Porto, Portugal  
<sup>2</sup> Center of Health Technology and Services Research (CINTESIS), Faculty of Medicine, University of Porto, Rua Dr. Plácido da Costa, 4200-450 Porto, Portugal  
E-mail address:

[patriciarafaelarodrigues@gmail.com](mailto:patriciarafaelarodrigues@gmail.com)

(R. Rodrigues).

**Aim:** In the present study, we aimed to explore the effect of high-sucrose diets on the dendritic trees of immature granule cells of the adolescent male rats.

**Introduction:** Adolescence is a period of high susceptibility to exogenous factors as the rat brain is still developing. Evidence shows that high-sucrose diets may be more detrimental to adolescent rats, therefore we intended to study immature granule cells in the hippocampal formation of these animals. For that, we used

doublecortin (DCX), a microtubule-associated protein expressed by neuronal precursor cells and immature neurons, which is used as a marker for neurogenesis.

**Methods:** At 4 weeks of age, adolescent male Wistar rats were randomly allocated to control group ( $n=7$ ) and to an high-sucrose (30% sucrose) diet group ( $n=4$ ; HS) during 4 weeks. After this period, rats were sacrificed and DCX immunocytochemistry was performed. The dendritic trees of the DCX-immunostained cells were drawn with the aid of a camera lucida. A metric analysis of the dendritic trees was performed, and the following parameters were quantified: total dendritic length, the total number of terminal segments, the total number of intermediate segments, mean length of terminal segments and mean length of intermediate segments.

**Results:** Our results show that the total dendritic length of HS adolescent rats was significantly reduced when compared with controls ( $p<0.03$ ). There were no other differences in the others parameters quantified.

**Conclusion:** In conclusion, the dendritic trees of immature neurons of the dentate gyrus of HS adolescent rats appear to be disturbed after the exposition of this diet. This data confirms previous evidence reporting adolescence as a susceptible period of the brain development with likely consequences in cognition. If that is so, and if the reported results can be extrapolated to man, public health interventions are necessary to advise adolescents concerning their diet.

**Acknowledgements:** This article was supported by ERDF through the operation POCI-01-0145-FEDER-007746 funded by the Programa Operacional Competitividade e Internacionalização – COMPETE2020 and by National Funds through FCT - Fundação para a Ciência e a Tecnologia within CINTESIS, R&D Unit (reference UID/IC/4255/2013).

<http://dx.doi.org/10.1016/j.pbj.2017.07.013>

### PS109

#### Looking for modulatory brain areas in the visual circuit related to freezing behaviour

Maria Roa

NERF (Neuro-Electronics Research Flanders),  
Belgium

E-mail address: [meriroa@hotmail.com](mailto:meriroa@hotmail.com).

**Aim:** I have been studying a visual circuit that is known to trigger freezing: the connection between the Retina, the Superior Colliculus and the Parabrachial Grey. The aim of this investigation is to look for cerebral nuclei that could be inputs of the SC and, therefore, regulate this behaviour. In other words, it is a search for modulatory brain areas in the circuit: Retina → SC → PBg.

**Introduction:** Information supplied by the retina initiates interactions in the brain that eventually lead to conscious perception of the visual scene, conventional reflexes such as adjusting the size of the pupil or triggering certain behaviours. Innate defensive behaviours evoked by threatening stimuli are essential to survival. When a danger suddenly appears, a mouse can either escape or freeze. I am interested in how the visual world cause freezing and why.

**Methods:** The tracing strategy used is based on two injections (stereotaxic surgery) with two different retrograde viruses. The first injection is in the PBg with a modified HSV (Herpes Simplex Virus) and the second one, 21 days later, with RVdG (Rabies Virus G-deleted) in the SC. The combined characteristics of these viruses allowed me to specifically follow the circuit. After perfusing the animals, slicing the brains and staining with specific antibodies attached to fluorochromes, I took images with a fluorescent confocal microscope.

**Results:** With a pertinent image processing and comparison with the brain atlas, I was able to identify which brain areas were mostly labelled: zona incerta, substantia nigra and L5 in V1 (visual cortex).

**Conclusion:** It is known that these three nuclei are involved in other visual pathways but this finding suggest that they also could have a role in freezing response to a visual stimulus. The current work is now focused on finding out how each one participates in modulating the behaviour.

**Acknowledgements:** This thesis is going to be evaluated by the University of Barcelona and it is supported by the KU Leuven, the experiments were performed at NERF (Neuro-Electronics Research Flanders) in the Karl Farrow's Laboratory.

<http://dx.doi.org/10.1016/j.pbj.2017.07.014>

### PS178

#### Cafeteria-diet effects on learning and memory, anxiety and fear response of the adolescent rat



André Ferreira<sup>1,2,\*</sup>, João Paulo Castro<sup>1,3</sup>,  
José P. Andrade<sup>1,2</sup>, Armando Cardoso<sup>1,2</sup>

<sup>1</sup> Department of Biomedicine – Unit of Anatomy,  
Faculty of Medicine, University of Porto, Alameda  
Prof. Hermâni Monteiro, 4200-319 Porto, Portugal

<sup>2</sup> Center of Health Technology and Services Research  
(CINTESIS), Faculty of Medicine, University of Porto,  
Rua Dr. Plácido da Costa, 4200-450 Porto, Portugal

<sup>3</sup> Physical Medicine and Rehabilitation Department,  
Centro Hospitalar Vila Nova de Gaia/Espinho, Vila  
Nova de Gaia, Portugal

E-mail address: [andre.ferreira@live.com.pt](mailto:andre.ferreira@live.com.pt)

(A. Ferreira).

**Aim:** We aimed to explore the effect of high caloric diets on adolescent male rats to mimic the feeding behavior of human adolescents in the Western world.

**Introduction:** Age of high-caloric diet exposure is an important factor for the cognitive and anxiety outcomes as key processes of brain development and maturation occur during adolescence. Evidence shows high-caloric diets to affect differently learning and memory performance in an age-dependent way, being more detrimental to adolescent rats.

**Methods:** At 4 weeks of age, 30 adolescent male Wistar rats were randomly allocated to control, high-sugar (HS) and high-fat high-sugar (HFHS) diet groups during 4 weeks. After this period, behavioral tests were performed to study: (1) anxiety behavior in the elevated plus-maze (EPM) and open field tests, (2) learning and memory processes in the Morris water maze (MWM) and novel object recognition test, (3) fear response in fear conditioning tests and (4) depression state in forced swim test.

**Results:** Our results show that only HFHS-treated rats presented more anxiety than control rats, spending more time in the closed arms and less time in open arms of EPM. Moreover, HFHS-treated animals presented an impairment of spatial learning in the final phase of acquisition and an impairment of spatial memory, since these rats spend less time in the target quadrant of MWM and cross less times the former position of the platform. There were no differences between groups regarding locomotor activity, fear acquisition and memory, object novelty detection and exploration, and depression state.

**Conclusion:** In conclusion, anxiety behavior and spatial learning and memory are particularly affected by a cafeteria-type diet in young rats. This data confirms previous evidence reporting adolescence as a susceptible period of brain development to neural insults. Furthermore, the results show that there are different cognitive



and emotional behavioral consequences between HS and HFHS diets.

**Acknowledgements:** This article was supported by ERDF through the operation POCI-01-0145-FEDER-007746 funded by the Programa Operacional Competitividade e Internacionalização – COMPETE2020 and by National Funds through FCT - Fundação para a Ciência e a Tecnologia within CINTESIS, R&D Unit (reference UID/IC/4255/2013).

<http://dx.doi.org/10.1016/j.pbj.2017.07.015>

Oncology & Molecular Biology Paralell Oral Session  
Friday, September 15th, 14h00

## PS081

### GE11 positive exosomes as a potential RNAi delivery system in clear cell Renal Cell Carcinoma

B. Adem<sup>1,\*</sup>, A.L. Teixeira<sup>2,3</sup>, F. Dias<sup>2,3</sup>, C. Ruivo<sup>1</sup>, R. Medeiros<sup>2,3,4,5,\*</sup>, S.A. Melo<sup>1,5,\*</sup>

<sup>1</sup> Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Portugal (i3S), 4200 Porto, Portugal, Institute of Pathology and Molecular Immunology of the University of Porto (IPATIMUP), 4200 Porto, Portugal

<sup>2</sup> Molecular Oncology and Viral Pathology Group, IPO-Porto Research Center (CI-IPOP), Portuguese Institute of Oncology of Porto (IPO-Porto), 4200-072 Porto, Portugal

<sup>3</sup> LPCC, Research Department Portuguese League Against Cancer (Liga Portuguesa Contra o Cancro-Núcleo Regional do Norte), 4200-177 Porto, Portugal

<sup>4</sup> Health Sciences Faculty, Fernando Pessoa University, 4249-004, Porto, Portugal

<sup>5</sup> FMUP, Medical Faculty of the University of Porto, Portugal

E-mail address: [badem@ipatimup.pt](mailto:badem@ipatimup.pt) (B. Adem).

**Aim:** Use GE11 positive (GE11+) exosomes as a targeted delivery system to EGFR overexpressing cells for the treatment of clear cell Renal Cell Carcinoma (ccRCC).

**Introduction:** ccRCC is the most prevalent subtype of renal cancer and the most lethal urologic tumor. Generally, it is radio-chemotherapy resistance, and frequently associated with relapse after 5–11 months upon targeted therapy treatment, which highlight the need to develop new therapeutic strategies. Exosomes, extracellular vesicles of 40–150 nm that mediate intercellular communication, have emerged as promising therapeutic tools due to their engineering potential and ability to evade the immune system.

**Methods:** EGFR is known to be overexpressed in ccRCC thus, the expression of GE11, a peptide that binds to EGFR, in exosomes membrane enable a targeted delivery of therapeutic molecules to EGFR overexpressing cells. Exosomes derived from HEK293T were engineered to express the GE11 peptide on their surface and incubated with normal or tumor renal cell lines.

**Results:** Our results revealed EGFR overexpression at the mRNA and protein levels in a ccRCC cell line, compared to a normal renal cell line. Furthermore, tumor cells presented increased protein levels of phosphorylated EGFR when compared to normal cells. These results support the hypothesis of using an EGFR-based exosomes delivery model, the GE11+ exosomes. A higher percentage of tumor cells internalized GE11+ exosomes compared to exosomes

derived from HEK293T cells transfected with control condition. Additionally, tumor cells exhibited an increased mean of fluorescence intensity compared to the control, suggesting that each cell uptakes more GE11+ exosomes in an EGFR-dependent manner. Importantly, GE11+ exosomes were internalized in a greater proportion by tumor cells rather than normal renal cell lines.

**Conclusion:** Overall, the use of GE11+ exosomes as a new delivery system is a promising therapeutic strategy for ccRCC treatment. Ultimately, these exosomes can be loaded with RNAi-based drugs to target deregulated genes in ccRCC.

<http://dx.doi.org/10.1016/j.pbj.2017.07.016>

## PS145

### Evaluation of combined cytoplasmic AR in tumour cells expression and tumour CD3 T-cells infiltrate as a prognostic score for patients with prostate cancer

V. Constâncio<sup>1,2,\*</sup>, M. McAllister<sup>2</sup>, S. Patek<sup>2</sup>, M. Underwood<sup>3</sup>, H. Leung<sup>4</sup>, J. Edwards<sup>2</sup>

<sup>1</sup> Biology Department, University of Aveiro, Portugal

<sup>2</sup> Institute of Cancer Sciences, Wolfson Wohl Cancer Research Centre, University of Glasgow, United Kingdom

<sup>3</sup> Department of Urology, Queen Elizabeth University Hospital, Glasgow, United Kingdom

<sup>4</sup> Beatson Institute of Cancer Research, United Kingdom

E-mail address: [veraconstancio@ua.pt](mailto:veraconstancio@ua.pt) (V. Constâncio).

**Aim:** We aimed to assess the prognostic value of using a cumulative score evaluating the expression of Androgen Receptor (AR) and the presence tumour inflammatory infiltrate as a prognostic marker for prostate cancer (PCa).

**Introduction:** PCa is the most common male cancer, in Europe. Currently, at diagnosis, only tumour-based factors, including clinical stage, tumour grade and circulating concentrations of Prostate-Specific Antigen (PSA) are used to predict PCa outcome. However, this can vary within patients sharing the same clinical conditions, leading to patient's over/under treatment. It is now recognized that cancer progression is also dependent on tumour's interaction with its microenvironment, specifically with immune cells. Therefore, the development of predictive biomarkers, capable of combining these two factors should be considered.

**Methods:** Immunohistochemistry for AR expression and CD3 T-cells was performed on biopsies from a cohort of 361 patients diagnosed with PCa. Semi-quantitative weighted histoscore and quantitative assessments were used.

**Results:** High cytoplasmic AR expression in tumour cells and high CD3-T cells presence were associated with reduced overall survival ( $p=0.000055$ , and  $p=0.004$ , respectively), with strong association ( $p=0.001$ ) on X2 analysis. When patients were grouped as having: both markers low or one low and low/moderate and one high, and both high, this cumulative prognostic score was strongly associated with overall survival ( $p=0.000001$ ), being the mean overall survivals: 7.1 years (95% CI 6.5–7.6), 6.0 years (95% CI 5.4–6.6) and 3.8 years (95% CI 2.4–5.0), respectively. Moreover, on multivariate analysis, it was considered a significant independent predictor of overall survival (HR 1.982, 95% CI 1.018–3.859,  $p=0.044$ ).

**Conclusion:** These results confirm the clinical utility of assessing both tumour and microenvironment characteristics when predicting patients' outcome, and suggest that the presence



of high cytoplasmic AR expression in tumour cells and CD3 T-cells predicts poor outcome for patients diagnosed with PCa.

<http://dx.doi.org/10.1016/j.pbj.2017.07.017>

## PS165

### ALDHs as potential biomarkers in myeloid neoplasms – Preliminary study

B. Macedo<sup>1,\*</sup>, J. Jorge<sup>2,3</sup>, R. Alves<sup>2,3</sup>,  
A.C. Gonçalves<sup>2,3,4</sup>, A.B. Sarmento-Ribeiro<sup>2,3,4,5</sup>

<sup>1</sup> Department of Chemistry, University of Aveiro, Portugal

<sup>2</sup> Laboratory of Oncobiology and Hematology, University Clinic of Hematology and Applied Molecular Biology, Faculty of Medicine, University of Coimbra, Portugal

<sup>3</sup> Center for Neuroscience and Cell Biology, IBILI, University of Coimbra, Portugal

<sup>4</sup> CIMAGO – Center of Investigation on Environment Genetics and Oncobiology, Faculty of Medicine, University of Coimbra, Portugal

<sup>5</sup> Clinical Hematology Service, University Hospital of Coimbra, Portugal

E-mail address: [barbaramacedo@ua.pt](mailto:barbaramacedo@ua.pt)  
(B. Macedo).

**Aim:** The aim of the study is to evaluate the expression of aldehyde dehydrogenase (ALDH) in patients with myelodysplastic syndromes (MDS) and acute myeloid leukemia (AML) to verify their potential as a marker for the diagnosis and/or prognosis of these diseases.

**Introduction:** ALDH superfamily is a group of 19 enzymes critical to the protection against toxic aldehydes, and have been associated with multiple diseases, namely in cancer. MDS are characterized by ineffective hematopoiesis associated with progressive peripheral blood cytopenias, and a predisposition toward leukemic transformation. MDS pathophysiology is a complex multistep process that involves genetic and epigenetic abnormalities in genes associated with differentiation, cellular proliferation, and apoptosis. Since ALDHs are involved in some of these biological processes, the deregulation of these enzymes may influence MDS and AML development.

**Methods:** To this end, we analyzed the expression levels of 8 ALDH isoforms, ALDH1A1, ALDH1A2, ALDH1B1, ALDH1L1, ALDH1L2, ALDH3A2, ALDH4A1, and ALDH16A1, in 31 patients (16 MDS and 15 LMA) and 19 healthy controls. ALDH expression levels were analyzed using RT-PCR and differentially expressed genes were quantified by qPCR. The statistical analysis was carried out by variance analysis and  $\chi^2$  test. Survival were analyzed by Kaplan Meier curves ( $p < 0.05$ ).

**Results:** Preliminary results indicate that all MDS patients express ALDH16A1 isoform whereas only 67% of controls ( $p < 0.05$ ) show expression of this isoform. Moreover, AML patients have lower ALDH1A2 expression levels than MDS and controls and only 20% of AML patients express this isoform (MDS=54% and controls=55%). The ALDH1L2 is only expressed in chronic myelomonocytic leukemia subtype of MDS. Furthermore, the expression of ALDH isoforms does not appear to influence patient overall survival.

**Conclusion:** According to these results, ALDH isoforms have differential expression patterns in MDS and AML patients when compared with controls and each other. Further studies are needed to prove their potential as a diagnostic/prognostic biomarkers.

<http://dx.doi.org/10.1016/j.pbj.2017.07.018>



## PS155

### Discovery of novel mechanisms of centrosome amplification and their therapeutic value in cancer



B.P. de Almeida<sup>1,2,\*</sup>, G. Marteil<sup>3</sup>,  
M. Bettencourt-Dias<sup>3</sup>, N.L. Barbosa-Morais<sup>2</sup>

<sup>1</sup> Departamento de Ciências Biomédicas e Medicina, Universidade do Algarve, Faro, Portugal

<sup>2</sup> Instituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal

<sup>3</sup> Instituto Gulbenkian de Ciência, Oeiras, Portugal

E-mail address:

[bernardo.almeida@medicina.ulisboa.pt](mailto:bernardo.almeida@medicina.ulisboa.pt)  
(B.P. de Almeida).

**Aim:** To understand the mechanisms of centrosome amplification and their therapeutic value in cancer.

**Introduction:** Centrosomes are the major microtubule-organising centres of animal cells. Centrosome amplification (CA) – the presence of more than two centrosomes in a cell – is a common feature in cancer<sup>1</sup> and was recently shown to be sufficient to drive tumorigenesis.<sup>2</sup> Recent work from the Bettencourt-Dias Lab has identified a new recurrent feature of cancer cells: centriole over-elongation, which also promotes CA. However, origins of those abnormalities and their therapeutic value remain poorly understood.

**Methods:** We have screened the NCI-60 panel of human cancer cell lines<sup>3</sup> for centriole number and individual length to test their frequency and interdependence. We have thereby also generated a metric capturing each abnormality level per cell line that we then correlated with the publicly available molecular (e.g. genomic, transcriptomic and proteomic) and drug-sensitivity quantitative profiles for that panel.

**Results:** Our single-centriole analyses showed that longer centrioles are more common in cells with CA and that cells do not control their overall centriolar mass when the centriole number increases. Moreover, cancer cell lines with longer centrioles proliferate slower due to an accumulation of cells in G1 phase, suggesting that centriole length defects could lead to a cell cycle delay in G1. In addition, our original genome-wide approach highlighted putative mechanisms associated with susceptibility to both abnormalities, such as the proteasome protecting cells from CA. Correlation with drug activity identified some compounds as potential therapeutic options to selectively target cells with higher incidence of centriole abnormalities.

**Conclusion:** This work provides the first single-centriole-level portrait of centriole abnormalities in cancer and contributes to the understanding of their molecular origins, namely by revealing novel molecular mechanisms in cell cycle biology. Given the cancer-specificity of these abnormalities, the identified compounds will inspire the development of drugs to selectively target cancer cells.

**Acknowledgements:** This work is supported by an EMBO Installation Grant to NLBM.

## References

- Chan JY. A clinical overview of centrosome amplification in human cancers. *Int J Biol Sci.* 2011;7:1122–44.
- Levine MS, et al. Centrosome amplification is sufficient to promote spontaneous tumorigenesis in mammals. *Dev Cell.* 2017;40:1–10.
- Shoemaker RH. The NCI60 human tumour cell line anticancer drug screen. *Nat Rev Cancer.* 2006;6:813–23.

<http://dx.doi.org/10.1016/j.pbj.2017.07.019>

## PS164

**Cytotoxic effects of parthenolide on lymphoid malignancies' cell lines**

J. Neves<sup>1,\*</sup>, J. Jorge<sup>2,3</sup>, R. Alves<sup>2,3,4</sup>,  
A.C. Gonçalves<sup>2,3,4</sup>, A.B. Sarmiento-Ribeiro<sup>2,3,4,5</sup>

<sup>1</sup> Department of Life Sciences, Faculty of Science and Technology, University of Coimbra, Portugal

<sup>2</sup> Laboratory of Oncobiology and Hematology (LOH), University Clinic of Hematology and Applied Molecular Biology, Faculty of Medicine, University of Coimbra, Portugal

<sup>3</sup> Center of Investigation on Environment Genetics and Oncobiology (CIMAGO), Faculty of Medicine, University of Coimbra, Portugal

<sup>4</sup> Center for Neuroscience and Cell Biology (CNC), University of Coimbra, Portugal

<sup>5</sup> Clinical Hematology Service, University Hospital of Coimbra, Portugal

E-mail address: [joanafbneves@gmail.com](mailto:joanafbneves@gmail.com)

(J. Neves).

**Aim:** The aim of this study was to evaluate the therapeutic potential of parthenolide (PRT), an NF- $\kappa$ B inhibitor, on acute lymphoblastic leukemia (ALL) and Burkitt Lymphoma (BL) cell lines and characterize the type of cell death induced and its molecular mechanisms.

**Introduction:** Playing an important role in the regulation of diverse biological processes such as cell proliferation and survival, nuclear factor kappa B (NF- $\kappa$ B) is closely associated with various human malignancies. Deregulated NF- $\kappa$ B signaling has been appointed as one important player in all stages of tumorigenesis. PRT has a dual anti-tumor effect – NF- $\kappa$ B pathway inhibition and oxidative stress induction – on a wide range of malignancies and could be a valid option for hematological cancer.

**Methods:** We used one BL (RAJI) and five ALL (697, CEM, JURKAT, MOLT-4 and KOPN8) cell lines. Cells were incubated in absence or presence of different concentrations of PRT in single dose and daily administration. Metabolic activity was assessed by Resazurin Assay. Cell death was analyzed by Optical Microscopy and Flow Cytometry (FC) using Annexin V/7-AAD double staining and JC-1 probe. Apoptotic proteins levels (FAS, FAS-L, BCL-2, BAX and activated caspase 3), cell cycle and oxidative stress parameters (superoxide anion, peroxides and reduced glutathione through the DHE, DCFH2DA and mercury orange probes, respectively) will be evaluated by FC.

**Results:** Preliminary results showed that PRT reduces the metabolic activity in time, dose and cell line dependent manner, being the KOPN8 and RAJI cells the most sensitive and JURKAT cells the lowest. In fact, the half maximal inhibitory concentration (IC<sub>50</sub>) at 72 h was 2  $\mu$ M for KOPN8 and RAJI, 3  $\mu$ M for CEM, 4  $\mu$ M for 697, 6  $\mu$ M for MOLT-4 and 12  $\mu$ M for JURKAT. These results may be related with the cell type and genetic background. Cell death analysis suggested that PRT induced apoptosis in these cell lines. Studies on the cell cycle and oxidative stress are still underway.

**Conclusion:** Our results suggest that PRT is a potential new targeted therapy in lymphoid malignancies, mainly ALL and BL.

<http://dx.doi.org/10.1016/j.pbj.2017.07.020>

## PS169

**WNT/ $\beta$ -catenin and Hedgehog signaling pathways as therapeutic targets in B-cell neoplasms**

C. Ferreira<sup>1,2,\*</sup>, J. Jorge<sup>2,4</sup>, R. Alves<sup>2,3</sup>,  
A.C. Gonçalves<sup>2,3,4</sup>, A.B. Sarmiento-Ribeiro<sup>2,3,4,5</sup>

<sup>1</sup> Department of Chemistry, Biochemical, University of Aveiro, Portugal

<sup>2</sup> Laboratory of Oncobiology and Hematology (LOH), University Clinic of Hematology and Applied Molecular Biology, Faculty of Medicine, University of Coimbra, Portugal

<sup>3</sup> Center for Neuroscience and Cell Biology, IBILI (CNC.IBILI), University of Coimbra, Portugal

<sup>4</sup> CIMAGO - Center of Investigation on Environment Genetics and Oncobiology, Faculty of Medicine, University of Coimbra, Portugal

<sup>5</sup> Clinical Hematology Service, University Hospital of Coimbra, Portugal

E-mail address: [catarina.d.ferreira@ua.pt](mailto:catarina.d.ferreira@ua.pt)

(C. Ferreira).

**Aim:** The goal of this study was to evaluate the therapeutic potential of WNT/ $\beta$ -catenin and Hedgehog inhibitors, IWR-1 and GDC-0449 respectively, alone and in combination, in B-cell neoplasms.

**Introduction:** B-cell neoplasms include, among others, the B-cell lymphomas and plasma cell disorders, such as multiple myeloma (MM), a malignant neoplasm originated by proliferation of monoclonal plasma cells; and diffuse large B-cell lymphoma (DLBCL), the most common form of non-Hodgkin lymphoma. Inappropriate activation of conserved embryonic signaling pathways, such as WNT/ $\beta$ -catenin and Hedgehog has been implicated in B-cell neoplasms. Hence, these pathways may constitute new potential candidate targets for MM and DLBCL therapy.

**Methods:** For this propose, H929 (MM) and FARAGE (DLBCL) cell lines, were cultured in absence and presence of different concentrations of IWR and GDC. Metabolic activity was evaluated using resazurin assay and cell death by optical microscopy (May-Grunwald staining) and flow cytometry (FC) (Annexin V/7-AAD staining). Cell cycle analysis was evaluated by FC, using a PI/RNase solution. Proteins related to apoptosis and some molecules related to WNT and HH signaling pathways were tested by FC. The expression levels of AXIN and SMO genes were analyzed by RT-PCR.

**Results:** Preliminary results showed that IWR-1 and GDC-0449 reduced metabolic activity in a time-, dose- and cell line dependent manner, when administrated alone or in combination. The IC<sub>50</sub> of IWR-1 and GDC-0449 in H929 cells was 40  $\mu$ M and 70  $\mu$ M, respectively, and 75  $\mu$ M and 57  $\mu$ M for FARAGE cell line, after 24 h of treatment. These compounds induce cell death mainly by apoptosis and showed an arrest in cell cycle at G<sub>0</sub>/G<sub>1</sub>. Complementary studies are still ongoing.

**Conclusion:** In conclusion, results suggest that IWR-1 and GDC-0449 are potential new targeted therapies that could be efficient in MM and DLBCL treatment.

<http://dx.doi.org/10.1016/j.pbj.2017.07.021>

Physiology & Immunology Paralell Oral Session  
Friday, September 15th, 14h00

## PS099

### Differences in aerobic capacity and spirometric parameters between athletes and nonathletes



V. Kostić

Department of Physiology, Faculty of Medicine,  
University of Novi Sad, Serbia  
E-mail address: [kostasm91@gmail.com](mailto:kostasm91@gmail.com).

**Aim:** To investigate if there are differences in aerobic capacity and spirometric parameters between athletes and nonathletes, and also differences in these parameters between anaerobic and aerobic athletes.

**Introduction:** Physical fitness is defined as ability of organism to increase level of metabolic processes due to increased level of metabolic needs. Aerobic capacity is measured by maximum level of oxygen consumption (VO<sub>2</sub>max), and it can be expressed by absolute (l/min) or relative (ml/kg/min) value. Pulmonary capacity has great evaluation importance for sport and health of general population.

**Methods:** Number of participants was 45 males, aged 18–35 years, divided into 2 groups: athletes and nonathletes. Athletes were divided by sport type in aerobic and anaerobic group of athletes. Testing was consisted of anthropometric measuring, spirometry and measuring of aerobic capacity on ergobicycle with mask, by principle of ramp test.

**Results:** Value of VO<sub>2</sub>max in group of athletes (55.46 ml/kg/min,  $p < 0.05$ ) was significantly greater than in group of nonathletes (37.78 ml/kg/min,  $p < 0.05$ ). Compared between all groups, VO<sub>2</sub>max showed significant difference in both aerobic (58.88 ml/kg/min,  $p < 0.05$ ) and anaerobic (52.04 ml/kg/min,  $p < 0.05$ ) athletes in relation to nonathletes (38.78 ml/kg/min,  $p < 0.05$ ). Spirometric parameters (FVC, FEV<sub>1</sub>) were significantly greater in group of nonathletes (5.481 L, 4.951 L,  $p < 0.05$ ) than in group of athletes (4.874 L, 4.635 L,  $p < 0.05$ ). Compared between all groups, we found significant difference in FVC between group of nonathletes (5.481 L,  $p < 0.05$ ) and anaerobic athletes (4.807 L,  $p < 0.05$ ), and in Tiffeneau index between group of anaerobic athletes (97.29%,  $p < 0.05$ ) and nonathletes (90.82%,  $p < 0.05$ ).

**Conclusion:** Values of anthropometric parameters are greater in group of nonathletes. Differences in body weight and body mass caused greater values of FVC and FEV<sub>1</sub> in group of nonathletes. Values of aerobic capacity are increasing with training. The greatest values of aerobic capacity are shown by aerobic athletes.

**Acknowledgements:** To Department of Physiology, Faculty of Medicine, Novi Sad for using their resources; To Athletic Club “Vojvodina”, Novi Sad and Triathlon Club “Tryogy” for participating; To Assist. Proff. Aleksandar Klačnja for mentorship.<sup>1–15</sup>

## References

1. Bowers RW, Fox EL, Foss ML. In: Bowers RW, editor. The physiological basis of physical education and athletics. 3rd ed. Boston: Saunders College Publishing; 1988.
2. Whyte G, Spurway N, MacLaren D. In: Whyte G, editor. The physiology of training. Churchill Livingstone; 2006.
3. Ranković G, Mutavdžić V, Toskić D, Preljević A, Kocić M, Nedin-Ranković G, et al. Aerobic capacity as an indicator in different kinds of sports. *Bosn J Basic Med Sci.* 2010;10:44–8.
4. Shin YS, Yang SM, Kim MY, Lee LK, Byoung-Sun Park LWD, Noh JW, et al. Analysis of the respirogram phase of Korean wrestling athletes compared with nonathletes for sports physiotherapy research. *Phys Ther Sci.* 2016;28:392–8.
5. Ponorac N, Matavulj A, Grujić N, Rajkovača Z, Kovačević P. Maximal oxygen uptake (VO<sub>2</sub> max) as the indicator of physicalworking capacity in sportsmen. *Acta Med Med.* 2005;44:17–20.

6. Pelemiš V, Mitrović N, Cicović B, Lolić D. Maximal oxygen consumption for different groups of athletes. *Sportske nauke i zdravlje.* 2011;1:52–7.
7. Albouaini K, Egred M, Alahmar A, Wright DJ. Cardiopulmonary exercise testing and its application. *Postgrad Med J.* 2010;83:675–82.
8. de Jong F. Lung function testing feature: spirometers. *Breathe.* 2008;4:251–4.
9. Myriantsefs P, Grammatopoulou I, Katsoulas T, Baltopoulos G. Spirometry may underestimate airway obstruction in professional Greek athletes. *Clin Respir J.* 2014;8:240–7.
10. Wilmore JH, Costill DL. *Physiology of sport and exercise.* 2nd ed. USA: Human Kinetics; 1999.
11. Verstappen F, Huppertz R, Snoeckx L. Effect of training specificity on maximal treadmill and bicycle ergometer exercise. *Int J Sport Med.* 1982;3:43–6.
12. Klisuras V. *Fundamentals of sports physiology belgrade.* Institut Za Sport; 2013.
13. Kausar A, Mudassir S, Badaam KM, Shete A, Khan S. Cardiorespiratory fitness of university volleyball players and sedentary youngpeople in marathwada region of Maharashtra Province in India. *J Clin Diagn Res.* 2015;9:20–1.
14. Herdy AH, Caixeta A. Brazilian cardiorespiratory fitness classification based on maximum oxygen consumption. *Arq Bras Cardiol.* 2016;106:389–95.
15. Franchini E, Vecchio FBD, Matsushigue KA, Artioli GG. Physiological profiles of Elite Judo athletes. *Sports Med.* 2011;41:147–66.

<http://dx.doi.org/10.1016/j.pbj.2017.07.022>

## PS070

### The assessment of body composition, energy demands and muscle strength in people on different dietary regimes



Dj. Milicev<sup>1,\*</sup>, M. Bogdan<sup>2</sup>, A. Rakovac<sup>1,2</sup>,  
V. Karan<sup>1,2</sup>, A. Klačnja<sup>1,2</sup>, M. Drapšin<sup>1,2</sup>

<sup>1</sup> Faculty of Medicine, University of Novi Sad, Serbia

<sup>2</sup> Department of Physiology, Faculty of Medicine,  
University of Novi Sad, Serbia

E-mail address: [milicev.dj@gmail.com](mailto:milicev.dj@gmail.com)

(Dj. Milicev).

**Aim:** The aim of this study was to determine whether there are any differences in body composition, energy demands and muscle strength between people on different dietary patterns.

**Introduction:** There are numerous types of diets: vegan, vegetarian, and non-vegetarian. Considering the dietary pattern, the assessment of the body composition and determining the resting metabolic rate are a major challenge for many researchers. Regarding the muscle strength of physically inactive participants related to dietary patterns, there is no current data in literature.

**Methods:** The study was conducted at the Department of Physiology, Faculty of Medicine University of Novi Sad from November 2016 to February 2017. The study included 45 healthy, physically inactive randomly selected respondents (15 vegans, 15 vegetarians, 15 on a mixed diet) aged 20–30 years. All respondents practiced their dietary regime for at least 6 months before research. Firstly, the anthropometric measurements were done, and later the body composition was assessed using bioelectrical impedance and by measuring skin folds. The resting metabolic rate was estimated using the indirect calorimetric method. The muscle strength was determined using the isoaccelerating dynamometer.

**Results:** The values of body mass index (BMI) between the group on a mixed diet ( $23.9 \pm 2.95 \text{ kg/m}^2$ ) and vegans ( $20.8 \pm 2.58 \text{ kg/m}^2$ ) showed a statistically significant difference ( $p < 0.05$ ). The BMI ( $21.3 \pm 2.63 \text{ kg/m}^2$ ) for vegetarians did not differ from the other groups. Statistically significant differences between groups in other parameters of body composition, resting metabolic rate and muscle strength were not found. A negative correlation was observed between total body fat, resting metabolic rate and muscle strength in all groups.

**Conclusion:** Diet differences between tested groups affected only the value of BMI between vegans and non-vegetarians. The impacts of different diets on other parameters of body composition,

resting metabolic rate and muscle strength were not confirmed by this study.

<http://dx.doi.org/10.1016/j.pbj.2017.07.023>

## PS146

### Titin phosphorylation by protein kinase G as a novel mechanism of diastolic adaptation to acute load



R. Rocha<sup>1,\*</sup>, J. Almeida-Coelho<sup>1</sup>, A.M. Leite-Moreira<sup>1</sup>, J.S. Neves<sup>1,2</sup>, N. Hamdani<sup>3</sup>, I. Falcão-Pires<sup>1</sup>, A.P. Lourenço<sup>1,4</sup>, W.J. Paulus<sup>5</sup>, W.A. Linke<sup>3</sup>, A.F. Leite-Moreira<sup>1,6</sup>

<sup>1</sup> Department of Physiology and Cardiothoracic Surgery & Cardiovascular Research Center, Faculty of Medicine, University of Porto, Portugal

<sup>2</sup> Department of Endocrinology, São João Hospital Center, Porto, Portugal

<sup>3</sup> Department of Cardiovascular Physiology, Ruhr University Bochum, Germany

<sup>4</sup> Department of Physiology, Institute for Cardiovascular Research, VU University Medical Center, Amsterdam, The Netherlands

<sup>5</sup> Department of Anesthesiology, São João Hospital Center, Porto, Portugal

<sup>6</sup> Department of Cardiothoracic Surgery, São João Hospital Center, Porto, Portugal

E-mail address: [rafaelm.rocha@hotmail.com](mailto:rafaelm.rocha@hotmail.com) (R. Rocha).

**Aim:** To evaluate acute adaptations of myocardial stiffness to acute stretch and characterize the underlying mechanisms.

**Introduction:** Systolic adaptation to myocardial stretch/volume overload is known, but whether the heart is also able to modulate its stiffness following such challenges remains unknown.

**Methods:** Left ventricle (LV) of intact rat Langendorff hearts, rabbit papillary muscles and myocardial strips from cardiac surgery patients were acutely stretched. Skinned cardiomyocytes from Stretched and Non-stretched myocardium were studied. Stretch by increased venous return or volume loading was assessed by echocardiography in healthy volunteers; pressure-volume hemodynamics in cardiac surgery patients and in a rat model of LV hypertrophy. Myocardial cGMP, phosphorylated vasodilator-stimulated phosphoprotein (VASP) and titin phosphorylation were quantified. Pharmacological studies assessed the role of NO and natriuretic peptides (NP).

**Results:** After stretch, end-diastolic pressure (EDP) or passive tension (PT) decreased over 15 min in all preparations. Skinned cardiomyocytes from Stretched hearts showed decreased PT – abrogated by protein phosphatase incubation – those from Non-stretched hearts showed decreased PT after protein kinase (PKG) incubation. Stretched samples showed increased cGMP levels and phosphorylation of VASP. Titin phosphorylation was increased in Stretched samples – attenuated by PKG inhibition (PKG<sub>i</sub>). PT decay after stretch was blunted by PKG<sub>i</sub> or by joint NP antagonism, NO synthase inhibition and NO scavenging. Moreover, it was remarkably attenuated in hypertrophic rat hearts which showed reduced titin phosphorylation and no increase with stretch. Healthy volunteers and cardiac surgery patients showed E/E' and EDP decrease after sustained stretch maneuvers, respectively.

**Conclusion:** We describe a novel physiological mechanism whereby myocardial compliance is increased in response to

stretch/volume overload, by titin phosphorylation through cGMP-PKG signaling. The mechanism was translated to human physiology and may be abolished in the hypertrophic heart (potential role in the pathophysiology of heart failure with preserved ejection fraction).

<http://dx.doi.org/10.1016/j.pbj.2017.07.024>

## PS172

### Cannabis sativa tetrahydrocannabinol (THC) impact on placental endocrine function



L. Midão<sup>1,3,\*</sup>, J. Maia<sup>1,2</sup>, M. Almada<sup>1,2</sup>, B. Fonseca<sup>1,2</sup>, D. Gonçalves<sup>4</sup>, J. Braga<sup>4</sup>, N. Teixeira<sup>1,2</sup>, G. Correia-da-Silva<sup>1,2</sup>

<sup>1</sup> Laboratório de Bioquímica, Faculdade de Farmácia Universidade do Porto, Porto, Portugal

<sup>2</sup> UCIBIO-REQUIMTE, Porto, Portugal

<sup>3</sup> Departamento de Química, Universidade de Aveiro, Aveiro, Portugal

<sup>4</sup> Departamento da Mulher e da Medicina Reprodutiva, Serviço de Obstetrícia, Centro Materno-Infantil do Norte- Centro Hospitalar do Porto, Porto, Portugal

E-mail address: [midao@ua.pt](mailto:midao@ua.pt) (L. Midão).

**Aim:** The main goal of this work is to understand the impact of THC on placenta endocrine function.

**Introduction:** Cannabis sativa-based medicines have been used to help ease pain, nausea and loss of appetite in cancer and HIV patients. Endocannabinoid system plays an important role in the regulation of female fertility and pregnancy. This system is implicated in proliferation, differentiation and apoptosis of placental stem cells, the trophoblasts (1). These mediate critical steps such as hormone production, fetal immune protection and increase in maternal vascular blood flow. Previous studies have shown that cannabis consumption during pregnancy is associated with intrauterine growth restriction, preterm labor and low birth weight. Moreover, tetrahydrocannabinol (THC), the main psychoactive compound of cannabis, is able to cross the placental barrier. However its effect on trophoblasts turnover and hormone production are unknown.

**Methods:** Term placental explants were treated with THC [1–40 μM] for 24 h to 72 h. The relative mRNA levels of 3β-HSD, aromatase, leptin and PP13 were determined by qRT-PCR. The protein expression levels of 3β-HSD, aromatase and leptin were assessed by Western Blot. Progesterone, estradiol and β-human chorionic gonadotropin (β-hCG) levels were measured by ELFA.

**Results:** After 24 h, PP13 mRNA levels were significantly increased at 40 μM of THC, while for leptin this effect was observed at 10 μM. Moreover, after 72 h aromatase mRNA levels were increased, while there was no effect on 3β-HSD. No differences were observed regarding progesterone whilst, an increase in estradiol and β-hCG with 40 μM at 72 h was detected.

**Conclusion:** These findings suggest that THC may impair trophoblast turnover and endocrine function which may affect pregnancy outcome. Moreover these results may contribute to disclose the cellular effects of cannabis-derived drugs.

**Acknowledgements:** Work financed by FEDER through COMPETE and FCT through PTDC/DTP-FTO/5651/2014-POCI-01-0145-FEDER-016562; FCT/MEC and FEDER, under PT2020 (UID/01/0145/FERDER/007728) and CCDR-N/NORTE2020/Portugal 2020 (norte-01-0145-FEDER-000024).<sup>1</sup>

## Reference

- Costa MA, Fonseca BM, Teixeira NA, Correia-da-Silva G. The endocannabinoid anandamide induces apoptosis in cytotrophoblast cells: involvement of both mitochondrial and death receptor pathways. *Placenta*. 2015;36:69–76.

<http://dx.doi.org/10.1016/j.pbj.2017.07.025>

## PS219

### Reactivity of the rat distal colon to autoantibodies targeting angiotensin type I receptors

R. Magalhães<sup>1,\*</sup>, A. Philippe<sup>2</sup>, R. Catar<sup>3</sup>,  
D. Dragun<sup>3</sup>, M. Morato<sup>4</sup>

<sup>1</sup> *Laboratory of Pharmacology, Department of Drug Sciences, Faculty of Pharmacy of University of Porto, Portugal*

<sup>2</sup> *Department of Nephrology and Critical Care Medicine, Charité University Medicine, Berlin, Germany*

<sup>3</sup> *Department of Nephrology and Critical Care Medicine, Charité University Medicine, Berlin, Germany; Berlin Institute of Health, Berlin, Germany*

<sup>4</sup> *Laboratory of Pharmacology, Department of Drug Sciences, Faculty of Pharmacy of University of Porto, Portugal; MedInUP - Center for Drug Discovery and Innovative Medicines, University of Porto, Porto, Portugal*

E-mail address: [rtmglhs@gmail.com](mailto:rtmglhs@gmail.com)

(R. Magalhães).

**Aim:** To describe the reactivity of the rat distal colon to AT1R-Abs and to compare it to that of Ang II.

**Introduction:** Agonistic IgG (IgG1 and IgG3 subclasses) autoantibodies against the angiotensin II type 1 receptor (AT1R-Abs) have been associated with hypertension, preeclampsia, placental ischemia, renal-allograft rejection and systemic sclerosis. It is thought that AT1R-Abs mimic the action of angiotensin II (Ang II) and contribute to the physiopathology of several diseases and the associated complications.

**Methods:** Male Wistar rats (9–12 weeks of age) were killed by decapitation and strips of the distal colon were mounted in organ baths along their longitudinal axis. Tissues were stretch to 1 g of resting force and isometric responses to AT1R-Abs (25, 50 and 100 mg/dl) obtained from sera of systemic sclerosis and renal-allograft rejection patients and to Ang II (10pM–1 μM) were recorded on a polygraph. The response of Ang II were expressed as % of the response to 125 mM potassium chloride (KCl).

**Results:** AT1R-Abs caused a long-lasting response. Very often, AT1R-Abs induced an increase in the frequency and amplitude of distal colon spontaneous contractions. Occasionally, AT1R-Abs caused a slight decrease in the resting tone and, more rarely, they caused colonic contraction. The effects of the AT1R-Abs seem to be attenuated by candesartan. The pattern of the response to Ang II was different; Ang II caused a fast developing contraction of the colon with an Emax of  $64.37 \pm 12.68$  (%KCl) and EC50 of  $1.22 \pm 0.20$  nM.

**Conclusion:** AT1R-Abs change the normal rhythm of spontaneous contractions of the rat distal colon but more studies are necessary to evaluate whether this reactivity is mediated by AT1 receptors. Moreover, Ang II cause a marked AT1 receptor-mediated contraction of the rat distal colon.

**Acknowledgements:** The authors acknowledge Mrs. Céu Pereira and Mrs. Mónica Caldas for excellent technical assistance.

<http://dx.doi.org/10.1016/j.pbj.2017.07.026>



Public Health & Medical Informatics Paralell Oral Session  
Friday, September 15th, 14h00

## PS224

### Intestinal colonization by antibiotic-resistant Gram negatives in children



C.S. Cruz<sup>1,\*</sup>, R. Mota<sup>1</sup>, D. Gonçalves<sup>1,2,3</sup>,  
H. Ferreira<sup>1,2</sup>

<sup>1</sup> *Microbiology, Department of Biological Sciences, Faculty of Pharmacy, University of Porto, Portugal*

<sup>2</sup> *UCIBIO, University of Porto, Portugal*

<sup>3</sup> *Superior Institute of Health of Alto Ave, Portugal*  
E-mail address: [cruz.carolinasantos@gmail.com](mailto:cruz.carolinasantos@gmail.com)  
(C.S. Cruz).

**Aim:** This study aims to further the knowledge of antibiotic-resistance in the commensal intestinal flora of children by studying the intestinal colonization by antibiotic-resistant Gram negative bacteria in portuguese children.

**Introduction:** Although it is known resistance to antibiotics is a growing problem worldwide, this scenario which constitutes a risk factor for infectious disease is an under-characterized reality in Portugal.

**Methods:** Faecal samples of 29 healthy children (4 months to 12 years-old) were collected from randomly selected localities of Portugal: Viana do Castelo ( $n=8$ ), Porto ( $n=6$ ), Braga ( $n=14$ ), Leiria ( $n=1$ ), from September 2016 to March 2017. Risk factors were assessed by questionnaire, namely antibiotic usage history and direct contact with dependent elders. Isolates were selected by spreading saline suspension (100 μL) on MacConkey agar and MacConkey agar with ampicillin (100 μg/mL), cefotaxime (2 μg/mL), and meropenem (1 μg/mL). Susceptibility profiles to β-lactam and non-β-lactam antibiotics were assessed by disk-diffusion methods according to the EUCAST. Presumptive identification of the isolates was performed with CHROMagar–Orientation culture media.

**Results:** In a total of 29 isolates (lactose fermenters ( $n=22$ ) and lactose non-fermenters ( $n=8$ )), 28 showed resistance to amoxicillin and 13 to amoxicillin with clavulanic acid. Of the 29 children analysed, 17 showed resistance to at least one of the antibiotics studied. Four children were colonized with bacteria resistant to cephalosporins ( $n=8$ ), two of which have daily contact with elders.

**Conclusion:** The results indicate that young children might be an important reservoir of commensals with clinically relevant resistance mechanisms. The clarification of this reality in Portugal could prove essential in the fight against silent dissemination of these threats and persistent infections.

<http://dx.doi.org/10.1016/j.pbj.2017.07.027>

## PS187

### Is the oral mycobiome of young adults influenced by the delivery mode?



P. Campos<sup>1</sup>, L. Costa<sup>1,\*</sup>, M. Ferreira<sup>1</sup>,  
C. Fernandes<sup>1</sup>, S. Ferreira<sup>1</sup>, I. Moreira<sup>1</sup>,  
R. Moreira<sup>1</sup>, M. Pereira<sup>2</sup>, B. Sampaio-Maia<sup>3</sup>

<sup>1</sup> *Faculty of Dentistry, University of Porto, Portugal*

<sup>2</sup> *ISPUP-EPIUnit – University of Porto, Portugal*

<sup>3</sup> *IS – Instituto de Investigação e Inovação em Saúde, University of Porto, Portugal*  
E-mail address: [analmcosta@hotmail.com](mailto:analmcosta@hotmail.com)  
(L. Costa).

**Aim:** To investigate whether the mode of delivery influences the oral yeast colonization in young adults.

**Introduction:** The human microbiome is a complex ecosystem that varies considerably across the body and between individuals.<sup>1</sup> Postnatally the child is exposed to microorganisms from maternal and environmental sources and influenced by infant feeding, developing its own microbiome that will continue evolving throughout life.<sup>2</sup> Several studies have been carried out to determine the influence of the mode of delivery on the oral microbiome, and some influence on bacterial colonization has been verified.<sup>3,4</sup> However, the influence on oral fungal colonization is still unknown.

**Methods:** In 200 healthy students from the Faculty of Dentistry of University of Porto, colonization by yeast in the oral cavity was evaluated by collecting unstimulated saliva. Yeast isolation was performed by pour-plaque technique using Sabouraud Agar medium supplemented with chloramphenicol and ChromAgar Candida medium for species identification. Statistical analysis was performed using the chi-square test and t-test for independent samples.

**Results:** Participants' mean age was  $21.61 \pm 1.86$  years old, with a total yeast prevalence of 37.5%. *Candida albicans* was the most isolated species present in 76.5% of the colonized participants. In comparison to caesarean section, the participants born by normal delivery presented higher oral yeast prevalence (41.6% vs. 25.8%,  $p=0.035$ ) and higher oral yeast load ( $13.68 \pm 38.02$  vs.  $1.69 \pm 0.62$  Log CFU/mL,  $p=0.030$ ).

**Conclusion:** Our results suggest that delivery mode influences the oral mycobiome throughout life, specifically, normal delivery appears to promote the oral yeast colonization.

## References

- Ding T, Schloss PD. Dynamics and associations of microbial community types across the human body. *Nature*. 2014;509:357–60.
- Sampaio-Maia B, Caldas IM, Pereira ML, Perez-Mongiovi D, Araujo R. The oral microbiome in health and its implication in oral and systemic diseases. *Adv Appl Microbiol*. 2016;97:171–210.
- Lif Holgerson P, Harnevik L, Hernell O, Tanner AC, Johansson I. Mode of birth delivery affects oral microbiota in infants. *J Dent Res*. 2011;90:1183–8.
- Ubeja RG, Bhat C. Mode of delivery and its influence on the acquisition of streptococcus mutans in infants. *Int J Clin Pediatr Dent*. 2016;9:326–9.

<http://dx.doi.org/10.1016/j.pbj.2017.07.028>

PS034

## Why, how and when are patients with Chromosomal anomalies hospitalized?



Manuel Gonçalves-Pinho<sup>1,2,\*</sup>,  
João Vasco Santos<sup>1,2</sup>, Sílvia Fernández<sup>1</sup>,  
Micaela Gregório<sup>1</sup>, Carla Pinto Moura<sup>3,4</sup>,  
Alberto Freitas<sup>1,2</sup>

<sup>1</sup> Department of Community Medicine, Information and Health Decision Sciences (MEDCIDS), Faculty of Medicine, University of Porto, Rua Dr. Plácido da Costa, s/n, 4200-450 Porto, Portugal

<sup>2</sup> Center for Health Technology and Services Research (CINTESIS), Rua Dr. Plácido da Costa, s/n, 4200-450 Porto, Portugal

<sup>3</sup> Department of Human Genetics, Faculty of Medicine, University of Porto/Centro Hospitalar São João, Porto, Portugal

<sup>4</sup> Institute for Research and Innovation in Health/Instituto de Investigação e Inovação em Saúde, University of Porto, Porto, Portugal  
E-mail address: [manuelpinho19@gmail.com](mailto:manuelpinho19@gmail.com)  
(M. Gonçalves-Pinho).

**Aim:** We aim to describe Chromosomal anomalies (CA) related hospitalizations characteristics and specific trends in

order to understand why, how and when are these patients hospitalized.

**Introduction:** CA affect approximately 2% of the world population.<sup>1</sup> Due to this low prevalence not many studies regarding hospitalizations are available in this set of conditions. Hospitalizations represent an overall health and prognosis indicator that may allow the implementation of specific health care policies regarding prevention measures to avoid CA-related hospitalizations.

**Methods:** A retrospective observational study was performed using a national hospitalization database that gathers all public hospital admissions between 2000 and 2014. CA were selected based on codes 758.0× to 758.7× codified by the International Classification of Diseases – 9th Revision – Clinical Modification. Birth date, sex, charges, admission/discharge date, discharge status, primary/secondary diagnoses were analyzed for each specific CA.

**Results:** CA related hospitalizations accounted for 0.08% of all the hospitalizations. Down syndrome represented 75.9% of all CA-related hospitalizations and 80.2% (approximately 30M€) of all the charges attributed to CA related hospitalizations. The median age of CA-related patients was 9.0 years old. The leading causes of hospitalization in different CA varied between pneumonia (3.6–18.6%) and live birth related diagnoses (7.9–52.5%). Mean number of hospitalizations ranged from 1.0 to 2.1 per patient and mean charges per hospitalization varied from 2 339 to 4 520 €.

**Conclusion:** CA hospitalizations have high mean charges per hospitalization, high length of stay and high in-hospital mortality. Down syndrome accounts for the majority of CA hospitalizations, representing the CA with higher economic burden in the health system. Klinefelter syndrome hospitalizations occur at a younger age than the described mean age of diagnoses in all Klinefelter syndrome patients, a novel finding not previously described.

**Acknowledgements:** We thank ACSS for providing the data on hospitalizations registered on public hospitals. Fernando Lopes, MD, for his support in the design of the study and João Paulo Oliveira, MD PhD, for his valuable insight regarding genetic epidemiology. We also thank project “NORTE-01-0145-FEDER-000016” (NanoSTIMA) that is financed by the North Portugal Regional Operational Programme (NORTE 2020), under the PORTUGAL 2020 Partnership Agreement, and through the European Regional Development Fund (ERDF).

## Reference

- Rimoin DL, et al. Nature and frequency of genetic disease. In: Emery and Rimoin's principles and practice of medical genetics; 2002. p. 55–9.

<http://dx.doi.org/10.1016/j.pbj.2017.07.029>

PS195

## Efficiency of web application and spaced repetition algorithms as an aid in preparing to practical examination of histology



Dominik Karch<sup>1,\*</sup>, Krzysztofa Kopyt<sup>1</sup>,  
Aleksandra Gauden<sup>1</sup>, Michal Nowakowski<sup>2</sup>

<sup>1</sup> Student Research Group – Jagiellonian University Medical College, Poland

<sup>2</sup> Jagiellonian University Medical College, Department of Medical Education, Poland  
E-mail address: [dexterdk@gmail.com](mailto:dexterdk@gmail.com)  
(D. Karch).

**Aim:** The aim of this study is to evaluate impact of using web application on the results of histology practical exam as well as to check if the SuperMemo-based algorithm is a useful tool in medical education.

**Introduction:** Students in medical disciplines are looking for new learning strategies. Computer applications are becoming more popular as they use a variety of methods to improve efficiency of studying. One of them is spaced repetition algorithm like Super-Memo.

**Methods:** We prepared web application which shows the photography of histological slide. Students had to decide if they have recognized the slide and the program was measuring time of each answer. Then the algorithm allocated new slide to display.

Users were randomly divided into two groups: study – where difficult slides were shown more frequently (SuperMemo2-based algorithm) and control – where the slides were displayed randomly.

Quality of the student's answers was evaluated according to the 6-point scale, where 0 means incorrect answer, and from 1 to 5 – correct answer depending on time.

We also took into consideration results of histology practical exam (0–15 points).

The level of statistical significance was set at  $p < 0.05$ .

**Results:** The study involved 204 first year medical students. The study group ( $n=98$ ) and control ( $n=106$ ) were similar in terms of the average number of responses in application (901 vs. 858;  $p=0.73$ ).

We have shown a statistically significant difference which indicate obtaining higher examination score by students who used our application – 11.8 vs. 10.98 ( $p=0.016$ ).

There was no superiority of spaced repetition algorithm over the random allocation of slides, based on the examination results (11.7 vs. 11.9;  $p=0.73$ ).

**Conclusion:** The usage of computer programs can be a valuable complement to traditional teaching methods. As we showed in this study it may have a measurable effect on examinations results of the students.

**Acknowledgements:** The approval of the Jagiellonian University Bioethics Committee was obtained.

The authors thank Prof. J.A. Litwin, Head of the Department of Histology Jagiellonian University Medical College for substantive supervision over the study.

<http://dx.doi.org/10.1016/j.pbj.2017.07.030>

## PS008

### The frequency of Human Parvovirus B19 infections in Vojvodina



M. Bugarski, T.A. Aleksandra Patić

*Department for Microbiology with Parasitology and Immunology, Faculty of Medicine, University of Novi Sad, Serbia*

*E-mail address: [marinabugarski@gmail.com](mailto:marinabugarski@gmail.com)*

(M. Bugarski).

**Aim:** Determinating the seroprevalence of IgG antibodies among residents of Vojvodina, as well as the incidence of acute infections in different age groups and with different diagnoses, especially in women of generative age and pregnant women.

**Introduction:** Human Parvovirus B19 is a cause of infections in patients of all age groups. Clinical manifestations vary from asymptomatic to manifest infections such as erythema infectiosum, arthropathy, heart problems, and infections in immunodeficient patients. Acute infections during pregnancy present a distinct problem, which can result in intrauterine fetal death or hydrops fetalis.

**Methods:** The data presented in this study are the result of serological testing for the presence of HP-B19 infections performed at the Institute of Public Health of Vojvodina, Centre for Virology, in the period from November 2015 to November 2016. Detection of specific IgG and IgM antibodies was completed by analysing 472

serum samples. Samples were tested using the ELISA test manufactured by VIRION, Germany, in the VIRION Analyzer I-2P device.

**Results:** Of the total number of tested subjects, an acute infection was detected in 10.8% of the cases (11.7% of pregnant women, and 7.14% of children). An acute infection was confirmed in 13.9% of the patients in a febrile state, and 7.1% of the patients diagnosed with arthritis, immune deficiency, and heart failure. Seroprevalence of IgG antibodies was confirmed in 42.8% of the tested subjects, 36.8% of pregnant women, 60.78% of non-pregnant women of generative age, and 11.03% of children. In the total sample, 46.4% of the results were negative.

**Conclusion:** It can be concluded that Human Parvovirus B19 exist and circulates in the population of Vojvodina. The use of rapid serological tests enables a specific etiological diagnosis, timely implementation of appropriate infection control measures, and an appropriate treatment of patients, especially those belonging to high risk groups like pregnant women are.

<http://dx.doi.org/10.1016/j.pbj.2017.07.031>

## PS211

### E-cigarette: An effective tool to quit smoking or an additional source of nicotine?



Miłosz Knura\*, Tomasz Kurowski, Jakub Lubański, Paulina Majek, Mateusz Jankowski

*Department of Epidemiology, Medical University of Silesia in Katowice, Poland*

*E-mail address: [miłosz.knura@wp.pl](mailto:miłosz.knura@wp.pl)*

(M. Knura).

**Aim:** We sought to evaluate the effectiveness of e-cigarette use as a tool to quit or reduce smoking.

**Introduction:** The electronic cigarettes (known as an “e-cigarettes”) gaining on popularity, especially among young people. Available evidence regarding the relationship between e-cigarette usage as a tool in smoking cessation are inconsistent.

**Methods:** A population based survey was performed, in a group of 3800 students from three Universities in Katowice, Poland. Self-prepared, previously validated questionnaire, included questions on e-cigarette smoking habits.

**Results:** Completed questionnaires were obtained from 3000 students (response rate 78.9%; mean age =  $21.5 \pm 2.1$  yrs) of which 70% were female (F) and 30% were male (M). E-smoking was declared by 3.5% of respondents (F: 3%, M: 4.9%;  $p=0.01$ ), wherein 1.5% of respondents smoked only e-cigarettes (F: 1.3%; M: 1.8%;  $p=0.3$ ) and 2.4% of subjects were dual smokers (F: 1.6%; M: 3%;  $p=0.01$ ). Almost one-third (33.7%) of e-smokers used e-cigarettes as an aid to quit smoking. Only 13.8% of e-smokers tired to give up e-smoking. Almost half of e-smokers (48.8%) tends to give up e-smoking in the nearest future. Reduction in cigarette consumption (mean  $6.5 \pm 5.0$  cigarettes/daily) was observed by 50.8% of dual smokers. Only 4.4% of e-smokers used e-cigarettes without nicotine. Since they started e-smoking, constant concentration of nicotine in e-liquid was indicated by 61.4% of e-smokers, 12.5% increased (mean  $8.7 \pm 5.1$  mg/ml) and 26.1% reduced (mean  $8.2 \pm 3.5$  mg/ml) nicotine content in usually used e-liquid. Among e-smokers, 48.8% reported an addiction to e-cigarettes.

**Conclusion:** Smoking cessation was not the main reason for e-cigarette use among most of e-smokers. Low percent of e-smokers who use a non-nicotine e-liquid and almost half of e-smokers who declared addiction to e-cigarette, suggests that e-cigarette is rather an additional source of nicotine than effective tool in smoking cessation.

<http://dx.doi.org/10.1016/j.pbj.2017.07.032>

Surgery Paralell Oral Session  
Friday, September 15th, 14h00

### PS130

#### The effect of medication intake on perforation rate in patients with colonic diverticulosis – A retrospective assessment



Aneta Turza\*, Justyna Tęczar, Rafał Król,  
Anna Gadamer

Students' Scientific Group at 2nd Department of  
General Surgery, Jagiellonian University Medical  
College, Poland

E-mail address: [aturza@onet.pl](mailto:aturza@onet.pl) (A. Turza).

**Aim:** The aim was to study the effect of drug intake on the frequency of perforation among patients with colonic diverticulosis.

**Introduction:** Diverticulosis is a common condition which incidence increases with age. One of the most severe complications with a high risk of late sequel and mortality is perforation of colonic diverticula. According to the current studies the use of some medications may affect the risk of perforation due to its influence on the intracolonic pressure and mucosal barrier function.

**Methods:** A retrospective review of 294 patients (mean age 68.6) with verified colonic diverticulosis was done. Included patients were admitted to 2nd Department of General Surgery JU MC during 2006–2016. Study enrolled 206 (70.1%) women and 88 (29.9%) men. Among investigated group 36 (12.2%) patients developed perforation.

The research regarded medications including NSAIDs, corticosteroids, calcium-channel blockers, statins, opioids, aspirin, anticoagulants and antiplatelet drugs. In addition, the data concerning comorbidity and the severity of the diverticulitis was collected.

**Results:** In the analysis the group of patients with perforation and the group with non-perforated diverticulosis were compared. Higher rates of the use of NSAIDs (13.89% vs. 3.88%; OR = 4;  $p = 0.01$ ; 95% CI = 1.28–12.46), opioids (11.11% vs. 1.55%; OR = 7.94;  $p < 0.001$ ; 95% CI = 1.89–33.3) and corticosteroids (22.22% vs. 8.14%; OR = 3.22;  $p = 0.01$ ; 95% CI = 1.31–7.96) were observed among the patients with perforation. The results revealed an inverse relation concerning the use of statins (5.56% vs. 22.09%; OR = 0.21;  $p = 0.02$ ; 95% CI = 0.05–0.89). Similar results were found in the review of available literature.

**Conclusion:** Medications used by patients with colonic diverticulosis affect the incidence of perforation. The administration of NSAIDs, corticosteroids and opioids is related to an increased rate of diverticular perforation. Conversely, statins may contribute to the decrease in the frequency of perforation. We can conclude that it is important to carefully administer drugs to patients with colonic diverticulosis.

<http://dx.doi.org/10.1016/j.pbj.2017.07.033>

### PS025

#### De novo atrial fibrillation following aortic valve replacement surgery is associated with decreased creatinine clearance and increased C-reactive protein levels



Mariana Fragão-Marques<sup>1</sup>, Francisca Saraiva<sup>1</sup>,  
João Oliveira<sup>1,\*</sup>, André P. Lourenço<sup>1</sup>,  
Adelino Leite-Moreira<sup>1,2</sup>, Inês Falcão-Pires<sup>1</sup>

<sup>1</sup> Departamento de Cirurgia e Fisiologia da Faculdade de Medicina da Universidade do Porto, Alameda Professor Hernâni Monteiro, Porto, Portugal

<sup>2</sup> Departamento de Cirurgia Cardiotorácica do Centro Hospitalar de São João, Porto, Alameda Professor Hernâni Monteiro, Porto, Portugal  
E-mail address: [oliveira.p.joao96@gmail.com](mailto:oliveira.p.joao96@gmail.com) (J. Oliveira).

**Aim:** We aim to assess the prevalence of de novo postoperative atrial fibrillation (POAF) in patients submitted to aortic valve replacement surgery (AVRS) and evaluate clinical and echocardiographic variables as predictors of POAF occurrence in this population.

**Introduction:** POAF is the most common complication following cardiac surgery, with peak incidence in the second day after the surgical procedure. Studies have demonstrated an increase in the incidence of stroke, hospital stay, health-associated costs and mortality in the group of patients experiencing POAF.

**Methods:** We conducted a cross-sectional study, that included all the patients submitted to AVRS during 2014 in a tertiary hospital, diagnosed with severe aortic valve stenosis without endocarditis, known history of atrial fibrillation, more than one major procedure, or other significant valve disease. Data were collected retrospectively and the statistical tests were performed according to the variable classification.

**Results:** The incidence of POAF in the 173 included patients was 45%, with the median time of occurrence being  $2.4 \pm 1.5$  days. A univariate analysis showed that the group of patients who developed POAF was older ( $p = 0.028$ ), had longer median in-hospital stay ( $p = 0.008$ ), had a significantly higher C-reactive protein (CRP) peak blood level ( $p = 0.025$ ) and a significantly lower minimum creatinine clearance ( $p = 0.026$ ) in the post-operative period when compared with those who did not develop POAF. A multivariate analysis confirmed age to be an independent predictor of POAF. (OR: 1.04, CI 95%: 1.00–1.09).

**Conclusion:** Our study suggests age, peak post-operative blood level of CRP and creatinine clearance as predictors of POAF occurrence and supports the hypothesis that POAF may be the result of inflammation, being one of the few studies that focuses on a population with isolated aortic stenosis. Our findings on increased hospital stay reinforce the idea of risk stratification and the use preventive measures in the higher risk groups.

<http://dx.doi.org/10.1016/j.pbj.2017.07.034>

### PS050

#### Predictors of early reoperation after meningioma removal



Paulina Donicz\*, Kornelia Kliś,  
Małgorzata Gackowska

Student Scientific Group at Department of  
Neurosurgery and Neurotraumatology, Jagiellonian  
University Medical College, Poland

E-mail address: [paulina.donicz@gmail.com](mailto:paulina.donicz@gmail.com)  
(P. Donicz).

**Aim:** The aim of our study was to establish predictors of unplanned early reoperations after meningioma removal.

**Introduction:** Complications after neurosurgical procedures which lead to reoperation are associated with poor treatment outcome and costs. The knowledge of risk factors for complications might allow to implement specific preventive measures. However those factor are still poorly defined, especially in terms of benign brain tumours.

**Methods:** We retrospectively analysed 177 patients, with histologically confirmed meningiomas, hospitalized between 2014 and 2016 who underwent craniotomy. From medical records

we obtained detailed medical history (previous diseases, medications, tumour characteristics, blood test results, surgery's details). Completeness of tumour resection was assessed using Simpson Grade. Early reoperation was defined as reoperation during the same hospital stay. We used  $\chi^2$  test for proportional values; *t*-student test, Mann–Whitney *U* test for continuous variables. To determine the potential predictors of early reoperation we used univariate and multivariate logistic regression analyses.

**Results:** A total of 13 (7.34%) patients underwent unplanned early reoperation. Those patients significantly more often had retromastoid craniotomy (25.00% vs. 6.40%;  $p=0.047$ ). And significantly more often suffered from ischemic heart disease (66.67% vs. 6.64%;  $p<0.01$ ) and atrial fibrillation (60% vs. 6.25%;  $p<0.01$ ). Reoperated patients also more often took heparin (50% vs. 6.74%;  $p<0.01$ ) and anticoagulants (66.67% vs. 6.21%;  $p<0.01$ ). In multivariate logistic regression analysis anticoagulants intake (OR: 31.463; 95% CI: 1.139–868.604;  $p=0.04$ ) and retromastoid craniotomy (OR: 6.642; 95% CI: 1.139–38.73;  $p=0.034$ ) remained independently associated with higher risk of early reoperation.

**Conclusion:** Patients who underwent retromastoid craniotomy, those with history of ischemic heart disease or atrial fibrillation and those who take heparin and anticoagulants are more likely to require early reoperation. Retromastoid craniotomy and anticoagulants intake are independent risk factors for early reoperation.

**Acknowledgements:** To our Tutors: Jarosław Polak, MD, PhD and Roger Krzyżewski, MD.

<http://dx.doi.org/10.1016/j.pbj.2017.07.035>

#### PS095

##### Prevalence of foramen arcuale and its clinical significance: A meta-analysis of 55,985 subjects

Przemysław A. Pękala<sup>1,2</sup>,  
Brandon Michael Henry<sup>1,2</sup>, Jakub R. Pękala<sup>1,2,\*</sup>,  
Wan Chin Hsieh<sup>1,3</sup>, Jens Vikse<sup>1,2</sup>,  
Beatrice Sanna<sup>4</sup>, Jerzy A. Walocha<sup>1,2</sup>,  
R. Shane Tubbs<sup>5</sup>, Krzysztof A. Tomaszewski<sup>1,2</sup>

<sup>1</sup> International Evidence-Based Anatomy Working Group, Krakow, Poland

<sup>2</sup> The Brain and Spine Lab, Department of Anatomy, Jagiellonian University Medical College, Krakow, Poland

<sup>3</sup> First Faculty of Medicine, Charles University, Prague, Czech Republic

<sup>4</sup> Faculty of Medicine and Surgery, University of Cagliari, Sardinia, Italy

<sup>5</sup> Seattle Science Foundation, Seattle, Washington, USA

E-mail address: [jr.pekala@gmail.com](mailto:jr.pekala@gmail.com)  
(J.R. Pękala).

**Aim:** The aim of this study was to deliver the most complex study on the prevalence of the FA and its clinical significance.

**Introduction:** Foramen arcuale (FA) is an osseous prominence formed in place of a sulcus for the vertebral artery on the posterior arch of the atlas. The presence of an FA can make a threat during neurosurgery by giving a false notion of a wider posterior arch when viewed dorsally during C1 lateral mass screw insertion.

**Methods:** An comprehensive search of the major electronic databases was performed in order to find and identify all studies which reported relevant data on the FA. No date or language restrictions were applied. Data on the prevalence, type (complete

and incomplete), side, gender, laterality, and morphometrics of the FA were extracted and pooled into a meta-analysis.

**Results:** A total of 127 studies ( $n=55,985$  subjects) were included into the quantitative analysis. The overall pooled prevalence of a complete FA was 9.1% (95%CI: 8.2–10.1), while the overall pooled prevalence of an incomplete FA was 13.6% (95%CI: 11.2–16.2). The complete FA was found to be most prevalent in North American (11.3%) and European (11.2%) populations, and least prevalent among Asian (7.5%) populations, especially Chinese (4.4%) and South Koreans (5.8%). In the presence of a complete FA, a contralateral FA (complete or incomplete) was found in 53.1% of cases.

**Conclusion:** The FA is a commonly present anatomical structure. Awareness of a complete variant of the FA during procedures performed on the atlas vertebra is essential in reducing the risk of iatrogenic injury. Therefore, risk for the presence of an FA should be considered by surgeons prior to procedures on the atlas in each patient according to gender and ethnic group. As such, we highly advise preoperative screening with CT as the gold standard for finding the presence of an FA.

<http://dx.doi.org/10.1016/j.pbj.2017.07.036>

#### PS136

##### Quality of Life and aortobifemoral bypass – Importance of the hypogastric arteries



C. Braz\*, R. Castro-Ferreira, P. Dias, S. Sampaio, J. Teixeira

Faculdade de Medicina da Universidade do Porto, Portugal

E-mail address: [carolina.c.braz@gmail.com](mailto:carolina.c.braz@gmail.com)

(C. Braz).

**Aim:** Evaluate SD after AFB and assess the importance of patent hypogastric arteries before the procedure.

**Introduction:** The aortobifemoral bypass (AFB) is one of the best options to revascularize patients with Aortoiliac Occlusive Disease (AIOD). The impact of this procedure in sexual function (SF) is unpredictable, with 20–80% of the patients reporting sexual dysfunction (SD) after surgery. There's still insufficient data to safely predict the development of SD after AFB and what the role of hypogastric arteries.

**Methods:** The study includes only male population submitted to AFB due to AIOD. Patients with major amputations after the surgery were excluded. The development of SD was evaluated by phone call. The quality of life before and after the procedure was evaluated by a standardized index score questionnaire (15D). Pre-operative patency of hypogastric arteries was appraised by assessing the patients imaging file. The arteries with direct anterograde flow were considered patent.

**Results:** Of a total of 53 patients, 40 were included in the study – exclusion causes were intrahospital death, natural cause death and major amputation. In the included group, 37% reported worsening, 26% improved and 37% did not notice any change in SF after surgery. If at least one of the hypogastric arteries was patent before surgery, 58% described worsening in SF compared to only 13% in the group with no sustained anterograde flow to the hypogastric arteries. 92% of the patients was not warned of the possibility of SD after surgery, being that 26% of these would have refused the procedure if they knew.

**Conclusion:** SD is a prevalent and often overlooked complication after open aortoiliac revascularization and it remains a major taboo in the surgeon/patient relation. The existence of at least one hypogastric artery with reserved anterograde flow before

surgery can strongly predict a higher risk of sexual dysfunction after surgery.

<http://dx.doi.org/10.1016/j.pbj.2017.07.037>

## PS197

### Compensatory renal overgrowth after unilateral nephrectomy in children



Krzysztofa Kopyt<sup>1,\*</sup>, Aleksandra Gauden<sup>1</sup>,  
Michal Ebisz<sup>1</sup>, Michal Jurczyk<sup>1</sup>, Adam Olesiak<sup>1</sup>,  
Piotr Soltysiak<sup>2</sup>, Wojciech Gorecki<sup>2</sup>

<sup>1</sup> Student Research Group – Jagiellonian University  
Medical College, Krakow, Poland

<sup>2</sup> Pediatric Surgery Clinic, Jagiellonian University,  
Krakow, Poland

E-mail address: [krzysztofa.kopyt@student.uj.edu.pl](mailto:krzysztofa.kopyt@student.uj.edu.pl)  
(K. Kopyt).

**Aim:** The aim of the study is to investigate the intensity of renal overgrowth after unilateral nephrectomy in children's population, as well as to check dependency between kidney's dimensions and patient's age.

**Introduction:** Solitary kidney after unilateral nephrectomy tends to overgrow. In adult population the dynamic of overgrowth and maximal dimensions are identified. In childhood there are no described patterns of the process of solitary kidney overgrowth.

**Methods:** Patients who had undergone unilateral nephrectomy in the University Children's Hospital of Cracow were enrolled. The length of the solitary kidney was compared with control group which was based on ultrasound examination of the kidney (left  $n = 1601$ , right  $n = 1635$ ) performed in the same clinic in children without kidney disease. All examinations were carried out with Philips Epiq 5G ultrasound unit with convex probe C5-1 MHz by a single physician (PS).

The comparison was analysed with t-student test for one or two means. 18 children (7 males) from the birth to the age of 17 who underwent in total 48 ultrasound examination after the nephrectomy were enrolled.

**Results:** There was significant difference between the mean of the kidney's length in patients after unilateral nephrectomy and control group. The difference was the most explicit in the groups at the age from 8 to 13 for the right kidney (difference range from 13 to 22 mm,  $p < 0.05$ ) and in the groups at the age from 8 to 12 for the left kidney (difference range from 11 to 19 mm,  $p < 0.05$ ). Solitary kidney in children after nephrectomy is significantly larger than in the control group.

**Conclusion:** The dynamic of solitary kidney overgrowth in children should be taken into consideration while performing the sonographic examination. Chronic kidney disease may be suspected when overgrowth of the solitary kidney is not present. Further research dealing with the dynamic of compensatory kidney overgrowth in children is indicated.

<http://dx.doi.org/10.1016/j.pbj.2017.07.038>

Internal Medicine Plenary Session  
Saturday, September 16th, 14h00

## PS067

### Cardiac effects of Ledipasvir plus sofosbuvir for Hepatitis C treatment in thalassemia



H. Karimi-Sari<sup>1,2,3,4,\*</sup>, A. Khosravi<sup>1,2,3,4</sup>,  
B. Behnava<sup>1,2,3,4</sup>, M. Abedi-Andani<sup>1,2,3,4</sup>,  
S.M. Alavian<sup>1,2,3,4</sup>

<sup>1</sup> Student Research Committee, Baqiyatallah  
University of Medical Sciences, Tehran, Iran

<sup>2</sup> Baqiyatallah Research Center for Gastroenterology  
and Liver Diseases (BRCL), Baqiyatallah University  
of Medical Sciences, Tehran, Iran

<sup>3</sup> Middle East Liver Diseases (MELD) Center, Tehran,  
Iran

<sup>4</sup> Atherosclerosis Research Center, Baqiyatallah  
University of Medical Sciences, Tehran, Iran

E-mail address: [dr.karimih@yahoo.com](mailto:dr.karimih@yahoo.com)  
(H. Karimi-Sari).

**Aim:** This study was designed to evaluate the effects of Ledipasvir plus Sofosbuvir on cardiac function of thalassemia patients.

**Introduction:** Hepatitis C (HCV) infection is much more prevalent in thalassemia patients because of blood transfusion. Thalassemia patients may also have cardiac abnormalities due to congenital problems, anemia, and increased burden of iron in their myocardium. HCV treatment has been revolutionized after introducing new direct acting antiviral (DAA) drugs, and data is limited about effects of these new drugs on patients' cardiac function.

**Methods:** In this study HCV-infected thalassemia patients which were selected for treatment with DAAs in HepCC-2 trial (NCT03061032) were evaluated prospectively. Fixed dose daily tablets of 90 mg-Ledipasvir plus 400 mg-Sofosbuvir (12/24week,  $\pm$ ribavirin) was prescribed for patients. All patients were evaluated by a unique echocardiography fellowship for collecting the echocardiography findings of before and after the treatment. Then effects of mentioned drugs on patients' cardiac function were evaluated.

**Results:** Thirty-two patients with mean age of  $24.2 \pm 6.4$  years were evaluated. The treatment response, which was evaluated by rapid virological response and sustained virological response rates, was 100%. The patients' left ventricular end-systolic diameter (LVESD) and volume (LVESV), global longitudinal strain (GLS) of LV and average, and right ventricle (RV) size were significantly increased after finishing the treatment ( $P < 0.05$ ). Changes in abovementioned parameters were not correlated with patients' myocardium iron load ( $P > 0.05$ ). There were no significant differences in before-after comparison of other echocardiographic parameters ( $P > 0.05$ ).

**Conclusion:** Ledipasvir-Sofosbuvir combination therapy was safe for our HCV-infected thalassemia patients and cause no serious cardiac events. But minimal changes in strain, size, and volume of left ventricle, and size of right ventricle may refer to needing more precise cardiac evaluations in these patients. Also, our patients' ejection fraction remained unchanged. Hence, we suggest more specific and long-term echocardiographic evaluations before and after treatment, if needed.<sup>1-3</sup>

## References

1. Jafroodi M, Davoudi-Kiakalayeh A, Mohtasham-Amiri Z, Pourfathollah AA, Haghbin A. Trend in prevalence of Hepatitis C virus infection among B-thalassemia major patients: 10 years of experience in Iran. *Int J Prev Med.* 2015;6.

- Poller W, Kaya Z, Muche M, Kasner M, Skurk C, Kappert K, et al. High incidence of cardiac dysfunction and response to antiviral treatment in patients with chronic Hepatitis C virus infection. *Clin. Res. Cardiol.* 2017:1–6.
- Hagiwara S, Nishida N, Watanabe T, Sakurai T, Ida H, Minami Y, et al. Outcome of combination therapy with sofosbuvir and ledipasvir for chronic type C liver disease. *Oncology.* 2017;92 Suppl. 1:3–9.

<http://dx.doi.org/10.1016/j.pbj.2017.07.039>

Neurosciences Plenary Session  
Saturday, September 16th, 14h00

### PS141

#### Paclitaxel-induced neuropathic pain: Unravelling the underlying mechanisms at the central nervous system

J. Ribeiro<sup>1,2,3,\*</sup>, J.T. Costa-Pereira<sup>1,2,3</sup>,  
I. Tavares<sup>1,2,3</sup>, I. Martins<sup>1,2,3</sup>

<sup>1</sup> Departamento de Biomedicina – Unidade de Biologia Experimental, Faculdade de Medicina da Universidade do Porto, Portugal

<sup>2</sup> i3S – Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Portugal

<sup>3</sup> IBMC – Instituto de Biologia Celular e Molecular, Universidade do Porto, Portugal

E-mail address: [jribeiro504@gmail.com](mailto:jribeiro504@gmail.com)

(J. Ribeiro).

**Aim:** Here we studied the effects of the cytostatic paclitaxel on: (i) the development of nociceptive and aversive behaviors; (ii) noxious-evoked-activation of spinal dorsal horn neurons and (iii) on descending noradrenergic modulation, which is the main spinal nociceptive inhibitory system.

**Introduction:** Chemotherapeutic drugs are widely used for cancer treatment but they also cause numerous deleterious side effects. Chemotherapy-induced neuropathy (CIN) is one of the most common side effects. The mechanisms underlying CIN are starting to be uncovered namely the alterations induced by cytostatics at the peripheral nervous system but the effects of these drugs at the central nervous system are still poorly studied.

**Methods:** Male Wistar rats were injected with paclitaxel (Taxol, 2.0 mg/kg) or the vehicle solution dimethyl sulfoxide on four alternate days. Nociceptive and aversive behaviors were assessed by the von Frey and conditioned place aversion (CPA) tests, respectively. Noxious-evoked-activation of spinal dorsal neurons was achieved at one month after CIN by evaluating the expression of c-fos expression upon cold stimulation. To study the descending noradrenergic pain modulation we assessed the effects of the  $\alpha$ 2-adrenoreceptor agonist clonidine at 1 and 10  $\mu$ g administered intrathecally, on the von Frey test. We further assessed the expression of the  $\alpha$ 2-adrenoreceptor and dopamine- $\beta$ -hydroxylase (DBH), a nor-adrenaline biosynthetic enzyme expressed in noradrenergic fibers, at the spinal dorsal horn.

**Results:** Paclitaxel induced mechanical allodynia and aversive behaviors. c-fos and DBH expression were increased in paclitaxel-treated animals while  $\alpha$ 2-adrenoreceptor expression remained unaltered. Clonidine induced antinociception at both doses with more pronounced effects in paclitaxel-treated animals.

**Conclusion:** Paclitaxel-treated animals showed neuropathic like-behaviors and increased spinal neuronal activation. It remains to ascertain if DHB upregulation results in increased spinal nor-adrenaline levels, but the increase of  $\alpha$ 2-AR antinociceptive potency in paclitaxel-treated animals indicates the recruitment of

descending inhibition probably as a buffer to increased spinal sensitization.

**Acknowledgements:** Funding: Norte 2020 refs: NORTE-01-0145-FEDER-000008.

<http://dx.doi.org/10.1016/j.pbj.2017.07.040>

Oncology & Molecular Biology Plenary Session  
Saturday, September 16th, 14h00

### PS203

#### Is there horizontal transfer of the oncogene BCR-ABL mediated by extracellular vesicles released by chronic myeloid leukemia cells?

A. Teixeira<sup>1,2,3,\*</sup>, D. Sousa<sup>1,2,4</sup>, C.P.R. Xavier<sup>1,2</sup>,  
M.H. Vasconcelos<sup>1,2,4</sup>

<sup>1</sup> i3S – Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Porto, Portugal

<sup>2</sup> Cancer Drug Resistance Group, IPATIMUP – Instituto de Patologia e Imunologia Molecular da Universidade do Porto, Porto, Portugal

<sup>3</sup> ICBAS – Instituto de Ciências Biomédicas Abel Salazar, Porto, Portugal

<sup>4</sup> Department of Biological Sciences, FFUP – Faculty of Pharmacy of the University of Porto, Porto, Portugal

E-mail address: [alexandrat@ipatimup.pt](mailto:alexandrat@ipatimup.pt)

(A. Teixeira).

**Aim:** The aims are to verify if: (i) EVs released by CML cells carry BCR-ABL in their cargo and if that BCR-ABL is captured by recipient cells; (ii) EVs released by a CML drug resistant cell line, with mutant BCR-ABL, may transfer mutant BCR-ABL and a resistant phenotype to sensitive cells.

**Introduction:** BCR-ABL, the fusion gene originated by the t(9;22) translocation, is responsible for Chronic Myeloid Leukemia (CML). BCR-ABL codes for a constitutively active tyrosine kinase (TK), deregulating downstream pathways and promoting cell survival. Imatinib mesylate (Gleevec), a TK inhibitor, is the gold standard treatment for CML; nevertheless, resistance to this drug often arises, mostly caused by additional point mutations on BCR-ABL and representing a major clinical drawback. It was recently suggested that drug resistance might be horizontally transferred by EVs, from resistant to sensitive cells.

**Methods:** A pair of drug-sensitive BCR-ABL+ cell line (KBM5), and its drug-resistant counterpart (KBM5-STI, harboring mutated BCR-ABL) were used in this study. EVs were isolated by ultracentrifugation and characterized by Dynamic Light Scattering, Nanoparticle Tracking Analysis, Transmission Electron Microscopy and Western Blot. The resazurin assay was used to assess drug response of drug resistant cells, drug sensitive cells and of drug sensitive cells following co-culture with EVs released by drug resistant cells. BCR-ABL levels were analysed by Western Blot.

**Results:** A dose-response curve to imatinib was performed in both cell lines, to confirm their different responses to the drug. Regarding EVs characterization, they had between 10 and 1000 nm and presented several markers of EVs with no evidence of cellular contaminants. Interestingly, BCR-ABL protein was detected in the EVs.<sup>1–9</sup>

**Conclusion:** These results suggest that there is selective packaging of BCR-ABL into EVs, promoting oncogenic protein shedding. Ongoing work will clarify if the EVs released by the resistant cells have mutant BCR-ABL and if they confer drug resistance to recipient sensitive cells.



## References

1. Goldman J, et al. *N Engl J Med.* 2003;349:1451–64.
2. Krause DS, et al. *N Engl J Med.* 2005;353:172–87.
3. Weisberg E, et al. *Nat Rev Cancer.* 2007;7:345–56.
4. Chávez-González A, et al. In: Alimoghaddam K, editor. *Stem Cell Biology in Normal Life and Diseases.* InTech; 2013 [Chapter 8].
5. Raimondo S, et al. *Cell Commun Signal.* 2015;13:8.
6. Sousa D, et al. *Trends Mol Med.* 2015;21:595–608.
7. Raposo G, et al. *J Cell Biol.* 2013;200:373–83.
8. Akers JC, et al. *J Neuro-Oncol.* 2013;113:1–11.
9. Théry C, et al. *Curr Protocols Cell Biol.* 2006:3–22.

<http://dx.doi.org/10.1016/j.pbj.2017.07.041>

Physiology & Immunology Plenary Session  
Saturday, September 16th, 14h00

### PS030

#### Assessment of left ventricular systolic and diastolic function in diabetic rat model using Electrocardiography-gated 18F-FDG PET imaging



Eissler Christoph<sup>1,2,3,\*</sup>, Nobuyuki Hayakawa<sup>3</sup>, Paula-Anahi Arias-Loza<sup>2</sup>, Hiroshi Wakabayashi<sup>3</sup>, Rudolf Werner<sup>1,3</sup>, Tetsuya Shinaji<sup>1,3</sup>, Constantin Lapa<sup>3</sup>, Theo Pelzer<sup>2</sup>, Takahiro Higuchi<sup>1,3</sup>

<sup>1</sup> Comprehensive Heart Failure Center, University Hospital Wuerzburg, Wuerzburg, Germany

<sup>2</sup> Internal Medicine I University Hospital Wuerzburg, Wuerzburg, Germany

<sup>3</sup> Nuclear Medicine University Hospital Wuerzburg, Wuerzburg, Germany

E-mail address: [christopheissler@web.de](mailto:christopheissler@web.de)  
(E. Christoph).

**Aim:** In this study, we explore the potential of ECG-gated 18F-FDG PET to assess LV systolic and diastolic function in a well-established rat model of type 2 diabetes.

**Introduction:** Left ventricular (LV) diastolic dysfunction, defined as a disruption of the normal filling pattern of the ventricle but normal systolic function, is one of the early signs of cardiac involvement in diabetic patients.

**Methods:** List-mode gated 18F-FDG PET imaging was performed on a rat model of type 2 diabetes (ZDF fa/fa) ( $n=6$ ) and ZL control rats ( $n=6$ ) at age of 13 weeks 15–30 min after tracer-administration (37 MBq) via tail vein under hyperinsulinemic-euglycemic clamp using a dedicated small animal PET system (Siemens Inveon) with ECG signal recording for 20 min. List-mode data were sorted and reconstructed into tomographic images of 16 frames per cardiac cycle. PET images were resized to match human-scale pixels. Left ventricular functional parameters were calculated using standard clinical software program (Heart Function View).

**Results:** Hyperinsulinemic-euglycemic clamp and post mortem tissue analysis demonstrated the development of diabetes in the ZDF rats and of significant myocardial hypertrophy in ZDF rats at age of 13 weeks ( $994 \pm 78$  mg vs.  $871 \pm 44$  mg in ZDF rats vs. ZL controls,  $p < 0.01$ , respectively). The PET images analysis showed a mild but significant decrease of LV PFR in the ZDF rats ( $10.4 \pm 0.5$  vs.  $11.8 \pm 0.4$  EDV/s in ZDF rats vs. ZL controls,  $p < 0.001$ , respectively), whereas no significant differences concerning LVEF and cardiac output (CO) could be detected between model and control rats (LVEF:  $60.0 \pm 4.5$  vs.  $63.7 \pm 4.1\%$ ,  $p = 0.25$  and CO:  $90917 \pm 14015$  vs.  $85208 \pm 17511$   $\mu$ l/min,  $p = 0.90$ , respectively).

**Conclusion:** In a rat model of type 2 diabetes, we demonstrated the ability of ECG-gated-18F-FDG PET together with a clinical ventricular edge detection software to assess reliable LV systolic and diastolic parameters and to detect the presence of a diastolic dysfunction in the diabetic rats.

<http://dx.doi.org/10.1016/j.pbj.2017.07.042>

Public Health & Medical Informatics Plenary Session  
Saturday, September 16th, 14h00

### PS045

#### Prevalence of dietary supplements and over-the-counter drug use in patients with arterial hypertension



Mateusz Łobacz\*, Marek Stopa, Magdalena Niemczyk, Karolina Rutkowska, Agata Radko

Jagiellonian University Medical College – Students' Scientific Group at the First Department of Cardiology, Interventional Electrocardiology and Hypertension, Poland

E-mail address: [lobacz.mateusz@gmail.com](mailto:lobacz.mateusz@gmail.com)  
(M. Łobacz).

**Aim:** Analysis of frequency of use of DS/OTC among patients with arterial hypertension as well as factors determining its use and patients' knowledge about possible interactions with conventional medication.

**Introduction:** Dietary supplements (DS) and over-the-counter drugs (OTC) are frequently advertised as a natural treatment of many disorders. DS/OTC can interfere with biotherapeutic action of prescribed medication and this is of particular concern in patients with cardiovascular disease, many of whom are on long term treatment.

**Methods:** The study was conducted in the Outpatient Hypertensive Clinic in the Tertiary Cardiac Center. Self-prepared questionnaire was administered among 151 hypertensive patients (58% females, age range 18–80 years). Regular DS/OTC use was defined as taking them at least 3 times per week.

**Results:** In the examined population regular use of DS/OTC was declared by 67% subjects. The most commonly, regularly used substances were minerals and microelements (60.4%), vitamins (48.5%), analgesics (18.8%), drugs increasing the immunity (18.8%), relieving the gastrointestinal symptoms (18.8%) and omega acids (18.8%). There were no differences in the frequency of DS/OTC use in relation to number of antihypertensive drugs, educational level, age and income. Women are more frequent regular users of DS/OTC than men ( $n = 65$  vs.  $n = 36$ ,  $p = 0.03$ ). Only 38% of responders always consulted the use of DS/OTC with a doctor. The majority of responders (52%) is not aware of possible influence of DS/OTC on antihypertensive medication or blood pressure control. Cost of DS/OTC in 23% of responders is equal or higher than cost of prescribed drugs.

**Conclusion:** Two thirds of hypertensive patients are regularly using DS/OTC. Half of them are not aware of possible interactions with antihypertensive therapy and influence of blood pressure control. The perception that nonprescription therapies are unnecessary to report during medication history taking should be changed. DS/OTC are the important position in the responders budget.

<http://dx.doi.org/10.1016/j.pbj.2017.07.043>

Surgery Plenary Session  
Saturday, September 16th, 14h00

### PS125

#### Do patients after bariatric surgery change their physical activity habits? A prospective one-year follow-up study



K. Jasińska\*, A. Wałkiewicz, D. Bugara

Students' Scientific Group at 2nd Department of General Surgery, Jagiellonian University Medical College, Poland

E-mail address: [katarzyna.jasinska100@gmail.com](mailto:katarzyna.jasinska100@gmail.com)

(K. Jasińska).

**Aim:** The purpose of this study was to assess whether patients have changed their physical activity habits one year after the bariatric surgery.

**Introduction:** Vast majority of obese patients have developed their condition by overeating and insufficient physical activity. Severe obesity leads to problems with locomotor system and constraint movability, resulting in closure of the vicious circle of gaining weight. Bariatric surgery is an effective weight loss method, but it is still unclear whether this procedure influences modification of physical activity routine.

**Methods:** 54 patients (55.56% females,  $n=30$ ) who underwent bariatric surgery at 2nd Department of General Surgery JU MC in Cracow, Poland from November 2015 to June 2016 were enrolled to this prospective study. Mean age, BMI and absolute waist circumference of participants were respectively:  $43.6 \pm 12.2$  y.o.,  $45.94 \pm 6.35$  kg/m<sup>2</sup> and  $128.39 \pm 13.45$  cm (female),  $146.9 \pm 17.21$  cm (male). One day before the procedure and one year later participants were asked to complete two standardized questionnaires: Paffenbarger Physical Activity Questionnaire and WHO Global Physical Activity Questionnaire, on the basis of which average physical activity in metabolic energy equivalents (MET-minutes) per week has been estimated

**Results:** One year after surgery MET-minutes has increased over 14 times (Me:299.75, Q1–Q3:225.78–358.38 vs. Me:4339.85, Q1–Q3:1590.6–7827.1,  $p < 0.00001$ ). Average time sitting or reclining has reduced from 480 to 300 min per day ( $p=0.00118$ ). Mean pace of walking has changed from  $<3.2$  km/h to average 3.2–4.8 km/h ( $p=0.00406$ ). Participants were also asked to rate on visual analogue scale their level of exertion during normal activities. This parameter has decreased from mean 5 (equivalent of strong effort) to mean 2.5 (equivalent of weak effort) ( $p=0.00004$ ).

**Conclusion:** Before the procedure none of participants has achieved recommended by WHO weekly level of 600 MET-minutes and after surgery 81% of them have exceed it. This data have shown significant positive changes in physical activity in patients who underwent bariatric surgery.

<http://dx.doi.org/10.1016/j.pbj.2017.07.044>

Internal Medicine Poster Session  
Friday, September 15th, 10h00

### PS085

#### Comparison of liver biopsy and non-invasive APRI test in assessing the stage of liver fibrosis in patients with chronic HCV infection



Nina Adzic\*, Mirjana Arapovic

Clinic for Infectious and Tropical Diseases "Prof. dr Kosta Todorovic", Clinical Center of Serbia, Faculty of Medicine University of Belgrade, Serbia

E-mail address: [adzicnina@gmail.com](mailto:adzicnina@gmail.com)

(N. Adzic).

**Aim:** The aim of this study was to evaluate the performance of non-invasive APRI score in predicting significant fibrosis and cirrhosis in patients with chronic HCV infection who underwent liver biopsy.

**Introduction:** Determining the stage of liver fibrosis is essential in managing patients with chronic hepatitis C virus. In chronic HCV infection, liver biopsy is the gold standard method for assessing stage of liver fibrosis, but it is invasive with potential complications. Non-invasive markers have been proposed and APRI score (aspartate aminotransferase (AST)-to-platelet ratio index) has been shown as a simple and inexpensive marker of liver fibrosis.

**Methods:** This retrospective study included 142 patients with chronic hepatitis C who had undergone liver biopsy from January 2013 to December 2015. Liver fibrosis was staged according to METAVIR (F0–F4) scoring system. The diagnostic performances of APRI score in predicting significant fibrosis (F2–F4) and cirrhosis (F4) were evaluated and compared by ROC curves.

**Results:** Fifty-three (37.3%) patients had significant fibrosis and 18 (12.7%) had cirrhosis. The areas under the ROC curve of APRI for predicting significant fibrosis and cirrhosis were 0.76 and 0.81. Using recommended cut-off values for APRI test, significant fibrosis could be identified in 26% and cirrhosis in 22% patients, but specificity for significant fibrosis was 88% and for cirrhosis 91%. Results have also shown that lower platelets count in our patients is associated with higher stage of fibrosis ( $p < 0.0001$ ).

**Conclusion:** APRI test shows low sensitivity and high specificity in the distinction between mild and significant fibrosis, and it shows good sensitivity in the evaluation of patients without cirrhosis and excellent specificity in patients with cirrhosis. Non-invasive biochemical tests and scores should be used only as additional criteria in differentiating the stage of liver fibrosis in chronic HCV infection, along with other non-invasive methods.

<http://dx.doi.org/10.1016/j.pbj.2017.07.045>

### PS144

#### Characteristics of patients with diagnosed chronic fungal rhinosinusitis surgically treated at the Clinic for otorhinolaryngology, Clinical centre of Vojvodina, in the past five years



D. Ignjić

University of Novi Sad, Faculty of medicine,  
Department of otorhinolaryngology, Serbia  
E-mail address: [dario.ignjic@gmail.com](mailto:dario.ignjic@gmail.com).

**Aim:** To determine the characteristics of patients with chronic fungal rhinosinusitis surgically treated at the Clinic for Otorhinolaryngology, Clinical Center of Vojvodina, from 2011 to 2016.

**Introduction:** The incidence of fungal rhinosinusitis in European counties is steadily growing. The reason behind this is the increased usage of immunosuppressive therapy, antibiotics and changes in everyday behaviors (increased stays in rooms with air-condition).

**Methods:** The study included 21 patients diagnosed with fungal rhinosinusitis. The patient's data was collected from their medical history.

**Results:** The mean age of the patients was  $45 \pm 16.51$ , with females being more often affected (11/21). The most commonly affected sinus was the maxillary sinus (54.67%), after that the sphenoid (20.83%), posterior ethmoid (18.5%), anterior ethmoid (8.33), and frontal (4.17%). Aspergillus was the most common cause (57.14%), mucormycosis was found once (4.78%). Staphylococcus aureus was isolated in 7 (33.33%) patients. Clinical symptoms were dominated by intensive facial pain and nasal secretion (found in all patients). Nasal congestion was present in 85.71% patients, less common was loss of sense of smell, in 47.62%. Endoscopic results showed significant differences between the characteristics of mucosa and mucus of the healthy and affected side of the patient's face. Significant differences are present in CT scans in all sinuses when the sinuses of the healthy and affected side of the patient's face were compared.

**Conclusion:** Clinical symptoms of patients with fungal rhinosinusitis were dominated by facial pain, nasal secretion and nasal congestion. Endoscopy shows pathological changes in the mucosa of the affected side of the patient's face, with viscous mucous secretions. Intraoperative findings show unilateral affection of the sinuses in all patients, most commonly in the maxillary sinus. The sphenoid sinus was less commonly affected, the ethmoid and frontal were rarely affected. Aspergillus is proven to be the most common cause.

<http://dx.doi.org/10.1016/j.pbj.2017.07.046>

## PS175

### Predictors for recurrent spontaneous intracerebral hemorrhage: A retrospective study

Catarina Castro Alves<sup>1,2,\*</sup>, Hipólito Nzwalo<sup>1,2</sup>, Ana Marreiros<sup>1,2</sup>

<sup>1</sup> Departamento de Ciências Biomédicas e Medicina, Universidade do Algarve, Portugal

<sup>2</sup> Algarve Biomedical Center, Universidade do Algarve, Portugal

E-mail address: [catarinacastro@yahoo.com](mailto:catarinacastro@yahoo.com) (C.C. Alves).

**Aim:** The aim of this work was to determine predictors for recurrence of intracerebral hemorrhage (ICH), which may allow the identification and more appropriate management of patients at higher risk of recurrent ICH.

**Introduction:** The prognosis of recurrent bleeding seems to be worse than that of the first ICH. However, only a limited number of studies report the frequency of recurrence in ICH and attempt to characterize which factors may be associated with tendency for repeated hemorrhagic events.

**Methods:** We analyzed data from 549 patients admitted for treatment of ICH at the Unit of Faro of the Algarve Hospital Center, followed over a period of 5 years. 189 patients with a Rankin at discharge equal to 6 were excluded from the analysis.

**Results:** We identified 24 patients (6.7%) with recurrent ICH. Recurrence was significantly more frequent in woman (10.9%) than in men (4.4%) ( $p < 0.05$ ). By comparing patients with recurrent ICH

with patients with isolated ICH, we found that recurrence was associated with more advanced ages at the time of the first ICH, but only for men (76 years for recurrent ICH and 68 years for isolated ICH) ( $p < 0.05$ ). There was a tendency towards recurrence when the bleeding was lobar (33% of recurrence, and 24% for non-lobar bleedings), although no statistical significance could be found. Other factors, such as previous hypertension were not associated with increased risk of recurrent ICH.

**Conclusion:** In this series of patients, we identified sex and age as predictors for ICH recurrence.

<http://dx.doi.org/10.1016/j.pbj.2017.07.047>

## PS133

### Randomized study to compare two methods of e-learning of ECG interpretation among medical students



Agnieszka Stępień\*, Michał Pacia, Sebastian Janiec, Wojciech Chemielak

Students' Scientific Group at the Department of Cardiac and Vascular Diseases IC UJ CM in The John Paul II Hospital in Krakow, Poland

E-mail address: [agaa.stepien@gmail.com](mailto:agaa.stepien@gmail.com) (A. Stępień).

**Aim:** To compare the effectiveness of two methods of ECG e-learning among medical students: collaborative e-learning (C-eL) and individual e-learning (I-eL).

**Introduction:** Electrocardiogram (ECG) interpretation is an essential skill in medicine. The best method of ECG education has not been determined.

**Methods:** Sixty 5th-year students from the Jagiellonian University Medical College were randomly assigned in a 1:1 ratio to the C-eL and I-eL groups. C-eL group students were further randomly divided into 6 subgroups of 5 students. Students from the I-eL group received by e-mail an ECG recording with comprehensive description every second day; at that time students from the C-eL group received the ECG recording without any description. C-eL students were encouraged to cooperate in analyzing the ECG in subgroups using internet platform and were expected to submit interpretation of the ECG recording to coordinator after 48 h. Afterwards they received comprehensive description of the ECG. Before starting the study all students participated in a pretest assessing their basic theoretic knowledge. The effects of e-learning were assessed at a final e-test. The main endpoint of the study was the number of students who passed the final e-test.

**Results:** Basic knowledge was similar in both study groups. Students from the I-eL group answered correctly to  $9.0 \pm 1.0$  ( $90 \pm 10\%$ ) and from the C-eL group to  $9.5 \pm 0.6$  ( $95 \pm 6\%$ ) questions,  $p = 0.07$ .

The main endpoint was achieved more frequently in the C-eL than in the I-eL group: 17 (63%) vs 10 (35.7%) students respectively,  $p = 0.045$ . C-eL group students, as compared to I-eL group students, achieved more points in the final e-test (12.3 vs. 11.0 points respectively,  $p = 0.036$ ) and also better results in ECG interpretation (4.1 vs. 3.4;  $p = 0.03$ ).

**Conclusion:** Collaborative e-learning of electrocardiography in 5th year medical students is superior to individual e-learning.

<http://dx.doi.org/10.1016/j.pbj.2017.07.048>



## PS096

### The influence of smoking cessation-related weight gain on cardiovascular risk in patients treated with drug-eluting stent after acute coronary syndrome

Aleksandra Buczyńska\*, Karol Kasprzycki, Aleksandra Pizun, Marta Tomica

Students' Scientific Group, The Department of Coronary Heart Disease, The John Paul II Hospital in Cracow Faculty of Medicine, Jagiellonian University Medical College, Poland

E-mail address: [abuczynska93@gmail.com](mailto:abuczynska93@gmail.com) (A. Buczyńska).

**Aim:** Aim of this study was to analyse association of smoking cessation influence on long-term clinical outcomes in patients with an acute coronary syndrome (ACS).

**Introduction:** Smoking and obesity are important cardiovascular risk factors. Patients often put on weight after quitting smoking.

**Methods:** 137 consecutive ACS patients of the Department of Coronary Heart Disease John Paul II Hospital in Cracow admitted between 2011 and 2013 were enrolled in the study. They had no previous history of Coronary Heart Disease and underwent Percutaneous Coronary Intervention with implantation of at least one Drug Eluting Stent. Telephone follow-up was carried out after a minimum time of 3 years. Patients were divided into 2 groups: patients who stopped smoking (91) and non-smokers (46) which were compared according to weight gain, increased of the BMI, morbidity of diabetes mellitus (DM), reasons of admission to hospital, occurrence of another ACS, stroke and neoplasm.

**Results:** The population consisted of 66% males, 34% females mean age 67 SD 11.29. Patients who used to smoke were significantly younger than non-smokers (64.99 vs. 71.37;  $P=0.048$ ). Both groups did not statistically differ in terms of gender and frequency of DM. The ex-smokers were admitted more frequently due to STEMI while in the non-smokers NSTEMI and Unstable Angina predominated. The weight and BMI in both groups did not differ on the date of ACS. However after 3 years there was a statistically significant difference: ex-smokers put on weight on average 1.3 kg while non-smokers lost 2.17 kg ( $p=0.01$ ). There was no association between the patients' history of smoking and occurrence of stroke, malignancy or another ACS.

**Conclusion:** Smoking cessation does not appear to influence long-term clinical outcomes after ACS. However it associates with weight gain which obviously increases cardiovascular risk. Our findings need further investigation and follow-up in a larger cohort of ACS patients.

<http://dx.doi.org/10.1016/j.pbj.2017.07.049>

## PS059

### Neonatal abstinence syndrome – Retrospective review

G. Knezović\*, N. Marić, V. Mijatović, A. Vejnović, V. Pavlović

Faculty of Medicine, University of Novi Sad, Department of Pharmacology, Toxicology and Clinical Pharmacology, Serbia

E-mail address: [goranknezovic@hotmail.com](mailto:goranknezovic@hotmail.com) (G. Knezović).

**Aim:** To evaluate the characteristics of newborns diagnosed with neonatal abstinence syndrome (NAS) and the characteristics

of their mothers in Vojvodina from 2012 to 2016., as well as the interrelationship of certain features.

**Introduction:** NAS is a collection of symptoms and signs that occur as a result of the sudden interruption of fetal exposure to certain substances (methadone, heroin, buprenorphine, etc.) that were used or abused by the mother during pregnancy.<sup>1,2</sup> It is manifested in a multitude of symptoms including central nervous system irritability, over-activity of the vegetative nervous system and dysfunction of the gastrointestinal tract.<sup>3,4</sup> The occurrence of NAS is closely related to the maintenance therapy of pregnant opioid addicts.<sup>1</sup>

**Methods:** This study analyzed medical records of women who gave birth at the Clinic of Gynaecology and Obstetrics in Novi Sad, whose children were diagnosed with NAS after birth, as well as the medical records of newborns treated at the Neonatology Department of the Institute for Child and Youth Health Care of Vojvodina diagnosed with NAS. Medical records included data from the medical history of the newborn and personal and gynaecological medical history of their mothers.

**Results:** A total of 41 cases of NAS were registered. An increase in incidence was noticed during the five-year period of about 15%. Mothers were mostly unemployed (80.49%). Slightly more than half of respondents (57.5%) during pregnancy were on one of substitution treatment modalities. The majority of newborns with NAS (75.61%) were male. The clinical picture was significantly more expressed in children whose mothers consumed methadone, compared to mothers who consumed heroin during pregnancy ( $p=0.0002$ ).

**Conclusion:** The incidence of diagnosed NAS cases is growing. Representation of male newborns with NAS is three times higher than female newborns. Methadone cause more NAS symptoms than heroin.

**Acknowledgements:** The Ministry of Education and Science of Republic Serbia (grant number 41012) supported this research work.

## References

- Kocherlakota P. Neonatal abstinence syndrome. *Pediatrics*. 2014;134:e547–61.
- Cramton REM, Gruchala NE. Babies breaking bad: neonatal and iatrogenic withdrawal syndromes. *Curr Opin Pediatr*. 2013;25:532–42.
- Ko JY, Patrick SW, Tong VT, Patel R, Lind JN, Barfield WD. Incidence of neonatal abstinence syndrome – 28 States, 1999–2013. *Morb Mortal Wkly Rep*. 2016;65:799–802.
- Bersani I, Corsello M, Mastandrea M, Patacchiola V, Foligno S, Garofalo V, et al. Neonatal abstinence syndrome. *Early Hum Dev*. 2013;89:s85–7.

<http://dx.doi.org/10.1016/j.pbj.2017.07.050>

## PS176

### Associations of epicardial adipose tissue thickness and cardiometabolic risk factors in STEMI patients treated with percutaneous coronary intervention

A. Gadeikytė<sup>1,\*</sup>, A. Varoniukaitė<sup>1</sup>, O. Gustienė<sup>2</sup>

<sup>1</sup> Lithuanian University of Health Sciences, Kaunas, Lithuania

<sup>2</sup> Department of Cardiology, Medical Academy, Lithuanian University of Health Sciences, Kaunas, Lithuania

E-mail address: [arvilegadeikyte@gmail.com](mailto:arvilegadeikyte@gmail.com) (A. Gadeikytė).

**Aim:** To evaluate EAT thickness in STEMI patients treated with percutaneous coronary intervention (PCI) and its associations with body mass index (BMI), blood lipids and acute left ventricular dysfunction.



**Introduction:** Epicardial adipose tissue (EAT), located between the myocardium and visceral layer of pericardium is an emerging risk factor for cardiometabolic diseases.

**Methods:** The retrospective study consisted of patients hospitalised for STEMI treated with PCI from 2014 to 2016. EAT thickness was measured from the parasternal long-axis view at end-diastole. Cholesterol levels were determined in a blood sample. According to median patients were divided in two groups: thin EAT group (<2.27 mm,  $n = 270$ ) and thick EAT group ( $\geq 2.27$  mm,  $n = 223$ ). Statistical analysis was performed with SPSS using Mann–Whitney test, *T*-test, logistic regression analysis. Values of cholesterol levels were evaluated by ROC curves.  $p < 0.05$  was significant.

**Results:** Total 492 patients (332 males,  $66.62 \pm 12.24$  year-old) were enrolled. Groups did not differ by age, gender, morbidity of diabetes mellitus and triglyceride levels. Patients had higher BMI ( $29.41 \pm 4.97$  vs.  $28.13 \pm 4.67$  kg/m<sup>2</sup>,  $p = 0.009$ ), total cholesterol ( $>4.82$  mmol/l: 35.2 vs. 26.4%,  $p = 0.024$ ), low density lipoprotein cholesterol ( $>2.5$  mmol/l: 45.8 vs. 33.3%,  $p = 0.004$ ) and reduced high density lipoprotein cholesterol (HDL-C) levels ( $\leq 1$  mmol/l: 24.4 vs. 10.4%,  $p = 0.009$ ) in thick EAT group. Logistic regression analysis revealed that higher BMI (OR = 1.532, 95% CI 1.008–2.328,  $p = 0.002$ ) and HDL-C  $\leq 1$  mmol/l (OR = 1.777, 95% CI 1.159–2.724,  $p = 0.008$ ) were associated with thicker EAT. Killip class  $\geq$ III was more frequent (17.6 vs. 10.3%,  $p = 0.02$ ) in thick than thin EAT group.

**Conclusion:** Increased EAT thickness was associated with obesity, cardiometabolic risk factors and influenced severity of left ventricular dysfunction.

<http://dx.doi.org/10.1016/j.pbj.2017.07.051>

#### PS123

#### Evaluation of spleen volume: Practical diagnostic role of linear measurements, 2D and 3D coefficients in computed tomography



Justyna Tęczar\*, Iwona Kucybała, Anna Gajdosz, Kamil Krupa, Jakub Wnuk, Maria Widomska

Students' Scientific Group at the Department of Diagnostic Imaging, Chair of Radiology, JU CM, Poland

E-mail address: [justyna.teczar@tlen.pl](mailto:justyna.teczar@tlen.pl)

(J. Tęczar).

**Aim:** The aim of the study was to find which linear measurements, field and volume coefficients correlate best with the real volume of the spleen and can be further used for determination of splenomegaly.

**Introduction:** Spleen is involved in a wide spectrum of abnormalities, which might lead to an increase in organ size. Splenic enlargement on CT is diagnosed basing on rather subjective criteria. The product of the length, estimated height and thickness of the spleen ("splenic index", cut-off  $\geq 480$ ) has also been proposed as an indicator for evaluating splenic size on CT.

**Methods:** Abdominal CT examinations of 153 patients' (77 females, 76 males) were retrospectively analysed in terms of maximal length, thickness, hilum thickness (axial plane), height (longest measurement in coronal plane), 90° height (maximum vertical height at coronal section), estimated height (number of axial scans where spleen was visible multiplied by the thickness of CT scans) (Impax Software) and real spleen volume (Vitrea software). Two-dimensional and three-dimensional coefficients were acquired through proper mathematical formulas. Splenomegaly cut-off: 314.5 ml. Pearson's correlation coefficient was calculated for the relationship between single, field, volume measurements and real volume (Statistica software).

**Results:** There was a statistically significant correlation between all single, field and volume measurements and real volume ( $p < 0.05$ ). For single measurements, the correlation is the strongest for height ( $r = 0.813$ , sensitivity 65%, specificity 91.7%, PPV 71.4%, NPV 95.6%). For two-dimensional, it is the coefficient calculated from length and 90° height ( $r = 0.918$ , 85%, 94.7%, 70.8%, 97.7%). For three-dimensional, it is the coefficient calculated from length, 90° height and hilum thickness ( $r = 0.919$ , 75%, 96.2%, 75%, 96.2%). Cut-off for splenic index from our calculations was  $\geq 1148$ .

**Conclusion:** Coefficient from length, 90° height and hilum thickness correlate best with the real volume of the spleen. Splenic index in our study is far from the perfection for clinical practice.

<http://dx.doi.org/10.1016/j.pbj.2017.07.052>

#### PS128

#### Influence of blood inflammatory parameters to erythropoietin resistance in haemodialysis patients



Skirmantė Rėkutė

Vilnius University Medicine Faculty, Lithuania

E-mail address: [skirmante.rekute@gmail.com](mailto:skirmante.rekute@gmail.com).

**Aim:** To evaluate the correlation between the inflammatory blood parameters and the resistance to EPO among the hemodialysis patients.

**Introduction:** Erythropoietin therapy is considered to be the standard treatment of anemia in chronic kidney disease patients, yet some patients do not respond well to this therapy. This is called EPO resistance and could be generally associated with the chronic inflammation.

**Methods:** A retrospective one single centre study, which analysed medical records of 30 HD patients who had advanced CKD and received EPO treatment in Vilnius University Hospital Santaros Clinics from 2016–2009 to 2016–2011. Data analysed – concentrations of C-reactive protein, neutrophils, lymphocytes, platelets, as well as EPO dose per kilo and hemoglobin concentration (measured at the beginning of the EPO therapy and one month after the treatment).

**Results:** Patients were grouped into 2 categories: 1 group ( $n = 14$ ) – concentration of hemoglobin increased, 2 group ( $n = 16$ ) – concentration decreased after treatment. In 1 group average concentration of platelets were statistically significantly ( $p = 0.039$ ) higher ( $230.2 \pm 73.70$ ), compared to 2 group ( $174.1 \pm 66.96$ ).

Furthermore, platelets concentration among patients with hemoglobin level of  $>100$  g/l ( $n = 17$ ) after one month of treatment were statistically significantly ( $p = 0.012$ ) higher ( $231.06 \pm 56.41$ ) compared to those patients with hemoglobin level of  $<100$  g/l ( $n = 13$ ) ( $160.08 \pm 78.17$ ) after treatment.

Additionally, patients with hemoglobin levels after one month of treatment  $100$ – $125$  g/l ( $n = 15$ ) were separated into two groups based on C-reactive protein level:  $>5$  (1 group) and  $<5$  (2 group). Average concentration of erythropoietin was statistically significantly higher in 1 group ( $n = 9$ ) ( $223.82 \pm 69.15$  VV/kg) than in 2 group ( $n = 6$ ) ( $116.68 \pm 59.68$  VV/kg).

Correlation analysis revealed that among patients with hemoglobin levels of  $<110$  g/l after treatment there is a statistically significant positive correlation (+0.428) between change of hemoglobin levels before and after treatment and erythropoietin dose and statistically significant ( $p = 0.023$ ) negative correlation (–0.481) with lymphocytes concentration in blood.

**Conclusion:** HD patients with a higher concentration of platelets respond to EPO therapy better than those with a lower concentration. Increased EPO dose results in higher Hgb

concentration, but the inflammatory environment could also lead to EPO resistance. Higher lymphocyte concentration in the blood results in lower Hgb concentration change during treatment. In order to achieve required Hgb change, the increase of CRP above the normal range may result in double the dose of EPO needed.

<http://dx.doi.org/10.1016/j.pbj.2017.07.053>

#### PS061

### The relationship between Calcium-Scor and the risk of coronary artery disease in patients with heart failure

Mahdi Safiabadi<sup>1,2,\*</sup>, Nasim Naderi<sup>1</sup>, Sepideh Taghavi<sup>1</sup>

<sup>1</sup> Department of Heart Failure and Transplantation, Shaheed Rajaei Cardiovascular, Medical and Research Center, Tehran University of Medical Sciences, Tehran, Iran

<sup>2</sup> Student Research Committee, Baqiyatallah University of Medical Sciences, Tehran, Iran  
E-mail address: [dr.masafi1990@yahoo.de](mailto:dr.masafi1990@yahoo.de)  
(M. Safiabadi).

**Aim:** The purpose of this study was evaluating relationship between coronary calcium score in detecting the risk of coronary artery disease in patients with heart failure.

**Introduction:** Heart failure (HF) is an abnormality of cardiac structure or function leading to failure of the heart to oxygen delivery. Angiography is discussed as a gold standard for diagnosis of coronary artery disease but Cardiac CT-Scan recently is typical imaging technique which is low-cost and non-aggressive technique to determine coronary artery calcification.

**Methods:** This is case-control study that was conducted in Services Hospital. All Patients referring to Heart failure department were EF (Ejection fraction)  $\leq 35\%$  and all of them previously examined by Coronary Angiography or Coronary CT-Angiography to know the coronary artery status. The case group was patients with CAD related heart failure and control group was patients with normal coronary or Non-CAD Related-HF. All patients in both groups were evaluated with Conventional CT-Scan for calculated the Calcium score.

**Results:** Ninety patients with HF divided into case group ( $n = 40$ ) and control group ( $n = 50$ ). The average of EF in case group was  $29.25 \pm 5.05$  and in control group was  $27.7 \pm 7.09$ . The amounts of calcium score in each Categories (Mild, Moderate, Severe and Extensive) in case group was 33%, 18%, 13% and 5%, but control group in Categories (Mild, Moderate, Severe) was 20%, 6% and 4% respectively.

There was a statistically significant correlation ( $r = 0.835$ ;  $p < 0.0001$ ) between calcium score and results of angiography. There was linear relationship between calcium score and age of patients with heart failure ( $r^2 = 0.807$ ). No significant difference was found between genders in terms of calcium score ( $p = 0.353$ ).

**Conclusion:** There was high correlation between calcium score and results of angiography. Calcium scoring is reliable tool for screening patients with CAD.

<http://dx.doi.org/10.1016/j.pbj.2017.07.054>



#### PS083

### Levels of 6-thioguanine nucleotides and clinical remission in inflammatory bowel disease – A systematic review and meta-analysis



M.M. Estevinho<sup>1,\*</sup>, J. Afonso<sup>1</sup>, I. Rosa<sup>2</sup>, P. Lago<sup>3</sup>, E. Trindade<sup>4</sup>, L. Correia<sup>5</sup>, C.C. Dias<sup>6</sup>, F. Magro<sup>1,7</sup>, on behalf GEDII (Portuguese IBD Group)

<sup>1</sup> Department of Pharmacology and Therapeutics, Faculty of Medicine of the University of Porto, Portugal

<sup>2</sup> Gastroenterology Department, Instituto Português de Oncologia de Lisboa, Lisboa, Portugal

<sup>3</sup> Gastroenterology Department, Centro Hospitalar do Porto, Porto, Portugal

<sup>4</sup> Department of Pediatrics, Centro Hospitalar São João, Porto, Portugal

<sup>5</sup> Department of Gastroenterology and Hepatology, Hospital de Santa Maria, University of Lisbon, Lisbon, Portugal

<sup>6</sup> Department of Community Medicine, Information and Decision in Health, Faculty of Medicine of the University of Porto, Portugal; CINTESIS - Centre for Health Technology and Services Research, Porto, Portugal

<sup>7</sup> Department of Gastroenterology, Faculty of Medicine, Centro Hospitalar São João, Porto, Portugal  
E-mail address: [mmestevinho@gmail.com](mailto:mmestevinho@gmail.com)  
(M.M. Estevinho).

**Aim:** This systematic review and meta-analysis aimed i) to assess the clinical value of 6-thioguanine nucleotides (6-TGN) thresholds (200, 225, 230, 235, 250 and 260 pmol/ $8 \times 10^8$  RBC); and ii) to compare mean 6-TGN concentrations between patients with active disease and those achieving remission.

**Introduction:** Thiopurines are widely used as immunosuppressive drugs in the management of inflammatory bowel disease even though their minimum effective dose and dose-response relationship remain controversial. In addition, the monitoring of thiopurines' pharmacological active metabolites is currently reserved for particular cases namely in refractory patients or when non-compliance or toxicity is suspected.

**Methods:** Literature search was carried out following PRISMA and Cochrane Collaboration Guidelines and four databases were used (PubMed, Web of Science, ScienceDirect and the Cochrane Central Register of Controlled Trials). Statistical heterogeneity was assessed using the I<sup>2</sup> statistic followed by subgroup and sensitivity analyses. Odds ratios (ORs) were computed under the random effects model.

**Results:** The systematic search identified 1384 records of which 25 matched the inclusion criteria and were retained for further analysis. From these, 22 were used in the cut-off comparisons while 12 were used in the 6-TGN mean differences analysis. The global OR for remission in patients with 6-TGN concentrations above the predefined thresholds was 3.95 (95%CI, 2.63–5.94;  $p < 0.001$ ). When considering each of the six thresholds individually, the OR was significant for levels above 235 pmol/ $8 \times 10^8$  RBC (OR = 2.25) and 250 pmol/ $8 \times 10^8$  RBC (OR = 4.71). Mean 6-TGN levels were significantly superior among patients achieving clinical remission, with a pooled difference of 63.37 pmol/ $8 \times 10^8$  RBC (95%CI, 31.81–94.93;  $p < 0.001$ ).

**Conclusion:** These results reinforce that 6-TGN levels are related to clinical remission and give an insight into the thresholds that may be used to guide clinical decisions.

**Acknowledgements:** This work was supported by the Portuguese Group of Studies in Inflammatory Bowel Disease (GEDII).

<http://dx.doi.org/10.1016/j.pbj.2017.07.055>

## PS020

### Influence of energy drinks on hemodynamic parameters in young healthy adults – Randomized double-blind placebo controlled cross-over study



M. Niemczyk\*, M. Stopa, M. Łobacz, K. Rutkowska, A. Radko

*Students' Scientific Group at the 1st Department of Cardiology, Interventional Electrophysiology and Arterial Hypertension in Cracow, Poland*  
E-mail address: [magdaniemczyk29@gmail.com](mailto:magdaniemczyk29@gmail.com)  
(M. Niemczyk).

**Aim:** Assessment of the influence of single dose of energy drink on blood pressure, heart rate, ECG, cardiac output and vascular compliance in healthy volunteers.

**Introduction:** An energy drink (ED) is a type of beverage containing stimulant drugs, caffeine, taurine, which is marketed as providing mental and physical stimulation. The popularity of product is increasing especially among teenagers and young adults. Some research suggest that its consumption may have negative effect on cardiovascular system.

**Methods:** A randomized double-blind placebo controlled cross-over study was conducted on 18 healthy volunteers (7 female, 11 male, mean age  $23.67 \pm 1.19$ ). Subjects received: 500 ml of energy drink containing 160 mg of caffeine, 2 g of taurine and 50 mg of guarana or 500 ml of placebo. Participants drank beverages in random order during two different meetings. Drinks did not differ in taste, smell and color. In all participants before and after consumption of a drink following procedures were performed: peripheral and central systolic and diastolic blood pressure (SBP and DBP) measurement, ECG recording, echocardiography, and pulse wave velocity analysis – in the same sequence and time intervals for every participant.

**Results:** ED consumption was related to significant increase of SBP in 75 min of observation compared to placebo ( $\Delta$ SBP for ED  $5.7 \pm 10.2$  mmHg vs  $-0.3 \pm 7.2$  mmHg for P,  $p=0.03$ ). ED caused increase in central SBP ( $107.8 \pm 13.2$  vs  $115.6 \pm 12.1$  mmHg,  $p=0.0005$ ), and central DBP ( $73.9 \pm 11.9$  vs  $78.1 \pm 10.2$  mmHg,  $p=0.02$ ). However comparison between placebo and ED revealed no significant differences in these parameters. The ECG parameters (HR, PQ, QRS and QTc intervals, axis of P wave, QRS complex, T wave) did not reveal significant differences between groups. There were no differences in echocardiographically determined cardiac output and LVEF.

**Conclusion:** Single dose ED consumption increases peripheral and central SBP. This effect is probably mediated by vascular wall properties and not by cardiac performance.

<http://dx.doi.org/10.1016/j.pbj.2017.07.056>

## PS055

### Analysis of genetic polymorphism 4a/b of the eNOS gene in infertile men



Miloš Lazić\*, Đorđe Radisavčević

*Institute of Human genetics, University of Belgrade School of Medicine, Germany*

E-mail address: [miloslazic1512@live.com](mailto:miloslazic1512@live.com)  
(M. Lazić).

**Aim:** The aim of our study was the analysis of genetic polymorphism 4a/b of the eNOS gene in infertile men with idiopathic infertility, correlation of genotype and phenotype in infertile men and comparing the results of testing of genetic polymorphism 4a/b with the results from the control group.

**Introduction:** Male infertility of unknown etiology represents a common medical and social problem, in whose basis lies a combination of genetic and environmental factors. Several recent studies have pointed to the possible connection of polymorphisms in eNOS gene and idiopathic male infertility.

**Methods:** The study included 50 infertile men with idiopathic infertility and 50 fertile controls. 4a4b polymorphism was detected by polymerase chain reaction (PCR).

**Results:** 4b4b genotype was detected in 27 (54%) patients and 36 (72%) controls, 4a4b genotype in 21 (42%) patients and 13 (26%) controls and 4a4a genotype detected in 2 (4%) patients and 1 (2%) control group participant. 4b allele frequency was 75% in the patient population and 85% in the control population, and frequency of allele 4a was 25% with patients and 15% in the control group. There was no statistically significant difference in the distribution of genotypes ( $p=0.062$ ) nor alleles ( $p=0.111$ ) between these two populations. Comparing 4a/b genotypes and serum concentration of FSH within patient group, we've detected a highly significant correlation ( $p<0.001$ ), where all carriers of 4b4b genotype had physiological concentration of serum FSH, while most of 4a4a and 4a4b carriers had higher serum FSH values.

**Conclusion:** Per our results VNTR (4a/b) is not connected to idiopathic male infertility in Serbian men, but they did show a highly significant correlation between serum FSH concentration and 4a/b genotype of infertile men.

<http://dx.doi.org/10.1016/j.pbj.2017.07.057>

## PS115

### Intravenous iron treatment effect to patients on hemodialysis



Vaida Kazlauskaitė\*, Skirmante Rekite

*Vilnius University, Faculty of Medicine, Lithuania*  
E-mail address: [kazlauskaitė.void@gmail.com](mailto:kazlauskaitė.void@gmail.com)  
(V. Kazlauskaitė).

**Aim:** To evaluate the coherence between intravenous iron therapy and the inflammatory indicators to patients on hemodialysis.

**Introduction:** when the kidney function is failing, the number, of patients who has a final stage kidney disease with anemia, is increasing. One of the most important reasons of anemia is iron deficiency. The iron treatment may be intravenous or oral. Though the oral treatment is cheaper, it may cause gastrointestinal disorders. Intravenous iron therapy has a better tolerance, but earlier studies had showed that it increases the risk of infections to patients on hemodialysis.

**Methods:** The retrospective study included 33 hemodialysis patients who undergone the intravenous therapy during the 2016-10 and 2016-12 in Vilnius university hospital. The absolute numbers of neutrophils and lymphocytes, C-reactive protein and procalcitonin were assessed before the treatment with intravenous iron and a month after it.

**Results:** we analyzed 13 men and 20 women, the mean age 59 years, the mean creatinine  $760 \mu\text{mol/l}$ , the mean hemoglobin  $105 \text{g/l}$ . By the test of Wilcoxon signed rank the means of neutrophils and C-reactive protein increased after the start of the treatment with iron (the mean of C-reactive protein increased

from  $12.8 \pm 12.96$  to  $27.4 \pm 41.17$ ;  $p=0.07$ ; the mean of neutrophils increased from  $4.45 \pm 1.52$  to  $6.86 \pm 12.11$ ;  $p=0.59$ ). The *T*-test showed that the means of procalcitonin increased from  $0.21 \pm 0.07$  to  $0.23 \pm 0.08$ ,  $p=0.04$ , the mean of lymphocytes increased from  $1.35 \pm 0.54$  to  $1.54 \pm 0.62$ ,  $p=0.1$ . Pearson correlation coefficient showed statistically insignificant positive correlation between the dose of medication and variation of procalcitonin.

**Conclusion:** The study has showed that inflammatory indicators increased after the intravenous iron therapy to patients on hemodialysis.

<http://dx.doi.org/10.1016/j.pbj.2017.07.058>

### PS038

#### Distribution and quantification of elements of the enteric nervous system in the distal rectum of neonates and infants



S. Lestarevic<sup>1,\*</sup>, M. Lazic<sup>1,2</sup>, R. Jankovic<sup>2</sup>

<sup>1</sup> School of Medicine, University of Belgrade, Serbia

<sup>2</sup> Institute of pathology, School of Medicine, University of Belgrade, Serbia

E-mail address: [sanjalestarevic@gmail.com](mailto:sanjalestarevic@gmail.com)

(S. Lestarevic).

**Aim:** Analysis of variations in the ENS of distal rectum in neonates and infants under the age of 6 months, with no previous history of intestinal dysmotility.

**Introduction:** The enteric nervous system (ENS) consists of numerous ganglia along the gastrointestinal tract. The most common disorder of ENS is Hirschsprung's disease (HD). Diagnostic problems may occur due to insufficient knowledge of the normal distribution of ganglion cells (GC) in the distal rectum.

**Methods:** The study analyzed ENS of distal rectum in autopsy samples of infants. The sections were stained with hematoxylin and eosin (H&E) and immunohistochemistry using the MAP-2 antibodies. All sections were analyzed at three levels: the level of anorectal junction (ARJ0), at 1 cm (ARJ1) and 2 cm (ARJ2) proximal to the ARJ0. We analyzed number of ganglia and GC, their distribution and thickness of the bundles of nerve fibers (BNF).

**Results:** GC were found at ARJ0 mainly within BNF of the intramuscular zone. Number of GC within BNF of intramuscular zone were lower at ARJ2 than ARJ1 (H&E:  $p=0.021$ ; MAP-2:  $p=0.017$ ). Number of GC in submucosal ganglia were significantly higher in ARJ1 and ARJ2 compared to ARJ0. In myenteric ganglia the number of GC were higher at ARJ1 compared to ARJ0 (H&E:  $p=0.002$ ; MAP-2:  $p=0.014$ ). Number of GC were significantly higher at ARJ2 compared to ARJ1 only in MAP-2 staining ( $p=0.009$ ). In submucosal plexus we observed higher number of ganglia at ARJ1 and ARJ2 ( $p=0.014$ , both) compared to ARJ0 at MAP-2. In myenteric plexus there were higher number of ganglia at ARP1 compared to ARP0 (H&E:  $p=0.006$ ; MAP-2:  $p=0.014$ ). Individual thicker BNF were found in submucosa.

**Conclusion:** In distal rectum of neonates and infants there are significant variations in number of ganglia in the submucosal plexus up to ARJ2 and in myenteric plexus up to ARJ1.

<http://dx.doi.org/10.1016/j.pbj.2017.07.059>

### PS220

#### Inflammatory bowel diseases: Nutritional status and its significance for the course of the disease



Magdalena Achtenberg, Urszula Skorus\*

Chair of Gastroenterology, Hepatology and Infectious Diseases, Jagiellonian University Medical College, Poland

E-mail address: [urszula.skorus@gmail.com](mailto:urszula.skorus@gmail.com)

(U. Skorus).

**Aim:** The aim of the study was to evaluate the association between the Body Mass Index (BMI) and the disease course of IBD patients.

**Introduction:** Inflammatory Bowel Disease (IBD) may lead to the underweight and malnourishment. However, the number of overweight and obese patients increases. Excess body weight connected with a pro-inflammatory state can modify the disease course.

**Methods:** Medical records from the University Hospital in Cracow Electronic System were screened from August 01, 2015 to December 31, 2016 in search of patients diagnosed with IBD. Data regarding the disease extension, occurrence of intestinal and extra-intestinal complications, number of days spent in the hospital annually and type of treatment was collected. The results were analyzed in the groups based on BMI ( $1 < 18.5$ ;  $2: 18.5-25$ ;  $3 > 25 \text{ kg/m}^2$ ).

**Results:** 150 patients with Crohn's disease (CD) and 151 with ulcerative colitis (UC) were included. The median number of days spent in the hospital annually was significantly higher in the underweight group (13 (IQR: 11) vs 7 (IQR: 17) vs 7 (IQR: 12);  $p < 0.01$ ). Overweight patients were less likely to receive anti-TNF or immunosuppressive treatment [anti-TNF (1: 35% vs 2: 38.36% vs 3: 18.29%; 1 vs 3;  $p=0.02$ ; 2 vs 3;  $p < 0.01$ ); immunosuppressive (1: 40.00% vs 3: 23, 17%;  $p=0.03$ )]. Patients with BMI  $> 25 \text{ kg/m}^2$  developed fistulas and bowel strictures less often [fistulas (1: 33.33% vs 2: 27.04% vs 3: 12.20%; 1 vs 3;  $p < 0.01$ ; 2 vs 3;  $p < 0.01$ ); strictures (1: 25% vs 2: 22, 64% vs 9.76%; 1 vs 3;  $p=0.01$ ; 2 vs 3;  $p < 0.01$ )]. Underweight UC patients had more extensive disease [left sided (1: 25% vs 2: 52.63% vs 3: 49.02%; 1 vs 2;  $p=0.02$ ; 1 vs 3;  $p=0.04$ ); pancolitis (1: 58.33% vs 2: 26.32% vs 3: 31.37%; 1 vs 2;  $p < 0.01$ ; 1 vs 3;  $p=0.02$ )].

**Conclusion:** Overweight seems to be associated with a milder clinical course of the disease in IBD patients. It is related to lower incidence of intestinal complications among CD and to less extensive intestine involvement in UC patients.

**Acknowledgements:** Dorota Cibor MD, PhD, Małgorzata Zwolińska-Wcisło MD, PhD, Associate Professor.

<http://dx.doi.org/10.1016/j.pbj.2017.07.060>

### PS023

#### Influence of glycoregulation and chronic degenerative complications of diabetes on bone mineral density



M. Džeba\*, A. Kovačić

Medical Faculty, University of Novi Sad, Serbia

E-mail address: [miiickey9.58@gmail.com](mailto:miiickey9.58@gmail.com)

(M. Džeba).

**Aim:** The aim of this study is to determine the correlation between duration of diabetes, glycoregulation and chronic degenerative complications of diabetes, on one side, and bone mineral density, on the other side.

**Introduction:** Diabetes mellitus is a state of chronic hyperglycaemia. In late stages of the disease, especially if it is not regulated well, chronic complications may occur, dominating the clinical picture. Osteoporosis is characterized by bone loss per volume unit leading to microarchitectonics disorder of the bone. Connection between diabetes and osteoporosis is very complex.

**Methods:** Medical documentation collected at daily hospital of Clinic of endocrinology, diabetes and metabolic disorders is used in this study. Sample includes 60 patients which have been diagnosed with diabetes mellitus, with or without complications, who underwent densitometry measurement (DEXA). Glycosylated hemoglobin (HbA1c), fasting glucose and postprandial glucose are used as parameters of glycoregulation.

**Results:** Average duration of diabetes is  $15.61 \pm 9.63$  years. Average value of HbA1c is  $8.5 \pm 1.79\%$ , average value of fasting glucose is  $9.23 \pm 2.94$  mmol/l and average value of postprandial glucose is  $11.35 \pm 4.27$  mmol/l. 67% of patients have one or more complications. Bone mineral density ( $\text{g}/\text{cm}^2$ ) of femoral neck and total have significant negative correlation with HbA1c ( $p < 0.01$ ). Bone mineral density of lumbosacral spine and femoral neck ( $\text{g}/\text{cm}^2$ ,  $T$ -score) have light negative correlation with postprandial glucose.

**Conclusion:** Bone mineral density and parameters of glycoregulation have negative correlation. Statistically significant correlations between bone mineral density and chronic degenerative complications of diabetes were not found.

<http://dx.doi.org/10.1016/j.pbj.2017.07.061>

#### PS022

##### Effect of autologous stem cell transplantation in patients with hematological malignancies



A. Kovačić\*, M. Džeba

Medical Faculty, University of Novi Sad, Serbia  
E-mail address: [alex.ak.bb@gmail.com](mailto:alex.ak.bb@gmail.com)  
(A. Kovačić).

**Aim:** The aim of this study is to analyse available medical data of patients diagnosed with multiple myeloma (MM), lymphoma Hodgkin (MB) and non-Hodgkin (NHL) and acute leukemia (AL), who underwent ASCT, and to compare the results with the results from other scientific works.

**Introduction:** Autologous stem cell transplantation (ASCT) with high dose chemotherapy is effective and safe approach in the treatment of different hematological malignancies. Nowadays, it is the standard therapy for multiple myeloma, lymphomas and acute leukemias.

**Methods:** Retrospective study included 84 patients diagnosed with MM, MH, LNH and AL who underwent ASCT in the period from 2004 to 2016. Data are presented in table and charts.

**Results:** In relation to the underlying disease, the distribution of respondents was as follows: 35 patients with MM, 24 with NHL, 20 with MH and 6 with AL. Large volume apheresis procedure had to 75 patients (89.3%), and 9 patients (10.7%) had conventional two-day procedure. The mean value of processed blood volume amounted to 13050 ml. The average number of MNC in the apheresis product was  $7.8 \times 10^8/\text{kg}$  bw, a CD34+ cells was  $12.11 \times 10^6/\text{kg}$  bw. After the application of conditioning regimens, depending on the underlying disease, neutrophils engraftment occurs at 11 day and platelets engraftment at 14 day.

**Conclusion:** Analyzing data of the patients with hematological malignancies and ASCT conducted, we conclude that the mentioned procedure is successful method of treatment, with low

transplant mortality and complications caused by the mentioned procedure.

<http://dx.doi.org/10.1016/j.pbj.2017.07.062>

#### PS237

##### Polymorphism of Kibra gene in patients with terminal renal insufficiency



N. Paovica\*, D. Bajovic, I. Novakovic

Institute of Human Genetics, Faculty of Medicine,  
University of Belgrade, Serbia  
E-mail address: [2carry.out2@gmail.com](mailto:2carry.out2@gmail.com)  
(N. Paovica).

**Aim:** The aim of this study was to determine whether there is a difference in frequencies of genotypes and alleles of KIBRA gene polymorphism, rs17070145 between patients with terminal renal insufficiency and normal population.

**Introduction:** KIBRA gene has a role in signal transmission that regulates apoptosis, proliferation, and movements of the cytoskeleton of cells. Due to its most common expression in kidney and brain, the name of this protein is Kibra (Kidney, BRAin). Polymorphism rs17070145 (substitution of thymine with cytosine in the ninth intron of the gene) is associated with Alzheimer's disease and memory, while its connection with kidney's diseases has not been tested yet. It is thought that allele C is the factor of predisposition in TRI.

**Methods:** Polymorphism rs17070145 was analyzed with Real Time PCR method using TaqMan probes and 50 people with TRI were involved. Results of gene analysis for the control group were taken from previous research. Frequencies of genotypes and alleles between patients with TRI and healthy examinees was compared with  $\chi^2$  (chi-square) test.

**Results:** The frequency of CC genotype among patients with TRI is 76%, CT genotype 22% and TT genotype 2%. Based on frequencies of genotypes, we found that frequency of C allele is 87%, while the frequency of T allele is 13%.

**Conclusion:** Results of  $\chi^2$  test show extremely statistically significant difference in frequencies of genotypes and alleles in patients with TRI in comparison with healthy people ( $P < 0.0001$ ). These results indicate that C alleles on locus rs17070145 in KIBRA gene are probably the significant factor of predisposition in the pathogenesis of TRI.<sup>1-3</sup>

#### References

1. Kremerskothen J, Plaas C, Buther K, et al. Characterization of KIBRA, a novel WW domain-containing protein. *Biochem Biophys Res Commun.* 2003;300:4.
2. Dunning K, Schurek EM, Schluter M, et al. KIBRA modulates directional migration of podocytes. *J Am Soc Nephrol.* 2008;19.
3. Maksimovic N. Analysis of polymorphisms in APOE, BDNF, BDNF and KIBRA genes and their correlation with memory performance in the population of students (Doctoral dissertation). NaRDuS – National Repository of Dissertations in Serbia.

<http://dx.doi.org/10.1016/j.pbj.2017.07.063>

Neurosciences Poster Session  
Thursday, September 14th, 16h00

#### PS088

##### D-Galactose high-dose administration and oral epigallocatechin-3-gallate effects on the dendritic trees of developing neurons of young male rats



F. Barreto<sup>1,2,\*</sup>, R. Rodrigues<sup>1,2</sup>, A. Cardoso<sup>1,2</sup>,  
J.P. Andrade<sup>1,2</sup>

<sup>1</sup> Department of Biomedicine – Unit of Anatomy, Faculty of Medicine, University of Porto, Alameda Prof. Hernâni Monteiro, 4200-319 Porto, Portugal

<sup>2</sup> Center of Health Technology and Services Research (CINTESIS), Faculty of Medicine, University of Porto, Rua Dr. Plácido da Costa, 4200-450 Porto, Portugal  
E-mail address: [fbarreto44@hotmail.com](mailto:fbarreto44@hotmail.com) (F. Barreto).

**Aim:** In the present study, we aimed to explore the effect of D-galactose administration and epigallocatechin-3-gallate (EGCG) on the dendritic trees of developing granule cells of the hippocampal formation (HF) of young male rats.

**Introduction:** The model of accelerated senescence with the administration of D-galactose is used in anti-aging studies. However, reports have questioned its effectiveness. To clarify this issue we used high-dose D-galactose on young rats and studied the immature granule cells stained with the neurogenesis marker doublecortin (DCX). We also used EGCG, a green tea catechin, to verify if there are neuroprotective effects in the D-galactose-treated animals.

**Methods:** At 4 weeks of age, male Wistar rats were allocated to a control group ( $n=7$ ), a D-galactose group (300 mg/kg body weight, intraperitoneally) ( $n=5$ ; GAL) and to a D-galactose + EGCG (oral solution, 2 grams/L) group ( $n=5$ ; gal + EGCG) during 4 weeks. After this period DCX immunocytochemistry was performed. The dendritic trees of immature granule cells were drawn with the aid of a camera lucida and a metric analysis of the dendritic segments of the dendritic trees was performed.

**Results:** No differences in all parameters quantified were found when controls and gal rats were compared. However, the results show that the total dendritic length of the dendritic trees of gal + EGCG rats was significantly reduced when compared with controls ( $p < 0.03$ ). There were no differences in the others dendritic parameters quantified.

**Conclusion:** D-Galactose did not induce disturbance of the neurogenesis as shown by the absence of alterations in the dendritic trees confirming our previous studies. Surprisingly, the addition of EGCG led to a reduced total dendritic length. This unexpected effect can be explained if we consider that the addition of the catechin acted as a second aggression leading to a disturbed dendritic tree of the immature neurons.

**Acknowledgements:** This article was supported by ERDF through the operation POCI-01-0145-FEDER-007746 funded by the Programa Operacional Competitividade e Internacionalização – COMPETE2020 and by National Funds through FCT - Fundação para a Ciência e a Tecnologia within CINTESIS, R&D Unit (reference UID/IC/4255/2013).

<http://dx.doi.org/10.1016/j.pbj.2017.07.064>

## PS079

### Analysis of variations in the F5, F2 and ACE genes among Latvian patients with ischemic stroke

Anna Inese Tutāne

Rīga Stradiņš University, Latvia

E-mail address: [annatutane@gmail.com](mailto:annatutane@gmail.com).

**Aim:** Evaluate thrombophilia causing genetic variants and ACE gene I/D variant impact on patients with ischemic stroke.

**Introduction:** Every year, 15 million people worldwide suffer a stroke that is the second leading cause of disability. Genetic variants in Leiden factor coding gene (F5) and in prothrombin gene (F2) cause inherited thrombophilia which is associated with increased

risk of intravascular thrombosis, thromboembolism and cerebral stroke. Angiotensin-converting enzyme (ACE) coding gene I/D variant is discussed among numerous conditions including stroke.

**Methods:** In the study there were included 115 patients with mean age  $70.3 \pm 11.0$  years, with diagnosed ischemic stroke. Control group for F5 and F2 gene variations consisted of 124 individuals with mean age  $55.6 \pm 14.6$  years. And for ACE gene variation 248 individuals with mean age  $56.8 \pm 11.4$  years. DNA was extracted from peripheral blood using standard phenol-chloroform method. Genotyping of F5 gene variant G1691A and F2 gene variant G20210A was performed using PCR-RFLP. ACE gene I/D variant genotyping were performed using PCR. Statistical analysis was performed using Fisher's exact test and SPSS v22.0 software.

**Results:** F2 gene variant were more frequent in patient group. Frequency in patients were 0.017 and in control group 0 ( $p = 0.038$ ). F5 gene variant frequency in both patients and control group were 0.012 ( $p > 0.05$ ). Seven patients (5.6%) had one variant in one of coagulation factors encoding genes comparing to three in control group (2.4%) ( $p > 0.05$ ). Mean age for patients with identified variations in F2 or F5 was not significantly different comparing to other patients ( $p > 0.05$ ). ACE gene I/D genotypes and allele frequencies in stroke patients were not significantly different from controls – I allele frequencies were 0.452 in patients versus 0.470 in controls ( $p > 0.05$ ).

**Conclusion:** Prothrombin encoding gene variant G20210A could be risk factor for ischemic stroke. F5 and ACE gene I/D genotypes are not associated with ischemic stroke.

<http://dx.doi.org/10.1016/j.pbj.2017.07.065>

## PS205

### The bioactive compounds from elderberry to modulate mitochondrial dysfunctions underlying Alzheimer's disease



Dina Neves<sup>1,\*</sup>, João Bernardo<sup>1</sup>,  
Patrícia Valentão<sup>1</sup>, Maria C. Oliveira<sup>2</sup>,  
David M. Pereira<sup>1</sup>, Paula B. Andrade<sup>1</sup>,  
Romeu A. Videira<sup>1</sup>

<sup>1</sup> REQUIMTE/LAQV, Laboratório de Farmacognosia, Departamento de Química, Faculdade de Farmácia, Universidade do Porto, Rua de Jorge Viterbo Ferreira, N.º 228, 4050-213 Porto, Portugal

<sup>2</sup> Centro de Química de Vila Real (CQ-VR), Departamento de Química; Escola de Ciências da Vida e do Ambiente, Universidade de Trás-os-Montes e Alto Douro (UTAD), P.O. Box 1013, 5001-801 Vila Real, Portugal  
E-mail address: [up201302607@ff.up.pt](mailto:up201302607@ff.up.pt) (D. Neves).

**Aim:** The specific objective of this work is to establish a correlation between the physical-chemical properties of the aqueous extract of elderberry (*Sambucus nigra* L.) and its ability to tune the cell redox state and to overcome mitochondrial dysfunctions, which are pathological events with high relevance in Alzheimer's disease (AD).

**Introduction:** Currently, there is no effective medicine to prevent or delay the progressive brain degeneration underlying cognitive decline and dementia that characterize AD. Previous works support the idea that the loss of mitochondrial functionality, connected with the decline of complex I activity, is able to promote AD phenotype through the activation of multiple pathophysiological pathways, including oxidative stress, neuroinflammation, and also tau and amyloid-beta pathologies. Thus, multi-targeted



approaches supported by mixtures of natural bioactive compounds should reveal more effectiveness than classical therapeutics for AD.

**Methods:** The polyphenolic profile of elderberry extract and of anthocyanin-enriched fraction was evaluated by HPLC-DAD, the optical properties by UV-vis and fluorescence spectroscopy and the redox behavior by cyclic voltammetry. Antioxidant properties were assessed in cell-free assays while the ability the elderberry extract to modulate the mitochondrial redox chain was evaluated in rat brain mitochondria.

**Results:** HPLC analyses showed that elderberry extract is a mixture of chemical compounds, particularly rich in anthocyanins. It exhibits intrinsic fluorescence properties with potential for bioimaging, reversible redox behavior and ability to scavenge DPPH, nitric oxide and superoxide radicals. The antioxidant, optical and redox properties of elderberry extract are strongly correlated to their content in anthocyanins. Bioenergetic studies show that elderberry extract has ability to promote the oxidation of NADH in aqueous phase and deliver electrons to ubiquinone or complex III in the inner-mitochondrial membrane, overcoming the complex I inhibition promoted by rotenone.

**Conclusion:** Elderberry anthocyanins have potential to be used in mitochondria-targeted formulations to modulate the pathophysiological changes underlying AD from their early stages.

**Acknowledgements:** This work received financial support from National Funds (FCT/MEC) through project UID/QUI/50006/2013, co-financed by FEDER through COMPETE, under the Partnership Agreement PT2020, and from NORTE 2020, under the PORTUGAL 2020 Partnership Agreement, through ERDF (NORTE-01-0145-FEDER-000024).

<http://dx.doi.org/10.1016/j.pbj.2017.07.066>

## PS016

### Effect of resveratrol on the cartilage and nociceptive system of Osteoarthritic animals



S. Rosas<sup>1,2,3,\*</sup>, T. Aguiar<sup>1,2,3</sup>, L. Almeida<sup>1,2,3</sup>,  
D. Nascimento<sup>1,2,3</sup>, S. Adães<sup>1,2,3</sup>,  
J.M. Castro-Lopes<sup>1,2,3</sup>, F.L. Neto<sup>1,2,3</sup>,  
J. Ferreira-Gomes<sup>1,2,3</sup>

<sup>1</sup> Department of Biomedicine - Experimental Biology Unit, Faculty of Medicine of the University of Porto, Porto, Portugal

<sup>2</sup> Pain Research Group, Institute for Molecular and Cell Biology (IBMC), Porto, Portugal

<sup>3</sup> i3S - Institute for Investigation and Innovation in Health, Porto, Portugal

E-mail address: [suu.rosas@gmail.com](mailto:suu.rosas@gmail.com)

(S. Rosas).

**Aim:** This study aims to evaluate the effect of RV on the nociceptive behavior, histopathological alterations at the knee and DRG neurons of OA rats.

**Introduction:** Osteoarthritis (OA) is a common degenerative joint disease and arthritic pain is a prominent symptom associated with reduced quality of life. Peripheral pain mechanisms seem to be involved, with cartilage lesions showing a repercussion in Dorsal Root Ganglia (DRG) neurons. Resveratrol, a polyphenol with proven anti-inflammatory, anti-oxidant and neuroprotective properties, has been shown to prevent development of OA and act as an antinociceptive agent. However, its systemic effects once the disease has fully developed remain unclear.

**Methods:** To evaluate this, OA was induced in 18 male Wistar rats through intra-articular injection of mono-iodoacetate (MIA) (day 0). Animals were allowed to develop the disease for two weeks,

after which followed a 4-week-long treatment with resveratrol or vehicle, administered intraperitoneally twice daily (10 mg/kg). Nociceptive behavior was quantified weekly using the CatWalk and Knee-Bend tests. Animals were sacrificed one week after the last treatment administration, their knees were dissected for histopathological analysis, and the DRG were dissected and processed for immunohistochemical evaluation of activating transcription factor 3 (ATF-3) neuronal expression.

**Results:** Resveratrol was unable to prevent cartilage degeneration but it significantly decreased ATF-3 expression. The nociceptive behavior of OA animals treated with resveratrol decreased during the first three weeks of treatment, in comparison to day 14 (before treatment was initiated), as shown by Knee-Bend scores. However, this tendency reverted as the disease progressed.

**Conclusion:** These results indicate that resveratrol may have antinociceptive effects in the early stages of the disease development, but it might not play such a relevant role once the disease has progressed. Thus, further studies are needed to fully understand the possible role of resveratrol in the different stages of OA.

**Acknowledgements:** The study was supported by the Chair on Pain Medicine of the Faculty of Medicine, University of Porto and by the Grünenthal Foundation – Portugal. By the time the study was conducted DN was receiving a doctoral grant (SFRH/BD/79497/2011) by Fundação para a Ciência e a Tecnologia (FCT), Portugal and SA had a doctoral grant by Fundação Calouste Gulbenkian, Portugal.

<http://dx.doi.org/10.1016/j.pbj.2017.07.067>

## PS173

### Bupivacaine treatment enhances the regeneration of the lesioned external urethral sphincter of the rat



J.P. Morais<sup>1,\*</sup>, M. Torrado<sup>2,3,4</sup>, A. Avelino<sup>2,3,4</sup>

<sup>1</sup> Escola das Ciências da Vida e Ambiente, Universidade de Trás-os-Montes e Alto Douro

<sup>2</sup> Department of Biomedicine, Experimental Biology Unit, Faculty of Medicine, University of Porto, Portugal

<sup>3</sup> Translational NeuroUrology Group, IBMC - Instituto de Biologia Molecular e Celular, Porto, Portugal

<sup>4</sup> i3S - Instituto de Investigação e Inovação em Saúde, Porto, Portugal

E-mail address: [joaopfmorais@gmail.com](mailto:joaopfmorais@gmail.com)

(J.P. Morais).

**Aim:** In this study we intent to verify if bupivacaine treatment can be used to enhance the repair of the lesioned urethral sphincter in rat.

**Introduction:** Stress urinary incontinence (SUI) is a major and frequent urinary dysfunction. It has been associated with external urethral sphincter (EUS) weakness due to several causes. Among them, ischemia and nerve lesion frequently associated with child-birth. The current treatments are mainly surgical but are far from being satisfactory. The local anesthetic bupivacaine is known to exert myotoxic action, followed by muscle regeneration with increased strength. This effect was already used in ocular muscles to treat strabismus. In the present study we evaluated the effect of bupivacaine application in the recovery of the damaged EUS.

**Methods:** A lesion of the external urethral sphincter (urethrol-ysis) was performed in adult female Wistar rats using established protocols. Two weeks after the lesion, the animals were injected in the EUS with 0.4 ml of 0.5% bupivacaine. Ten days later, the whole urethra was removed, fixed and sectioned in paraffin wax. Sections

were stained with hematoxylin and eosin, Masson's trichrome and immunoreacted for markers of striated and smooth muscle (sarcomeric actin and smooth muscle actin, respectively).

**Results:** Two weeks after urethrolisis, a marked reduction of muscle fibers in the EUS was detected. A recovery was evident in lesioned, bupivacaine injected animals when compared with lesioned and saline-injected controls.

**Conclusion:** Our data show that bupivacaine application in the lesioned external urethral sphincter accelerates its recovery. This finding opens a therapeutic opportunity to treat stress urinary incontinence.

**Acknowledgements:** This study has been funded by FEDER - Fundo Europeu de Desenvolvimento Regional funds through the COMPETE 2020 - Operacional Programme for Competitiveness and Internationalisation (POCI), Portugal 2020, and by Portuguese funds through FCT/Ministério da Ciência, Tecnologia e Ensino Superior in the framework of the project "Institute for Research and Innovation in Health Sciences" (POCI-01-0145-FEDER-007274).

<http://dx.doi.org/10.1016/j.pbj.2017.07.068>

## PS074

### In vivo and in silico study of allicin as a stroke prevention



Alfryan Janardhana\*, M. Naufal Al Hasan,  
N. Edvin Prawira

DR. dr. Yuyun Yoeniwati P.W. M.kes., Sp Rad (K),  
Indonesia

E-mail address: [ryanjanardhana@gmail.com](mailto:ryanjanardhana@gmail.com)  
(A. Janardhana).

**Aim:** To prove allicin effect to prevent stroke in insilico and invivo method.

**Introduction:** Stroke is a disease that can cause permanent disability, even death. Atherosclerosis is one of the cause of stroke. One way to prevent stroke is to treat atherosclerosis using allicin. Allicin works by inhibiting Integrin Alpha Beta-3 and ApoE proteins. Therefore, allicin can be considered as an alternative in stroke prevention.

**Methods:** This is true experimental study with two methods. Allicin was taken from pubchem, while ApoE with code (1EA8), Integrin Alpha Beta-3 with code (1JX5) taken from protein data bank. Afterwards, the ligands and macromolecule were docked with Pyrx. Analysis was done using Discovery Studio. Pharmacokinetic study, allicin compounds were analyzed with ACD/I-Lab. During invivo study, rats were induced with high fat diet for 8 weeks and were given allicin with dose 5, 10, 20 mg/kg BW during 6 weeks. Rat blood, carotid artery, and brain were analyzed for lipid profile, foam cells in blood vessels, and immunohistochemically to see BDNF.

**Results:** Pharmacokinetic results showed that allicin has oral bioavailability above 70%, distributed through lipoproteins and a few albumins. Allicin can penetrate through membrane and cytoplasm, affecting its target. Pharmacodynamically, allicin can bind to active site of ApoE on 149 leucine, and to active site of ApoE on 173 serine. Allicin bound with active site of ApoE will increase ApoE expression, thus lowering lipid profile except HDL. Meanwhile, allicin bound with active site of Integrin Alpha Beta-3 blocked platelet aggregation. Decreasing Integrin Alpha Beta-3 was proven by invivo results where foam cells were decreasing. These events caused a decrease foam cell in common artery, causing no brain hypoxia and increased BDNF. Invivo test showed a decrease in foam cells on 10 mg/kg BW. On the contrary, the brain showed an increase in BDNF amount on 20 mg/kg BW.

**Conclusion:** Based on insilico and invivo studies, allicin can be considered as a preventive treatment to stroke by inhibiting atherosclerosis development by increasing ApoE, lowering Integrin Alpha Beta-3 protein, and increasing BDNF.<sup>1-5</sup>

## References

1. American Heart Association. Atherosclerosis; 2014 [http://www.heart.org/HEARTORG/Conditions/Cholesterol/WhyCholesterolMatters/Atherosclerosis\\_UCM\\_305564\\_Article.jsp#.WAixhZN95sM](http://www.heart.org/HEARTORG/Conditions/Cholesterol/WhyCholesterolMatters/Atherosclerosis_UCM_305564_Article.jsp#.WAixhZN95sM)
2. Goldstein, et al. History of discovery: the LDL receptor. NIH Public Access. 2009;29:431–8.
3. Xiao-Hua, et al. Foam cell in atherosclerosis. Elsevier; 2013.
4. Freeman WM, Patel KM, Brucklacher RM, Lull ME, Erwin M, Morgan D, et al. Neuropsychopharmacology. 2008;33:1807–17.
5. Aruoma OI, Sun B, Fujii H, Neergheen VS, Bahorun T, Kang KS, et al. Low molecular proanthocyanidin dietary biofactor Oligonol: its modulation of oxidative stress, bioefficacy, neuroprotection, food application and chemoprevention potentials. Biofactors. 2006;27:245–65.

<http://dx.doi.org/10.1016/j.pbj.2017.07.069>

## PS148

### Anxiety-like behavior in elevated plus maze upon sleep fragmentation of one light phase of rats' circadian cycle



A. Leković\*, A. Ademović, Ž. Grubač, N. Šutulović,  
B. Knežević, M. Novaković

Institute of Medical Physiology "Richard Burian",  
Belgrade University School of Medicine, Serbia  
E-mail address: [aleksa.lekovic@gmail.com](mailto:aleksa.lekovic@gmail.com)  
(A. Leković).

**Aim:** To assess anxiety-like behavior in rats after twelve hours of sleep fragmentation during the light phase of sleep-wake cycle, using the elevated plus maze test.

**Introduction:** Hallmark of sleep fragmentation is set of frequent, brief arousals, which modulates sleep architecture without significant diminishment of total sleep time. Anxiety is recognized as comorbidity in numerous disorders, including some of those related to sleep quality. Sleep fragmentation may be appropriate model of sleep alteration pattern in some disorders, yet its effects on behavioral alterations have not been broadly investigated.

**Methods:** Sleep fragmentation was achieved by treadmill method lasting for 12 h, during the light phase of the day (starting at 8 AM). Wistar albino male rats were randomly divided into: sleep fragmentation group (SF,  $n=8$ , treadmill programmed to alternately work 30 s ON and 90 s OFF every 2 min); activity group (AC,  $n=8$ , treadmill programmed to alternately work 10 min ON and 30 min OFF); and treadmill control group (TC,  $n=8$ , rats stayed in the treadmill set to OFF mode and conditions equivalent to cages). Immediately after the sleep fragmentation regimen, elevated plus maze test was performed. To assess anxiety-like behavior, we measured time spent in the open arms, as well as number of transitions between open and closed arms.

**Results:** SF group spent significantly less time in the open arms compared to both, AC and TC group ( $p < 0.001$ ). SF rats also had significantly less number of transitions between closed and open arms of the maze, compared to AC and TC group ( $p < 0.05$ ). Moreover, no significant difference was observed in any of the measured parameters between TC and AC group.

**Conclusion:** The results of our study indicate that acute 12-h sleep fragmentation induced anxiety-like behavior in rats in elevated plus maze. Further research should help us better understand impact of this phenomenon on psychiatric disorders.

<http://dx.doi.org/10.1016/j.pbj.2017.07.070>

## PS082

**Pain and bladder dysfunction in an animal model of multiple sclerosis**

H. Cavaleiro<sup>1,\*</sup>, R. Silva<sup>1</sup>, R. Oliveira<sup>1,2,3</sup>,  
A. Coelho<sup>1,2,3</sup>, F. Cruz<sup>2,3,4</sup>, C.D. Cruz<sup>1,2,3</sup>

<sup>1</sup> Department Biomedicine – Experimental Biology Unit, Faculty of Medicine, Porto, Portugal

<sup>2</sup> Translational NeuroUrology Group, Instituto de Investigação e Inovação em Saúde, Porto, Portugal

<sup>3</sup> Instituto de Biologia Molecular e Celular, Porto, Portugal

<sup>4</sup> Dept. of Urology Hospital São João, Porto, Portugal  
E-mail address: [helenacavaleiro@hotmail.com](mailto:helenacavaleiro@hotmail.com)  
(H. Cavaleiro).

**Aim:** Here, we investigated if MS-induced pain and bladder dysfunction can be attenuated by TRPV1 desensitization with RTX.

**Introduction:** Multiple sclerosis (MS) is the most prevalent neurological disorder in young people, causing irreversible disability and producing substantial economic and social impact. Among the most incapacitating symptoms, neuropathic pain and bladder dysfunction are reported by the majority of patients. The transient receptor potential vanilloid 1 (TRPV1) is a receptor described to have an important role in neuropathic pain, bladder dysfunction and inflammation. TRPV1 desensitization with agonists, such as resiniferatoxin (RTX), has been shown to improve bladder function and reduce behavioural signs of pain in various animal models of disease. In the context of MS, a recent study showed that TRPV1 knockout mice were protected from disease progressions, presenting delayed disease onset, myelin preservation and reduced clinical scores.

**Methods:** Experimental Auto-immune Encephalitis (EAE) was induced by a single injection in the flank of a solution of myelin basic protein (MBP) in Complete Freund's adjuvant (CFA). Behavioural tests were performed to evaluate symptoms. One month after MS-induction, animals were anesthetized and cystometries performed. Two other groups of MS animals received intrathecal RTX or vehicle and also submitted to behavioural tests and cystometries. At end of experiments, tissue was collected and processed.

**Results:** EAE rats developed neuropathic pain, as shown by the presence of mechanical allodynia and hypersensitivity to thermal stimuli. Cystometries performed at this time point showed signs of neurogenic detrusor overactivity. These clinical signs were accompanied by decreased spinal expression of MBP and increased activity of astrocytes and microglia. Preliminary observations suggest that intrathecal RTX improved cutaneous hypersensitivity and bladder function. These results suggest that TRPV1 might be involved in pain bladder dysfunction accompanying MS and that its modulation could have therapeutic relevance.

**Conclusion:** These results suggest that TRPV1 might be involved in pain bladder dysfunction accompanying MS and that its modulation could have therapeutic relevance.

<http://dx.doi.org/10.1016/j.pbj.2017.07.071>

## PS227

**Neurogenesis in a rat model of sporadic Alzheimer's disease**

Sara L. Paulo<sup>1,2,\*</sup>, Rui S. Rodrigues<sup>1,2</sup>,  
Liana Shvachiy<sup>3</sup>, Filipa F. Ribeiro<sup>1,2</sup>,  
Susana Solá<sup>4</sup>, Ana M. Sebastião<sup>1,2</sup>, Sara Xapelli<sup>1,2</sup>

<sup>1</sup> Instituto de Farmacologia e Neurociências, Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal

<sup>2</sup> Instituto de Medicina Molecular (iMM), Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal

<sup>3</sup> Cardiovascular Autonomic Function (CAF) lab, Cardiovascular Center of the University of Lisbon (CCUL), Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal

<sup>4</sup> Research Institute for Medicines (iMed.Ulisboa), Faculdade de Farmácia, Universidade de Lisboa, Lisboa, Portugal

E-mail address: [sara.lnp@gmail.com](mailto:sara.lnp@gmail.com) (S.L. Paulo).

**Aim:** Characterize adult hippocampal neurogenesis in a rat model of the initial stages of sporadic Alzheimer's disease (AD).

**Introduction:** Sporadic late-onset AD is the most common cause of dementia, that can be characterized by a progressive cognitive decline, with a noteworthy episodic long-term memory impairment at early stages, accompanied by an excess accumulation of amyloid beta (A $\beta$ ) peptide in the brain. Present treatment options are very limited, so understanding AD pathophysiology is essential for exploring efficient therapies. Adult hippocampal neurogenesis is thought to play a crucial role in hippocampus-dependent cognitive abilities, namely learning/memory, although how this process is modulated in AD remains unclear.

**Methods:** An A $\beta$ 1–42 peptide solution was intracerebroventricularly injected into the rats' lateral ventricle (the same volume of vehicle was injected to controls). Moreover, rats were injected with 5-bromo-2'-deoxyuridine (BrdU) intraperitoneally to study cell proliferation and differentiation. Two weeks after A $\beta$ 1–42 injection, the open field (OF) test and the novel object recognition (NOR) test were performed. Further behaviour tests are currently being performed, including the elevated plus maze (EPM), the Y-maze forced alternation test, and the Morris water maze (MWM) test. Focusing on the dentate gyrus, immunohistochemical analysis is presently being performed to investigate cell proliferation, neuronal differentiation and neuroblast/neuron morphology. Additionally, the presence of A $\beta$ 1–42 monomers and oligomers will be assessed by western-blot and the eventual occurrence of A $\beta$ 1–42 aggregates by histology.

**Results:** Our results show that the A $\beta$ 1–42 injection did not affect locomotor activity, as assessed by the OF test. Furthermore, this injection did not affect exploratory drive or episodic long-term memory performance, as indicated by the NOR test.

**Conclusion:** Since the NOR test is dependent from several brain regions besides the hippocampus that might not be affected in our model, additional behaviour tests as well as cellular and molecular analysis are needed to further characterize this model.

<http://dx.doi.org/10.1016/j.pbj.2017.07.072>

## PS086

**Hydroalcoholic extract of *Dorema aucheri* leaves prevents weakening of the brain antioxidant defense system and inhibits oxidative damage in rat model of ischemic stroke**

Mohammad Ehsan Bayatpoor<sup>1,\*</sup>,  
Javad Rasouli Vani<sup>2</sup>,  
Mohammad Taghi Mohammadi<sup>3</sup>

<sup>1</sup> Baqiyatallah Research Center for Gastroenterology and Liver Diseases (BRCGL), Baqiyatallah University of Medical Sciences, Tehran, IR Iran

<sup>2</sup> Department of Physiology and Biophysics, School of Medicine, Baqiyatallah University of Medical Sciences, Tehran, Iran

<sup>3</sup> Middle East Liver Disease (MELD) Center, Tehran, IR Iran

E-mail address: [ehsanbayatpoor@yahoo.com](mailto:ehsanbayatpoor@yahoo.com) (M.E. Bayatpoor).

**Aim:** The aim of this article is Hydroalcoholic extract of *Dorema aucheri* leaves prevents weakening of the brain antioxidant defense system and inhibits oxidative damage in rat model of ischemic stroke

**Introduction:** The production of free radicals is the principal mechanism of brain injury in ischemic stroke. The present study tried to identify whether pretreatment with hydroalcoholic extract of *Dorema aucheri* (DA) leaves potentiates the brain antioxidant system and decreases brain infarction and oxidative damage during cerebral ischemia–reperfusion.

**Methods:** Three groups of rats were randomly selected (each group;  $n = 12$ ); sham, control ischemic and ischemic pretreatment groups. Treated rats received freshly hydroalcoholic extract of DA (200 mg/kg/day) for 14 days. Then, cerebral ischemia–reperfusion was achieved by 90 minutes middle cerebral artery (MCA) occlusion followed by 24 h reperfusion. Infarct volume and contents of malondialdehyde (MDA), glutathione and nitrate (NOx) as well as superoxide dismutase (SOD) and catalase activities were assessed after 24 h reperfusion

**Results:** The contents of MDA and nitrate significantly increased in the ischemic hemispheres by 34% and 14%, respectively. Brain ischemia decreased the glutathione content (20%) and activities of catalase (38%) and SOD (14%) in ischemic hemispheres compared to sham rats. Treatment with DA before MCA occlusion significantly decreased the infarction in cortex and striatum by 63% and 75%, respectively, compared to control. DA considerably reduced the contents of MDA and nitrate in ischemic hemispheres by 28% and 11%, respectively, compared to control rats. Treatment with DA also increased the glutathione content (7%) and activities of catalase (46%) and SOD (16%) of ischemic hemispheres.

**Conclusion:** The present study revealed that pretreatment with hydroalcoholic extract of DA leaves prevents weakening of the brain antioxidant defense system and decreases the brain damage during cerebral ischemia–reperfusion.

<http://dx.doi.org/10.1016/j.pbj.2017.07.073>

## PS150

### The Levator Auris Longus (LAL) muscle as an accessible system to study the effects of Botulinum Toxins in vivo

Torrado Marília<sup>1,\*</sup>, Cruz Célia Duarte<sup>1,2,3</sup>, Avelino António<sup>1,2,3</sup>

<sup>1</sup> Departamento de Biomedicina – Unidade de Biologia Experimental, Faculdade de Medicina da Universidade do Porto, Portugal

<sup>2</sup> Translational NeuroUrology, IBMC – Instituto de Biologia Molecular e Celular, Universidade do Porto, Portugal

<sup>3</sup> Instituto de Investigação e Inovação em Saúde - i3S, Universidade do Porto, Portugal

E-mail address: [marilia.torrado@gmail.com](mailto:marilia.torrado@gmail.com) (T. Marília).

**Aim:** In the present work, we aimed to find a reproducible model to study the effects of Botulinum neurotoxins (BoNTs), that allowed a widespread visualization of the intoxicated nerve terminals.

**Introduction:** Despite the well-established successful use of BoNTs to treat a variety of human conditions, their mechanism of action is still not fully understood. Thus, there is an emergent need of new and accurate models to study the effects of BoNTs. However, considering their potential lethality, it is challenging to find reproducible models to study the local application of BoNTs in living animals that allow a widespread visualization of the intoxicated nerve terminals.

In these work, we studied the innervation pattern and the effect of BoNTs in a group of small subcutaneous cranial muscles that are responsible for moving the pinna in rodents. Although all are easily accessible and manipulated, we focused on the levator auris longus (LAL).

**Methods:** Animals were injected subcutaneously with the indicated doses of BoNT/A, in the cranial muscles area. Muscles were then dissected and prepared for wholemount staining for Synapsin-I, cleaved SNAP-25 (synaptosome-associated protein of 25 kDa) and  $\beta$ 3-tubulin.

**Results:** Detection of cleaved SNAP-25, the end-product of the catalytic action of BoNT/A, was possible even with injections as low as 0.1 ng. Mapping of the injected muscle showed the effect of BoNT/A in the majority of the endplate population. Also, seven days after BoNT/A injection, a sprouting process was evident, a landmark of regeneration.

**Conclusion:** BoNTs delivery to the LAL is a sensitive, simple and reproducible model to study the mechanisms of action of these toxins as it allowed the evaluation of BoNT/A effects throughout the entire muscle, without sampling bias. Thus, we forward that the LAL manipulation may constitute an excellent model to clarify the mechanisms of action of BoNTs in the neuromuscular system.

**Acknowledgements:** This study has been funded by FEDER - Fundo Europeu de Desenvolvimento Regional funds through the COMPETE 2020 - Operational Program for Competitiveness and Internationalization (POCI), Portugal 2020, and by Portuguese funds through FCT - Fundação para a Ciência e a Tecnologia/Ministério da Ciência, Tecnologia e Ensino Superior in the framework of the project “Institute for Research and Innovation in Health Sciences” (POCI-01-0145-FEDER-007274).

<http://dx.doi.org/10.1016/j.pbj.2017.07.074>

## PS188

### NLRP3 inflammasome as a potential target to reduce epileptic-like activity



L. Ribeiro-Rodrigues\*, D.M. Rombo, A.M. Sebastião, C.A. Valente

Faculdade de Medicina, Universidade de Lisboa e Instituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, Portugal  
E-mail address: [leonor\\_rr@hotmail.com](mailto:leonor_rr@hotmail.com) (L. Ribeiro-Rodrigues).

**Aim:** Decipher how inflammation drives epilepsy and how NLRP3 targeting impacts epileptic-like activity.

**Introduction:** Epilepsy is one of the most common neurological diseases in worldwide. Inflammation was linked to the presence of inflammasomes, cytosolic multiprotein complexes, which promote the release of proinflammatory cytokines, namely IL-1 $\beta$ . Although a feedback loop has been described between inflammation and epilepsy, the role of inflammasomes in epilepsy is still unknown. NLRP3 is the most studied inflammasome,<sup>1</sup> activated by a two-signal process: 1) a priming signal (as lipopolysaccharides – LPS), which enhances the expression of NLRP3 and pro-IL-1 $\beta$ ; and 2)

an activating signal (as ATP), which promotes the formation of the complex.

**Methods:** Organotypic slices were used to assess the interplay between inflammation and epilepsy. Slices were exposed to different concentrations of LPS (5, 10 and 20 ng/mL), either alone or in the presence of ATP (1 mM). LPS-induced inflammation was characterized using molecular-based assays, such as ELISA to quantify IL-1 $\beta$ , CBA to measure TNF- $\alpha$ , and western blot to assess the expression of Iba-1, GFAP, NLRP3/ASC, and  $\alpha$ II-Spectrin. Field potential recordings were used to evaluate the epileptic-like activity of the slices and the effect of MCC950, a NLRP3 selective inhibitor,<sup>2</sup> was assessed.

**Results:** Results obtained by ELISA showed a significant increase in IL-1 $\beta$  concentration in slices exposed to 10 ng/ml LPS/1 mM ATP. TNF- $\alpha$ , assessed by CBA, was also significantly increased in this condition, corroborating the inflammatory phenotype. No changes in NLRP3 expression were observed by immunoblot analysis, but ASC, one component of the inflammasome, showed a decreased expression in LPS/ATP exposed slices, suggestive of its binding to NLRP3 and thus to complex formation.

Furthermore, epileptic-like activity, measured by field potential recordings, was blocked by MCC950 (10  $\mu$ M).

**Conclusion:** We demonstrate that LPS induces an inflammatory phenotype in organotypic slices. NLRP3 blockade eliminated the epileptic-like activity of the slices.

## References

- Walsh JG, Muruve DA, Power C. Inflammasomes in the CNS. *Nat Rev Neurosci.* 2014;15:84–97.
- Coll RC, Robertson AAB, Chae JJ, et al. A small-molecule inhibitor of the NLRP3 inflammasome for the treatment of inflammatory diseases. *Nat Med.* 2015;21:248–55.

<http://dx.doi.org/10.1016/j.pbj.2017.07.075>

## PS110

### Study of the modulatory CNS regions in the visual circuit Retina-Superior Colliculus-Lateral Posterior nucleus triggering freezing behavior



Ares Sellés Rius

*NERF (Neuro-Electronics Research Flanders), Karl Farrow's Laboratory, Belgium*  
E-mail address: [aresseri.28@gmail.com](mailto:aresseri.28@gmail.com).

**Aim:** The goal is to understand the neuronal networks organization from the sensory input to the freezing behavior through the identification of modulatory brain regions that project to the Superior Colliculus.

**Introduction:** The behavior of an animal can be triggered by signals in its visual environment. Threatening visual stimulus evoked innate defense behaviors as freezing behavior. This project is focused in one visual-guided behavioral circuit that links the retina visual information with the Lateral Posterior thalamic nucleus(LP) via Superior Colliculus(SC).

**Methods:** The experimental approach is based on retrograde viral tracing techniques. Using the stereotaxic surgery, the first injection with a Herpes Simplex Virus expressing TVA receptor and glycoprotein G was done at LP. After 21 days, the second injection was done at the SC with a Rabies Virus coated by EnvA and lacking of glycoprotein G. The combination of these viruses allowed the restriction of the viral tracing to the circuit of interest. Subsequently, the experimental procedure continued perfusing the mouse, slicing the brain and staining it. Finally, the slices were scanned using the fluorescent confocal microscope.

**Results:** The resulting images presented labeled cells in all brain areas that sent inputs to collicular neurons that are projecting to LP. The main nuclei identified were the Periaqueductal gray, the primary visual cortex and the Substantia nigra, suggesting their modulatory role in freezing responses.

**Conclusion:** The main areas labeled are sending excitatory projections to SC to reinforce the freezing behavior. Also, Ntsrl-GN209-Cre mice used in combination with flox-HSV for the first injection restricted more the viral tracing, specifically to the Ntsrl+ Wild-field neurons of SC which were already known that project to LP. The results were not completely consistent with the non-flox-HSV injections but the main nuclei named above were also labeled. These results suggest that the flox-HSV is necessary to exclude nonspecific labeling of projections from SC-LGN.

**Acknowledgements:** The exposed project was done in Karl Farrow's laboratory, at NERF (Neuro-Electronics Research Flanders) in Leuven, Belgium. It was a Bachelor's thesis, supported by both Katholieke Universiteit Leuven and University of Barcelona.

<http://dx.doi.org/10.1016/j.pbj.2017.07.076>

## PS048

### The influence of antipsychotics therapy and sociodemographic characteristics on cognitive performances in acute phase of schizophrenia



Milica Erdevički\*, Nataša Jovičić

*Medical Faculty of Novi Sad*

E-mail address: [mimaerdevicki1@gmail.com](mailto:mimaerdevicki1@gmail.com)

(M. Erdevički).

**Aim:** The main purpose of this research was to examine the influence of sociodemographic characteristics (gender, age, level of education, heredity, alcohol and psychoactive substances), and the effect of different therapies on cognitive capabilities of patients diagnosed with schizophrenia.

**Introduction:** Schizophrenia, as one of the most common psychiatric diseases, is characterized by generalized cognitive damage with various degrees and in all domains of cognitive functioning. Cognitive dysfunction is one of the main causes of poor social and professional functioning for patients with schizophrenia.

**Methods:** The research involved 50 patients with acute phases of schizophrenia from the Psychiatric Clinic in Novi Sad. The primary instrument for the research was the standardized test for examination of cognitive impairments, Mini-Mental Scale Examination (MMSE).

**Results:** Acquired data correlated with MMSE score, noting the degree of cognitive impairments in patients, particularly significant with relation to age and duration of illness. Gender, level of education and type of used antipsychotics were not significantly correlated with MMSE score.

**Conclusion:** During this research it is found that aging and longer illness duration bear significant correlation to higher levels of cognitive impairment.

<http://dx.doi.org/10.1016/j.pbj.2017.07.077>

## PS190

### Voluntary inhibition of saccadic eye movements: EEG study



A. Fedotova\*, M. Slavutskaya

*Lomonosov Moscow State University, Department of Higher Nervous Activity, Russia*

E-mail address: [fedotova.brain@gmail.com](mailto:fedotova.brain@gmail.com)  
(A. Fedotova).

**Aim:** The aim of our study was to find out EEG markers of inhibitory control in human.

**Introduction:** The voluntary inhibition is an important component of cognitive control. It is strong in healthy adults and weak in people with schizophrenia. The cortical mechanisms of inhibition are associated with event-related potentials (ERPs). In the case of a saccadic response some new EEG correlates of inhibition could be found.

**Methods:** Sixteen healthy right-handed subjects (18–22 years) participated in the study. We used a modified “Go/No go delay” paradigm with long interstimulus interval (2800–3000 ms). The task involved two types of target stimuli (“Go”, “No go”) with 50% probability. EEG and saccades were recorded simultaneously. ERPs were determined by means of coherent averaging relative to target stimulus onset. The EEG brain mapping was used to depict spatial dynamics of P1.

**Results:** P1 peak latency was 90–140 ms and tended to increase in cases of inhibition (by  $6 \pm 0.5$  ms,  $p < 0.05$ ). In the “No go” situation P1 amplitude was significantly lower than that in case of “Go” stimulus presentation (by  $3.3 \pm 0.7$  mV,  $p < 0.05$ ). Regardless of the place where “No go” stimulus appeared, P1 amplitude was significantly higher on the right hemisphere, that is known to be the dominant one for inhibitory control. The EEG mapping data demonstrate the “bottom-up” spreading of P1 foci in “No go” conditions. It also indicates inhibitory processes.

**Conclusion:** The spatiotemporal parameters of P1 component in “Go/No go delay” paradigm reflect inhibitory processes. Therefore, P1 can be used as EEG marker of inhibitory violations in the clinical research. Our current research involves as subjects the patients with schizophrenia and ultra-high risk patients, as they demonstrate weakened the inhibitory processes. The data would contribute to the reliable diagnostics of schizophrenia at its early stages and to the plausible correction of cognitive impairments.

**Acknowledgements:** I would like to express my sincere gratitude to my supervisor D.Sc. in Biology Maria V. Slavutskaya.

<http://dx.doi.org/10.1016/j.pbj.2017.07.078>

### PS135

#### Factors influencing the outcome of endovascular embolization of anterior communicating artery aneurysms



I. Kucybala\*, K. Krupa, J. Polak, J. Wnuk

Jagiellonian University Medical College, Cracow,  
Poland

E-mail address: [iwona.kucybala@gmail.com](mailto:iwona.kucybala@gmail.com)  
(I. Kucybala).

**Aim:** The aim of the study was to assess the influence of morphologic parameters of anterior communicating artery aneurysms and the method of embolization on the success rate of procedure.

**Introduction:** Endovascular embolization of anterior communicating artery aneurysms is currently considered as primary management tool and the improvement of procedural success rate is crucial.

**Methods:** Treatment process of 109 patients undergoing endovascular embolization of anterior communicating artery aneurysm was retrospectively analysed. All procedures were performed between August 2006 and December 2016 in Department of Interventional Radiology of University Hospital in Cracow (Poland).

The mean age of patients was  $56.7 \pm 15.2$  years (range 28–91), 50.5% of patients were female. Used methods of embolization: coiling alone, balloon-assisted coiling, stent-assisted coiling, Y-stenting + coiling. Evaluated morphologic parameters: width of the neck, maximal height, maximal width, shape of aneurysm, dome orientation. The outcome of the procedure was assessed with Raymond–Roy occlusion classification. Data were analysed using chi-square test and Student’s *t*-test. Statistical significance was set at  $p < 0.05$ .

**Results:** Coiling alone significantly improved outcome of embolization considered as better score in Raymond–Roy occlusion classification, compared to other methods ( $1.4 \pm 0.5$  vs.  $1.6 \pm 0.7$ ;  $p = 0.034$ ). In case of irregular aneurysms (85.7% vs. 34.6% (regular aneurysms);  $p = 0.025$ ; OR = 2.615) and those with posterior orientation of the dome (76.9% vs. 36.5% (anterior orientation);  $p = 0.005$ ; OR = 5.810) incomplete embolization (Raymond–Roy class II and III) was significantly more frequent. Within the group of discharged patients, only 33.3% undergone control radiologic examination – 40.7% conventional angiography, 59.3% MR angiography. In that group, 81.5% of aneurysms had better or the same class in Raymond–Roy classification and 18.5% had worse outcome. We did not discovered any statistically significant factor contributing to that phenomenon.

**Conclusion:** Coiling alone is the most efficient method in terms of the aneurysm occlusion rate. Irregular shape of the aneurysm and posterior orientation of the dome significantly hinder the embolization of aneurysm.

<http://dx.doi.org/10.1016/j.pbj.2017.07.079>

### PS097

#### Antidepressive potential of aqueous extract of common vervain (*V. officinalis* L. Verbenaceae) and molecular docking studies of its main components as potential antidepressive agents



N. Lasica<sup>1,\*</sup>, V. Raicevic<sup>2</sup>

<sup>1</sup> Department of Pharmacology, Toxicology and Clinical Pharmacology, Faculty of Medicine, University of Novi Sad, Serbia

<sup>2</sup> Department of Chemistry, Biochemistry and Environmental Protection, Faculty of Sciences, University of Novi Sad, Serbia

E-mail address: [nebojsa.lasica@gmail.com](mailto:nebojsa.lasica@gmail.com)  
(N. Lasica).

**Aim:** Assessment of antidepressive effect of aqueous extract (AE) of common vervain and its main constituent – verbascoside and elucidation of underlying mechanism of action.

**Introduction:** Common vervain is a plant used in traditional medicine. Its AE contains a vast number of compounds, hence its significant pharmacological potential.

The monoamine hypothesis is the central theory of depression, and a majority of conventional antidepressants act on the monoaminergic system.

**Methods:** Experiments were conducted on Swiss albino sexually mature male mice. There were 6–8 animals in each of 5 subgroups (imipramine; fluoxetine; two different doses of AE – AE I, II; and VS;). Forced Swimming Test (FST) and Tail Suspension Test (TST) were used to assess the antidepressive effect.

Molecular docking experiments were performed using the programme AutoDock 4.2, with 3D structures of crystallized proteins from the PDB database and 3D structures of ligands generated by the software Avogadro 2.0.8.0.

**Results:** Immobilisation time (IT) in FST after the administration of imipramine was shorter than the control, same as for subgroups treated with AE I, II and VS. In the subgroup treated with fluoxetine, IT in TST was shorter than the control time, and the same was observed in subgroups treated with AE I, II and VS.

Significant binding energies were found for Serotonin Reuptake Transporter (SERT) and verbenalin (–7.20 kcal/mol) and verbasco-side (–6.61 kcal/mol), and for the Leucine Transporter (LeuT), the homologue of the noradrenaline reuptake transporter, and verbenalin (–6.27 kcal/mol) and caffeic acid (–5.85 kcal/mol).

**Conclusion:** In both pharmacodynamic tests the antidepressive effect of AE and VS has been confirmed. Verbenalin and verbasco-side binding energies and poses in interaction with SERT were similar to those of paroxetine. For LeuT, verbenalin showed both a similar binding energy and pose to that of imipramine, whereas caffeic acid showed only a similar binding energy.<sup>1–4</sup>

## References

1. Abascal K, Yarnell E. Nervine herbs for treating anxiety. *Altern Complement Therap.* 2004;10:309–15.
2. Akanmu MA, Honda K, Inoue S. Hypnotic effects of total aqueous extracts of *Vervain hastata* (Verbenaceae) in rats. *Psychiatry Clin Neurosci.* 2002;56:309–10.
3. Belmaker RH, Galila Agam. Major depressive disorder. *N Engl J Med.* 2008;358:55–68.
4. Makino Y, Kondo S, Nishimura Y, Tsukamoto Y, Huang ZL, Urade Y. Hastatoside and verbenalin are sleep-promoting components in *Verbena officinalis*. *Sleep Biol Rhyth.* 2009;7:211–7.

<http://dx.doi.org/10.1016/j.pbj.2017.07.080>

## PS213

### Thermal denaturation profiles of proteome and blood serum of rats with drug-induced dementia. A DSC study



N. Nizamova\*, S. Abarova, L. Traikov, R. Koynova, B. Tenchov

*Dept Medical Physics and Biophysics, Medical University – Sofia, 1431 Sofia, Bulgaria*  
E-mail address: [nezi.nizamova@gmail.com](mailto:nezi.nizamova@gmail.com)  
(N. Nizamova).

**Aim:** The aim of this study is to evaluate the effect of scopolamine on different brain segments using DSC.

**Introduction:** In this work, probes from different brain segments of rats with drug-induced dementia were characterized by differential scanning calorimetry (DSC) and their thermodynamic properties were determined.

**Methods:** Male Wistar rats were injected with scopolamine for 14 consecutive days in order to induce drug model of dementia. After being decapitated, their brains were divided into the following segments: telencephalon, mesencephalon and cerebellum. Afterwards, the brain supernatants of the latter 3 segments were examined by DSC and compared with the controls.

**Results:** The DSC measurements revealed large differences between the denaturation profiles of rat brain supernatants and blood serum. The thermograms of brain tissues displayed clearly expressed low-temperature exothermic transitions with peaks in the range 35–45 °C which are missing in blood serum samples. There were differences between the thermograms of the separate brain segments as well. The thermodynamic parameters of the denaturation profiles were also determined.

**Conclusion:** These measurements show that DSC is an appropriate method with great potential for detection and characterization of the changes taking place at molecular level in different tissues, especially in brain tissues affected by neurodegenerative disorders.

**Acknowledgements:** This research was made with the support of Science Fund at Medical University-Sofia, Bulgaria, Project 8-C/2016 and National Science Fund Project DN03/13/2016.

<http://dx.doi.org/10.1016/j.pbj.2017.07.081>

## PS231

### Effects of Vitamin D on the expression of markers of principal neurons, interneurons and astrocytes in cerebral cortex and hippocampus in gerbils exposed to transient global cerebral ischemia



M. Malinic\*, G. Jevtic Dozudic

*Institute of Clinical and Medical Biochemistry, Faculty of Medicine, University of Belgrade*  
E-mail address: [marija.malinic@gmail.com](mailto:marija.malinic@gmail.com)  
(M. Malinic).

**Aim:** Examination of the effects of vitamin D pretreatment on the expression of markers of principal neurons (NeuN), inhibitory interneurons (PV) and astrocytes (GFAP) in cerebral cortex and hippocampus in gerbils who were exposed to transient global cerebral ischemia.

**Introduction:** Brain ischemia may cause serious damage to the cells in the central nervous system. Vitamin D has an important role in brain injury treatment due to its neuroprotective effects.

**Methods:** Gerbils were divided in 5 groups: control group; two groups that underwent ischemia and then reperfusion for three (I/R3d) and seven days (I/R7d) and two groups that were treated with vitamin D before I/R (vitD + I/R3d and vitD + I/R7d). Complete blood supply to the brain was cut off for 10 minutes and reperfusion lasted 3 and 7 days. They were daily treated with vitamin D for 7 days prior ischemia. Expression of proteins was detected using Western blot.

**Results:** No changes were detected in expression of NeuN markers in cortex of experimental groups, while there was increase in expression in hippocampus in groups I/R7d and vitD + I/R7d in comparison to the control group and group vitD + I/R3d. Expression of PV in cortex was significantly reduced in group I/R7d in comparison to group I/R3d, whereas in hippocampus the expression was significantly higher in group vitD + I/R3d than in group I/R3d. Expression of GFAP has significantly risen in all groups in comparison to the control group whereas in hippocampus there was a rise in groups vitD + I/R3d, I/R7d and vitD + I/R7d in comparison to the control group. There was also a rise of GFAP expression in groups treated with vitamin D (vitD + I/R3d and vitD + I/R7d) in comparison to those that have not been treated (I/R3d, I/R7d).

**Conclusion:** Vitamin D has positive effect on astrocytes in both structures of gerbils that underwent global cerebral ischemia, especially in hippocampal region.

<http://dx.doi.org/10.1016/j.pbj.2017.07.082>

## PS238

### Identification of genetic modifiers of somatic CAG instability in Huntington's Disease by in vivo CRISPR – Cas9 genome editing



A. Azevedo<sup>1,2</sup>, M. Kovalenko<sup>2</sup>, M. Andrew<sup>2</sup>, F. Zhang<sup>3</sup>, J. Lee<sup>2,4</sup>, V. Wheeler<sup>2,4</sup>, R. Mouro Pinto<sup>2,4,\*</sup>

<sup>1</sup> University of Porto

<sup>2</sup> Center for Genomic Medicine, Massachusetts General Hospital

<sup>3</sup> Broad Institute of Harvard and MIT

<sup>4</sup> Harvard Medical School

E-mail address: [rmouropinto@mgh.harvard.edu](mailto:rmouropinto@mgh.harvard.edu)

(R. Mouro Pinto).

**Aim:** To develop an experimental platform for *in vivo* investigation of candidate genetic modifiers of somatic CAG instability in Huntington's disease.

**Introduction:** Huntington's disease (HD) is an autosomal dominant neurodegenerative disorder caused by a CAG repeat expansion within the huntingtin gene (*HTT*).<sup>1</sup> Despite being a monogenic disorder, for which the mutation has been known for some time now, no cure or disease-modifying therapy is available, indicating that novel approaches are critical.

Somatic CAG repeat instability, characteristic of mutant *HTT* alleles, is inversely correlated with patient age of onset and may contribute to HD pathogenesis.<sup>2,3</sup> This phenotype, common to other trinucleotide repeat disorders,<sup>4</sup> was previously shown to be DNA mismatch repair (MMR) dependent.<sup>5</sup> The DNA repair machinery was further implicated as a modifier of HD age of motor onset in a recent genome wide association study, underlining its promise as a relevant disease mechanism that could potentially be therapeutically targeted.<sup>6</sup>

In this study, we are developing a CRISPR/Cas9-based approach that will enable the investigation of candidate genetic modifiers of HD age of onset as potential modifiers of somatic CAG repeat instability in a HD mouse model.

**Methods:** We have developed CRISPR reagents against known and candidate genetic modifiers of somatic CAG instability in Huntington's disease. In preliminary experiments, we treated HD mice with CRISPR reagents against *Mlh1* and investigated the level of gene editing achieved as well as the impact on liver CAG instability.

**Results:** We were able to significantly suppress the CAG expansion process in the liver of HD mice by knocking out the *Mlh1* gene in our *in vivo* CRISPR platform. The efficiency achieved in modifying the instability phenotype makes us very confident that we will be able to test and validate additional candidate modifiers. To that end, we have already validated reagents for efficient knockout of a subset of known and candidate modifier genes and we have developed assays that will allow detailed characterization of gene editing at these sites.

**Conclusion:** We have successfully developed an *in vivo* CRISPR-Cas9-based platform that allows for knocking out genes of interest in the liver of adult mice, and consequently perturb the somatic CAG expansion process. We will next use this tool to test the role that candidate genes might play in that disease-relevant process. While the scope of this project was liver oriented, future work will also be aimed at targeting the striatum which is the main site of HD-related pathology.

## References

- Gusella JF, et al. Genetic modifiers of Huntington's disease. *Mov Disord.* 2014;29:1359–65.
- Kennedy L, et al. Dramatic tissue-specific mutation length increases are an early molecular event in Huntington disease pathogenesis. *Hum Mol Genet.* 2003;12:3359–67.
- Swami M, et al. Somatic expansion of the Huntington's disease CAG repeat in the brain is associated with an earlier age of disease onset. *Hum Mol Genet.* 2009;18:3039–47.
- Schmidt MHM, et al. Disease-associated repeat instability and mismatch repair. *DNA Repair (Amst).* 2015;38:117–26.
- Pinto RM, et al. Mismatch repair genes *Mlh1* and *Mlh3* modify CAG instability in Huntington's disease mice: genome-wide and candidate approaches. *PLoS Genet.* 2013;9:e1003930.
- Genetic Modifiers of Huntington's Disease (GeM-HD) Consortium. Identification of genetic factors that modify clinical onset of Huntington's disease. *Cell.* 2015;162:516–26.

<http://dx.doi.org/10.1016/j.pbj.2017.07.083>

Oncology & Molecular Biology Poster Session  
Sunday, September 17th, 10h00

## PS021

### Regulation of transcription factor MEF2C by RNA binding protein HuR



Z. Anyu<sup>1</sup>, G. Shi<sup>1</sup>, A. Xie<sup>1</sup>, D. Aksoy<sup>2,\*</sup>, S. Dudley<sup>1</sup>

<sup>1</sup> Cardiovascular Research Center, The Warren Alpert Medical School of Brown University, Providence Rhode Island, United States

<sup>2</sup> Marmara University School of Medicine, Istanbul, Turkey

E-mail address: [sdilsadaksoy@gmail.com](mailto:sdilsadaksoy@gmail.com)

(D. Aksoy).

**Aim:** We hypothesized that HuR RNA binding protein regulates MEF2C expression through association with MEF2C mRNA.

**Introduction:** MEF2C is earliest expressed member of the MADS-box super family during heart development. In the postnatal heart, decreased expression of MEF2C has been associated with myotonic dystrophy type 1 (DM1) heart disease. Hu proteins are known to regulate a wide range of gene expression by modulating mRNA's half-lives.

**Methods:** We use Human Fetal Cardiomyocyte cell line RL14. Cells are transfected with Superfect Transfection Reagent(Qiagen). And RNA Isolation performed by using RNeasy Plus Mini Kit. Real Time quantitative PCR (q-PCR) analysis performed using Fast SYBR Green Master Mix.

**Results:** Over expression of HuR in cardiomyocytes derived from primary human fetal ventricle increased MEF2C mRNA 47.3% ( $p = 0.01$ ). Knocking down of HuR by siRNA decreased MEF2C mRNA by 62% ( $p = 0.01$ ). RNA Immunoprecipitation showed HuR associated with MEF2C mRNA.

**Conclusion:** Our results suggest that RNA binding protein HuR associates with MEF2C mRNA in cardiomyocytes. And also HuR positively regulates MEF2C mRNA expression.

<http://dx.doi.org/10.1016/j.pbj.2017.07.084>

## PS024

### The effect of prenatal Vitamin C deficiency on endochondral ossification in guinea pigs



N. Rakočević

Medical Faculty, University of Novi Sad, Serbia

E-mail address: [rakocevicnatali@gmail.com](mailto:rakocevicnatali@gmail.com).

**Aim:** The aim of the research is to investigate the effect of prenatal vitamin C deficiency on endochondral ossification in guinea pigs.

**Introduction:** Vitamin C is an essential nutrient which inter alia enables the synthesis of collagen and therefore endochondral ossification. Throughout years a lot of research has been published investigating the exact role of vitamin C and the impairment developed due to its deficiency. However there is insufficient data about the effect of prenatal deficit of vitamin C on the developing bone structures.

**Methods:** The study encompassed 14 fertilized female albino guinea pigs. Their diet was comprised of vitamin C-free food and ad libitum water enriched with vitamin C. The 10th day of fertilization, experimental group was depleted of vitamin C. Deprivation lasted until the 50th day, after which the females were sacrificed and their fetuses were taken out. Forelegs of fetuses were fixed and dehydrated, after which they were embedded in paraffin and

longitudinal sections were made. The stain used for histology was Alcian&Alizarin.

**Results:** The development of long bones in vitamin C deficient guinea pigs are considerably stagnant. Hyaline cartilage models are significantly shortened. Ossification in the diaphyses of carpal and metacarpal bones are absent, and the organization of the epiphyseal plates is very irregular with the reduction of number of chondrocytes. Moreover, there are numerous haemorrhagic regions and subperichondrial bleeding with separation of perichondrium.

**Conclusion:** Deprivation of vitamin C during inrauterine period disables normal development of long bones. Disorder of hyaline cartilage models was seen, as well as the disorder of ossification.

<http://dx.doi.org/10.1016/j.pbj.2017.07.085>

#### PS060

##### Peculiarities of expression of apoptosis markers in the tissues of primary fallopian tubes carcinoma



Franklin Unawunwa\*, Natalia Hyriavenko, Anna Korobchanska, Mykola Lyndin, Vladyslav Sikora

Sumy State University

E-mail address: [unawunwafranklin@yahoo.com](mailto:unawunwafranklin@yahoo.com) (F. Unawunwa).

**Aim:** immunohistochemical analysis of apoptosis markers in the tissue of PFTC.

**Introduction:** Primary fallopian tubes carcinoma is a rare case among oncological diseases of female genital organs, but the mortality rate is rather high. Nowadays, the prognostic factors of this neoplasia are not fully determined. The data on the p53 and bcl2 proteins expression and their use as prognostic factors in patients with malignant tumors of many locations are contradictory.

**Methods:** the study was conducted on 66 samples of fallopian tubes tumor tissue. To study the apoptosis peculiarities of tumor cells the mouse monoclonal antibodies for bcl-2 (clone 100/D5) and p53 (clone SP5) were used. Mathematic calculations were done using Microsoft Excel 2010 with AtteStat 12.0.5.

**Results:** The high expression of p53 was found in patients of all clinical stages. Mutations of p53 increased with spreading of the neoplastic process. Strong correlation of p53 presence in tumor samples and clinical stage of the disease was determined ( $r=0.77$ ). In contrast to the abovementioned protein the study of bcl-2 showed the moderate negative correlation between this protein and the stage of the disease ( $r=-0.54$ ). Analysis of the dependence of p53 expression with the presence or absence of lymph nodes metastasis showed a direct correlation between the indicators ( $r=0.25$ ). Thus the level of p53 expression in patients with N1 was  $80.6 \pm 2.7\%$  compared with the N0 group ( $29.7 \pm 3.6\%$ ). The stage of neoplasia differentiation is in moderate direct correlation with p53 expression ( $r=0.58$ ) and in inverse with – bcl-2 ( $r=-0.64$ ).

**Conclusion:** Expression of p53 depends on neoplasia spreading and stage of tumor differentiation. The expression of p53 is an independent prognostic marker for N-status and helps to classify the patients into “risk” groups.

**Acknowledgements:** Supervisor: A. M. Romaniuk, prof., doctor of medical sciences, Department of Pathological Anatomy, Medical Institute, Sumy State University.

<http://dx.doi.org/10.1016/j.pbj.2017.07.086>

#### PS064

##### Analysis of combined impact of doxorubicin and menadione on human leukaemia Jurkat T cells



Alexandru Ionut Duta\*, Ioana Teodora Tofolean, Ramona Madalina Babes, Constanta Ganea, Irina Baran

“Carol Davila” University of Medicine and Pharmacy, Department of Biophysics, Bucharest, Romania  
E-mail address: [alexxxduta@yahoo.com](mailto:alexxxduta@yahoo.com) (A.I. Duta).

**Aim:** The anti-proliferative effect and the mechanism of action of doxorubicin(DOX) in combination with menadione(MD) were studied in Jurkat T cells, a model for acute lymphoblastic leukaemia (ALL).

**Introduction:** Doxorubicin is a well-characterized and successful antineoplastic drug commonly used in various cancer treatments, including ALL. Menadione has proven a strong proapoptotic effect in Jurkat cells.<sup>1-3</sup>

**Methods:** Cell cycle, apoptosis/necrosis and the oxidative status were assessed by flow cytometry on propidium iodide, Annexin V-FITC/PI and CM-H2DCFDA/7-AAD labelled cells, respectively.

**Results:** Oxidative stress induced within 4h by MD ( $IC_{50}=11.5 \mu M$ ) was reduced in the presence of 500 nM DOX ( $IC_{50}=22.0 \mu M$ ). After treatments of 18 h, DOX induced cell cycle arrest displaying a trimodal distribution; successive G2/M, S and G0/G1 blockage was produced with an  $IC_{50}$  of 49 nM, 464 nM and 1866 nM, respectively, whereas in the presence of 7.5  $\mu M$  MD, increasing levels of DOX mainly induced S-phase arrest. Within 18 hours of exposure, DOX induced apoptosis in a biphasic dose-dependent manner ( $K_d=335$  nM and 3.29  $\mu M$ , respectively). Addition of 7.5  $\mu M$  MD enhanced apoptosis at  $<300$  nM DOX, but reduced cell death at higher levels of DOX. However, 48 h after drug removal the apoptotic rate was considerably higher in cells exposed to DOX:MD, which also showed consistent fractions of early apoptosis (up to 44%). The efficacy of DOX was doubled by MD( $K_d=46.5$  nM in the presence, and  $K_d=99$  nM and 143 nM in the absence of MD).

**Conclusion:** Data indicate that clinically relevant levels of MD and DOX in combined treatments can exert considerable cytotoxic impact on Jurkat cells, via cell cycle arrest and apoptosis induction. These findings could encourage new therapeutic strategies to improve the therapeutic index of doxorubicin in ALL treatments.

**Acknowledgements:** This work was supported by a fellowship of the Romanian Ministry of Education, UEFISCDI, for Young Researchers, project number 8/2016.

#### References

1. Baran I, et al. Cell Biochem Biophys. 2010;58:169–79.
2. Baran I, et al. Leukemia Res. 2014;38:836–49.
3. Tofolean IT, et al. Pharmacol Res. 2016;103:300–17.

<http://dx.doi.org/10.1016/j.pbj.2017.07.087>

#### PS068

##### Effect of symptom interval and demographic characteristics on initial stage of malignant tumors in children



R. Grujicic\*, O. Djurmez, M. Trkulja, J. Lazić, M. Bjelić

School of Medicine, University of Belgrade, Serbia

E-mail address: [robertogrujicic@gmail.com](mailto:robertogrujicic@gmail.com)  
(R. Grujicic).

**Aim:** The aim of our retrospective study was to determine the influence of demographic and clinical characteristics of patients, initial stage of disease and tumor size on symptom period in children with malignant tumors.

**Introduction:** One of the main goals in pediatric oncology is timely diagnosis, cause it allows prompt and more effective treatment and significantly decreases the number of complications. The majority of children with malignant tumors have specific or non-specific symptoms certain time period before the diagnosis which can point towards malignant disease.

**Methods:** Our study included 296 children with malignant tumors, diagnosed and

treated between 2005 and 2016 in University Children's Hospital in Belgrade. Collected data included sociodemographic parameters, variety of symptoms and its duration, initial stage of disease and size of the tumor.

**Results:** The most frequent tumors were as follows: neuroblastoma, Hodgkin and non-Hodgkin lymphoma and kidney tumors. Non-Hodgkin lymphoma was diagnosed more frequently in boys, while Ewing sarcoma and primitive neuroectodermal tumors were seen mostly in girls. The majority was admitted at IV stage (30.1%) in opposite to 13.5% of patients in I stage. The average symptom interval was 87.7 days (median 46; SD= 164), from 5 to 2190 days. We have proven that following factors have significant effect on the extent of symptom interval: age ( $p < 0.001$ ), type of tumor ( $p < 0.05$ ), its localization ( $p < 0.001$ ), specific symptoms ( $p < 0.05$ ), and referral from primary health care unit in comparison to secondary one ( $p < 0.05$ ).

**Conclusion:** The results of our study give a new insight in symptom interval of children with malignant tumors in our country. More detailed comprehension of patients' characteristics, their diseases, healthcare system and their effect on symptom interval could significantly contribute to early diagnosis, as well as decreased number of complications at admission and during treatment.<sup>1–6</sup>

## References

- Atanaskovic Z, Kocев N, Penev G. The burden of disease and injury in Serbia. Beograd: Narodna biblioteka Srbije; 2003. p. 94–102.
- Little J. Epidemiology of Childhood Cancer. International agency for research on cancer; 1999. p. 342–50.
- Dang-tan T, Franco EL. Diagnosis delays in childhood cancer. *Cancer*. 2007;110:703–13.
- Dang-tan T, Trotter H, Mery LS, et al. Determinants of delays in treatment initiation in children and adolescents diagnosed with leukemia or lymphoma in Canada. *Int J Cancer*. 2010;126:1936–43.
- Wallach M, Balmer A, Munier F, Houghton S, Pampallona S. Shorter time to diagnosis and improved stage at presentation in Swiss patients with retinoblastoma treated from 1963 to 2004. *Pediatrics*. 2006;118:1493–8.
- Veneroni L, Mariani L, Vullo S, Lo, et al. Symptom interval in pediatric patients with solid tumors: adolescents are at greater risk of late diagnosis. *Pediatr Blood Cancer*. 2013;60:605–10.

<http://dx.doi.org/10.1016/j.pbj.2017.07.088>

## PS069

### Impact of prior malignancies on the outcome of colorectal cancer: Revisiting clinical trial eligibility criteria

Anas M. Saad<sup>1,\*</sup>, Muneer J. Al-Husseini<sup>1</sup>,  
Hadeer H. Mohamed<sup>1</sup>, Mohamad A. Alkhatat<sup>1</sup>,  
Mohamad Bassam Sonbol<sup>2</sup>,  
Omar Abdel-Rahman<sup>3</sup>

<sup>1</sup> Faculty of Medicine, Ain Shams University, Cairo, Egypt



<sup>2</sup> Mayo Clinic Cancer Center, Phoenix, Arizona, USA

<sup>3</sup> Faculty of Medicine, Ain Shams University, Clinical Oncology, Cairo, Egypt

E-mail address: [anassaad256@gmail.com](mailto:anassaad256@gmail.com)

(A.M. Saad).

**Aim:** To study the impact of prior malignancies on the survival of subsequent CRC.

**Introduction:** Colorectal cancer (CRC) is the third most common cancer in the US.<sup>1–3</sup> Some studies have correlated a prior history of malignancy with an increased incidence of CRC. Patients with history of cancer are generally excluded in clinical trials. This practice, not only affects clinical trials accrual, but also limits the potential therapeutic options for this population. The rationale behind this exclusion is that a history of malignancy could potentially interfere with the study outcomes.<sup>4</sup> However, little is known about its real impact on survival of subsequent CRC.

**Methods:** We identified patients with CRC diagnosed between 1973 and 2008 using the National Cancer Institute's SEER database.<sup>5,6</sup> Outcomes of interest were overall survival and cause-specific survival of subsequent CRC in general, and specifically stage IV disease. Unadjusted Kaplan-Meier test and multivariable covariate-adjusted Cox models were used to assess the eligibility of enrollment of stage IV CRC patients in clinical trials.

**Results:** Overall, 550,325 patients with CRC were identified, of whom 31,663 patients had a prior malignancy. Both, history of prior non-leukemic malignancy and prior leukemia were associated with a worse overall survival (HR = 1.165 95% CI = 1.148–1.183,  $P < 0.001$ ) and (HR = 1.825 95% CI = 1.691–1.970,  $P < 0.001$ ), respectively. However, a history of any prior non-leukemic malignancy showed a favorable colorectal-specific survival (HR = .930 95% CI = .909–.952,  $P < 0.001$ ). Analysis of stage IV CRC showed that a history of any prior non-leukemic malignancy was not associated with a significant difference in overall survival but having a history of leukemia showed a worse overall survival (HR = 1.535, 95% CI = 1.303–1.809,  $P < 0.001$ ).

**Conclusion:** Clinical trials should take these results into consideration when including/excluding stage IV CRC patients with prior malignancies.

## References

- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA: Cancer J Clin*. 2016;66:7–30.
- Siegel RL, Miller KD, Fedewa SA, Ahnen DJ, Meester RGS, Barzi A, et al. Colorectal cancer statistics, 2017. *CA: Cancer J Clin*. 2017;67:177–93.
- Siegel R, DeSantis C, Jemal A. Colorectal cancer statistics, 2014. *CA: Cancer J Clin*. 2014;64:104–17. <http://dx.doi.org/10.3322/caac.21220>.
- Laccetti AL, Pruitt SL, Xuan L, Halm EA, Gerber DE. Effect of prior cancer on outcomes in advanced lung cancer: implications for clinical trial eligibility and accrual. *J Natl Cancer Inst*. 2015;107.
- Surveillance Research Program, National Cancer Institute SEER\*Stat software ([www.seer.cancer.gov/seerstat](http://www.seer.cancer.gov/seerstat)) version 8.3.3.
- Surveillance, Epidemiology, and End Results (SEER) Program ([www.seer.cancer.gov](http://www.seer.cancer.gov)) SEER\*Stat Database: Incidence – SEER 18 Regs Research Data+Hurricane Katrina Impacted Louisiana Cases, Nov 2015 Sub (1973–2013 varying) – Linked To County Attributes – Total U.S., 1969–2014 Counties, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2016, based on the November 2015 submission.

<http://dx.doi.org/10.1016/j.pbj.2017.07.089>

## PS071

### Intervention of diabetes mellitus and metabolic risk factors in AMPK-PGC1 $\alpha$ -SIRT3 pathway in the human corpus cavernosum

A. Santos Pereira<sup>1,3,\*</sup>, A.R. Rodrigues<sup>1</sup>, B. Rocha<sup>1</sup>,  
N. Tomada<sup>2</sup>, A.M. Gouveia<sup>1,4</sup>, D. Neves<sup>1</sup>



<sup>1</sup> Department of Biomedicine - Experimental Biology Unit, Faculty of Medicine of the University of Porto, Al. Prof. Hernâni Monteiro, 4200-319 Porto, Portugal and Instituto de Investigação e Inovação em Saúde (I3S) Rua Alfredo Allen, 208, 4200-135 Porto, Portugal

<sup>2</sup> Department of Urology, Central Hospital of S. João-Alameda Prof. Hernâni Monteiro, 4200-319 Porto, Portugal

<sup>3</sup> Faculty of Medicine, University of Porto, Portugal

<sup>4</sup> Faculty of Nutrition and Food Sciences, University of Porto, Portugal

E-mail address: [andressasp16@gmail.com](mailto:andressasp16@gmail.com)

(A. Santos Pereira).

**Aim:** This study aims to investigate the signaling pathway AMP-activated protein kinase (AMPK)-Peroxisome proliferator-activated receptor-gamma coactivator (PGC)1 $\alpha$ -sirtuin (SIRT)3 in the human corpus cavernosum (HCC) between healthy individuals and those with cardiovascular disease risk factors (CVDRF).

**Introduction:** SIRT3 is a mitochondrial NAD<sup>+</sup>-dependent protein-deacetylase involved in the regulation of cellular metabolism.<sup>1,2</sup> As a key factor in AMPK and PGC1- $\alpha$  activation in stress, the decrease in SIRT3 expression or activity is associated with diverse pathologies and aging. Actually, SIRT3 expression was found decreased in HCC of aged individuals with CVDRF.<sup>3</sup> CVDRF such as diabetes mellitus (DM), dyslipidemia, hypertension and obesity strongly associate to endothelial dysfunction, which early manifests as erectile dysfunction (ED).<sup>4</sup>

**Methods:** HCC's samples from individuals aged 40-60 years, submitted to programmed urological surgeries at Hospital São João-Porto, were divided in three groups (n=4): (1)-controls without ED or CVDRF; (2)-DM patients; and (3)-patients with three or more CVDRF including DM. Dual immunolabelling of SIRT3 and superoxide dismutase (SOD)2 with alpha-actin was carried out. As well, levels of SIRT1, SIRT3, SOD2, PGC1 $\alpha$ , NADPH oxidase (Nox)1, phospho-AMPK and AMPK were assessed by Western-blotting(WB).

**Results:** We observed SIRT3 and SOD2 expression in  $\alpha$ -actin-labelled fusiform muscle cells in all groups. The semi-quantification by WB demonstrated a significant decrease in SOD2 expression in group 3 relatively to controls, as well as, an increased tendency of Nox1 and PGC1 $\alpha$  and a decreasing trend in phospho-AMPK in groups 2 and 3. No differences in SIRT1 and SIRT3 were observed among groups.

**Conclusion:** This study suggests that CVRF including DM increase oxidative stress in HCC owing to a decrease in SOD2 expression and concomitant increment in Nox1. Further studies with an increased number of HCC samples will be necessary to elucidate the role of the AMPK-PGC1 $\alpha$ -SIRT3 signaling pathway in the response to oxidative damage.<sup>5</sup>

**Acknowledgements:** Adriana R Rodrigues was supported by QREN-POPH, FSE and "Fundação para a Ciência e Tecnologia" (SFRH/BPD/92868/2013).

## References

1. Frye RA. Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity. *Biochem Biophys Res Commun.* 1999;260:273–9.
2. Schwer B, et al. The human silent information regulator (Sir)2 homologue hSIRT3 is a mitochondrial nicotinamide adenine dinucleotide-dependent deacetylase. *J Cell Biol.* 2002;158:647–57. PMC.
3. Freitas M, et al. Effects of aging and cardiovascular disease risk factors on the expression of sirtuins in the human corpus cavernosum. *J Sex Med.* 2015;12:2141–52.
4. Guay AT. ED2: erectile dysfunction = endothelial dysfunction. *Endocrinol Metab Clin North Am.* 2007;36:453–63.

5. Yu L, et al. Melatonin ameliorates myocardial ischemia/reperfusion injury in type 1 diabetic rats by preserving mitochondrial function: role of AMPK-PGC-1 $\alpha$ -SIRT3 signaling. *Sci Rep.* 2017;7.

<http://dx.doi.org/10.1016/j.pbj.2017.07.090>

## PS075

### Examination of antiproliferative effects of the horseradish extracts



L. Đurić<sup>1,\*</sup>, D. Četojević-Simin<sup>2</sup>, M. Milanović<sup>1</sup>

<sup>1</sup> University of Novi Sad, Faculty of Medicine, Department of Pharmacy, Novi Sad, Serbia

<sup>2</sup> University of Novi Sad, Faculty of Medicine, Experimental Oncology Department, Oncology Institute of Vojvodina, Sremska Kamenica, Serbia  
E-mail address: [djuriclarisa@gmail.com](mailto:djuriclarisa@gmail.com)

(L. Đurić).

**Aim:** The aim of the study was to investigate in vitro the antiproliferative effects of the horseradish juice and pulp using different solvents for the extraction.

**Introduction:** Horseradish (*Armoracia rusticana*, Brassicaceae) is a perennial herbal plant, which is widely used in human nutrition, as well as in a traditional medicine. Horseradish is a rich source of bioactive compounds such as isothiocyanates, that have proved to be significant antitumor agents.

**Methods:** Samples were prepared by the Kupchak extraction method, and the antiproliferative effects of the horseradish juice and pulp extracts were examined on the human tumor cell line MDA-MB-231 (ER-, human breast adenocarcinoma). Cell growth was determined by measuring the total protein by colorimetric sulforhodamine B assay. The obtained results (expressed as mean  $\pm$  SD) were analyzed by Tukey HSD test and the differences were considered statistically significant at  $p < 0.05$ .

**Results:** According to the IC<sub>50</sub> parameter (the concentration that inhibited the cell growth by 50%), as an important indicator of the antiproliferative effects, the most pronounced antitumor activity was observed for chloroform juice extract (IC<sub>50</sub> = 5.52  $\pm$  1.47  $\mu$ g/ml). In addition, highly potent was chloroform pulp extract (IC<sub>50</sub> = 19.44  $\pm$  3.82  $\mu$ g/ml), as well as the dichloromethane juice (IC<sub>50</sub> = 26.50  $\pm$  4.15  $\mu$ g/ml) and pulp (IC<sub>50</sub> = 26, 01  $\pm$  2.45  $\mu$ g/ml) extracts. On the other hand, significantly lower in vitro antitumor effect was noticed for the butanol pulp extract (IC<sub>50</sub> = 114.52  $\pm$  0.28  $\mu$ g/ml). IC<sub>50</sub> values for butanol juice extract, as well as water juice and pulp extracts were higher than 500  $\mu$ g/ml.

**Conclusion:** The obtained results suggest that *A. rusticana* is as a significant source of antitumor agents, especially liposoluble isothiocyanates and as such, it should be recommended for further use in a human nutrition and prevention of cancer.

<http://dx.doi.org/10.1016/j.pbj.2017.07.091>

## PS080

### Contribution of the determination of numeric value of adc map in early detection of prostate cancer



Dj Perovic

Faculty of Medicine, University Novi Sad, Serbia  
E-mail address: [djukaperovic@yahoo.com](mailto:djukaperovic@yahoo.com).

**Aim:** To define the range of ADC values for the absence of malignant disease, as well as to determine the threshold of ADC values for suspected prostate cancer.

**Introduction:** Prostate cancer is the second most diagnosed cancer, and the second most common cancer-cause of deaths in men worldwide. The apparent diffusion coefficient (ADC) derived from DWI has been shown to improve the detection of prostate cancer and is the primary imaging method for the differentiation between low to high grade cancers. ADC values show reduction with increasing Gleason's score.

**Methods:** Prospective study included 60 subjects. Male patients were divided into the groups with pathohistology verified benign and malignant lesions (aged, 46–81; average age, 67.7 years) with abnormal PSA values (>4 ng/ml), and into control group (aged, 44–81; average age 65.3) with normal PSA values (0–4 ng/ml). Prostates were first examined on MRI, determining the diffusion values on ADC map, by placing the region of interest (ROI), through the middle of lesions. Later, the TRUS-guided biopsies were performed. Three intersections of the prostate (apex, middle, and base) were observed, and at total of 12 places (4 places per section) the mentioned methods were indicated.

**Results:** Statistically significant difference ( $p < 0.05$ ) between the groups of patients with malignant and benign lesions in relation to the ADC values of the apex, base and middle of prostate. ADC values of malignant lesions at apex were in range 952–1030, at base  $859\text{--}977 \times 10^{-6} \text{ mm}^2/\text{s}$ , while in benign lesion values at apex were in range 1234–1336, and at base  $1096\text{--}1183 \times 10^{-6} \text{ mm}^2/\text{s}$ .

**Conclusion:** Determination of the numerical value of ADC map represents a significant additional diagnostic parameter for prostate cancer. All values in the range of 1179–1229 for base, 1063–1139 for middle, and  $1199\text{--}1379 \times 10^{-6} \text{ mm}^2/\text{s}$  for apex were considered normal. Values between the range of  $857\text{--}1030 \times 10^{-6} \text{ mm}^2/\text{s}$  have been suspected for possible presence of the prostate cancer.

<http://dx.doi.org/10.1016/j.pbj.2017.07.092>

## PS101

### Deoxycytidine kinase expression in AML blasts and its relationship to leukemia-free and overall survival

L. Ng<sup>1,\*</sup>, C. Chan<sup>2</sup>, T. Au<sup>2</sup>, C.K. Cheng<sup>3</sup>, K.F. Mo<sup>2</sup>, W. Li<sup>2</sup>, K. Lei<sup>2</sup>, T. Mok<sup>2</sup>, M. Ng<sup>3</sup>, R. Raghupathy<sup>2</sup>

<sup>1</sup> Faculty of Medicine, The Chinese University of Hong Kong (CUHK)

<sup>2</sup> Partner State Key Laboratory of Oncology in South China, Sir YK Pao Centre for Cancer, Department of Clinical Oncology, Hong Kong Cancer Institute and Prince of Wales Hospital, CUHK

<sup>3</sup> Blood Cancer Cytogenetics and Genomics Laboratory, Department of Anatomical and Cellular Pathology, Prince of Wales Hospital, CUHK, Hong Kong

E-mail address: [chitat.ng@gmail.com](mailto:chitat.ng@gmail.com)

(L. Ng).

**Aim:** Study the correlation of expression of genes involved in cytarabine metabolism to leukemia-free (LFS) and overall survival (OS) in AML.

**Introduction:** Cytarabine is the backbone of AML therapy. Understanding the roles and polymorphisms of genes involved in cytarabine metabolism and resistance in AML will facilitate development of novel therapeutics.

**Methods:** Adults less than 60 years with non M3 AML were included. Archived diagnostic marrow samples were studied for expression of 10 genes involved in cytarabine metabolism by RT-qPCR; gene expression normalized to GAPDH was compared

using the unpaired t-test with Welch correction. SNP rs4694362 in deoxycytidine kinase (DCK) gene was tested using Taqman assay. Median time to relapse and survival was calculated by Kaplan Meier method.

**Results:** 21 Han Chinese patients (median age: 50) were identified; 15 were male; 16 had intermediate risk cytogenetics; 5 had a blast count of over  $100 \times 10^9/\text{L}$  at diagnosis. 17 patients achieved CR; 12 after first induction. No difference in gene expression was seen between CR ( $n = 12$ ) versus non CR ( $n = 9$ ) with first induction. 17 CR patients were followed for a median duration of 67.5 months; median time to relapse was 15 months. 1 patient who underwent allogeneic transplant in CR1 was excluded. Higher mean DCK expression was seen in patients with LFS longer than ( $n = 7$ ) versus less than 2 years ( $n = 9$ ) ( $1.91 \pm 0.67$ ,  $p = 0.01$ ) and in those with OS longer than ( $n = 8$ ) versus less than 3 years ( $n = 8$ ) ( $2.02 \pm 0.69$ ,  $p = 0.01$ ). DCK rs4694362 TT genotype was less prevalent than CT in patients with >2 year LFS; but not statistically significant. (49% vs 60%,  $p = 0.6$ ).<sup>1,2</sup>

**Conclusion:** DCK phosphorylates cytarabine to its active metabolite. Our work shows that higher DCK expression is correlated to LFS and OS in AML. The role of DCK SNP rs4694362 should be explored further in the Chinese.

## References

1. Cai J, Damaraju VL, Groulx N, et al. Two distinct molecular mechanisms underlying cytarabine resistance in human leukemic cells. *Cancer Res.* 2008;68:2349–57.
2. Abraham A, Varatharajan S, Karathedath S, et al. RNA expression of genes involved in cytarabine metabolism and transport predicts cytarabine response in acute myeloid leukemia. *Pharmacogenomics.* 2015;16:877–90.

<http://dx.doi.org/10.1016/j.pbj.2017.07.093>

## PS108

### Analysis of MTHFR C677T polymorphism significance in patient preparation for the in vitro fertilization procedure



Amalija Stojanovic\*, Stevan Stojanovic, Suzana Sredic

*Institute for Human Genetics, Faculty of Medicine, University of Belgrade, Serbia*

E-mail address: [amalija.stojanovic@gmail.com](mailto:amalija.stojanovic@gmail.com)

(A. Stojanovic).

**Aim:** The aim of this study is to determine if in relation to general population, there is a statistically significant difference in the frequency of alleles and genotypes of the C677T polymorphism of MTHFR gene, amongst women with unknown cause of infertility, who are undergoing in vitro fertilization preparation.

**Introduction:** Methylene tetrahydrofolate reductase (MTHFR) is an enzyme coded by MTHFR gene. Polymorphism of MTHFR gene C677T leads to decreased function of MTHFR enzyme, which is associated with high level of homocysteine and low concentration of folate, which can undermine female reproductive function and affect the outcome of in vitro fertilization.

**Methods:** The study included the experimental group consisted of 31 women and the control group consisted of 100 women. C677T polymorphism was detected via PCR/RFLPS method. The statistical difference in genotype and allele frequencies was conducted using the Chi-square test.

**Results:** The comparison of genotypes amongst the experimental and control group has not shown a statistically significant difference ( $p > 0.05$ ). Frequency of the MTHFR677 TT genotype is 22.6% in the experimental group, and 12.0% in the control group, while the allele T frequency amongst the experimental group was

41.9%, and the frequency of the aforementioned allele amongst the control group was 34.5%.

**Conclusion:** The results of this study show that there is no statistically significant correlation between MTHFR C677T polymorphism in women with infertility of unknown cause, who are undergoing in vitro fertilization preparation, but also underline the need for further research.

**Acknowledgements:** Mentor: Professor Dr. Ivana Novaković.

<http://dx.doi.org/10.1016/j.pbj.2017.07.094>

#### PS114

### Assessing the oxidative modification of proteins in inflamed placenta combined with iron deficiency anemia in the pregnant through histochemical method with bromophenol blue based on Mikel Calvo



Y. Karliichuk\*, V. Ilika

Department of Pathological Anatomy and Morphology, USA

E-mail address: [j.m.karliichuk@gmail.com](mailto:j.m.karliichuk@gmail.com)

(Y. Karliichuk).

**Aim:** To set features of OMB in the cytoplasm of decidua cells in basal plate of the placenta at chorioamnionitis with iron deficiency anemia in pregnant women by means of histochemical methods combined with computer microspectrophotometry.

**Introduction:** Decidua cells are important cells to the placenta, playing a significant role both in the physiology of pregnancy and during inflammation. The processes of oxidative modification of proteins (OMB) in inflammation are associated with increased levels of oxygen free radicals, which alter the properties of these macromolecules while oxidating amino groups of proteins. Anemic condition is accompanied by intensification of free radical processes in the blood and tissues, and iron deficiency additionally significantly modifies these processes.

**Methods:** 125 studied placentas, to compare the studied placental physiology of pregnancy and monitoring iron deficiency anemia without inflammation.

A histochemical reaction of bromophenol blue for “acidic” and “basic” proteins by Mikel Calvo was set in histological sections 5 µm thick.

Delta Optical Evolution 100 and Olympus SP-550UZ were used to obtain a digital copy of the image. Ratio R/B, which is the ratio between the amino and carboxyl groups in proteins, was determined by “ImageJ”.

Unpaired Student’s test calculated arithmetic mean and its error.

**Results:** When assessing visual histochemical preparations decidua cells are clearly stained, that is suitable for quantitative research, cell boundaries are defined through clear cell membrane coloring and contrasting color around decidua cells fibrinoid. Nuclei and nucleoli were visualized fairly well. “Basic” proteins prevailed in nucleoplasm, while “sour” in the nucleolus.

The decidua cells’ cytoplasm specific color has been mostly granular in nature, and spectral characteristics and optical density of color varied greatly.

Factor R/B at physiological pregnancy ( $n = 20$ ) was  $-1.04 \pm 0.008$  and in iron deficiency anemia ( $N = 21$ )  $-1.06 \pm 0.009$   $P > 0.05$ . In acute chorioamnionitis ( $n = 23$ )  $-1.08 \pm 0.009$ , and combined with iron deficiency anemia ( $N = 21$ )  $-1.09 \pm 0.009$   $P > 0.05$ . Regarding chronic chorioamnionitis ( $n = 20$ ) ratio  $-1.24 \pm 0.011$ , and combined with iron deficiency anemia ( $N = 21$ )  $-1.64 \pm 0.016$   $P < 0.001$ .

**Conclusion:** Conclusion. The intensity of OMB increases only in chronic form of chorioamnionitis in the decidua cells cytoplasm,

and combined with iron deficiency anemia significant performance increase has been observed.

<http://dx.doi.org/10.1016/j.pbj.2017.07.095>

#### PS121

### Comparison of Ras/Raf/MAPK signaling pathway in primary tumour and lymph node metastases – A report on an experimental study of two colorectal cancer cell lines (SW480 and SW620) and tissue samples



K. Kałuzińska\*, A. Wach, P. Frączek

Chair of Medical Biochemistry, USA

E-mail address: [ka.kaluzinska@gmail.com](mailto:ka.kaluzinska@gmail.com)

(K. Kałuzińska).

**Aim:** To compare the presence of mutations in essential genes of CRC pathogenesis pathway between tissues derived from the primary tumour site and lymph node metastases.

**Introduction:** Colorectal cancer (CRC) remains the third most commonly diagnosed malignancy worldwide and a leading cause of cancer - related death. One of the pivotal pathways leading to CRC development is Ras/Raf/MAPK which is regulated by the receptor for the EGF. Mutations in these genes predict lack of response to EGFR-targeting monoclonal antibodies. However it is a common practice to assess only the primary tumour site, while mutations in metastasis may also affect the response to treatment.

**Methods:** The study was conducted on 10 patient-derived tissue samples and two ATCC human CRC cell lines obtained from the same individual: SW480 (primary tumour) and SW620 (lymph node metastasis). Cell lines were cultured according to the protocol. Genomic DNA and RNA were isolated, and PCR and RT-PCR were conducted. Primers for PCR included the following fragments: KRAS (exons 2,3,4), NRAS (exons 2,3,4), BRAF (exon 15); and for RT-PCR: KRAS, NRAS, BRAF and EGFR. Restriction enzymes were used. Proteins were extracted, purified and Western-Blot (RAS, RAF, MAPK) was performed.

**Results:** For SW480 we detected a mutation in exon 3 of NRAS gene, whereas SW620 presented a wild type. The level of Ras protein remained the same. Raf protein expression was abundant in the primary tumour site as compared to the lymph node metastasis, whereas MAPK protein presented the opposite level of expression.

**Conclusion:** The analysis of Ras-Raf-MAPK pathway may suggest that along with the tumour progression, the dominating signal is located at deeper levels of signaling pathway. Due to existing differences in key molecular points between the primary tumour and its metastases, in the era of targeted therapy, pre-treatment assessment of both sites has a potential to become a standard of care.<sup>1,2</sup>

#### References

1. Van Cutsem E, Cervantes A, et al. ESMO consensus guidelines for the management of patients with metastatic colorectal cancer. *Ann Oncol.* 2016;27:1386–422.
2. Negru S, Papadopoulou E, Apeessos A, et al. KRAS, NRAS and BRAF mutations in Greek and Romanian patients with colorectal cancer: a cohort study. *BMJ Open.* 2014;4:e004652. <http://dx.doi.org/10.1136/bmjopen-2013-004652>.

<http://dx.doi.org/10.1016/j.pbj.2017.07.096>

#### PS122

### PI3K-Akt and Ras-Raf-MAPK signaling in colorectal cancer – Comparison of activity in primary tumor tissues and primary tumour – Derived human colorectal cancer cell lines



A. Wach\*, K. Kałuzińska, P. Frączek

Chair of Medical Biochemistry, Estonia

E-mail address: [adamwach27@gmail.com](mailto:adamwach27@gmail.com)

(A. Wach).

**Aim:** The study aimed to compare the differences in activity of PI3K-Akt and Ras-Raf-MAPK pathways, and changes in the Ras-Raf-MAPK activity after PI3K-Akt silencing, between different cell lines and tissue samples from primary tumour sites of human CRC.

**Introduction:** Alterations in EGFR-related Ras-Raf-MAPK and PI3K-Akt pathways are involved in the pathogenesis of up to 55% and 15% colorectal cancers (CRC) respectively. The Ras-Raf-MAPK pathway mutations are assessed before introducing a standard anti-EGFR treatment, as they indicate lack of response. However, the autonomic activity of alternative PI3K-Akt pathway may also have an impact on the effectiveness of targeted therapy.

**Methods:** The study was carried out on three ATCC human CRC cell lines derived from primary tumours (COLO320, SW480 and HT29) and ten patient tissue samples. Cell lines were cultured according to the protocol. Genomic DNA and RNA were isolated, PCR and RT-PCR were performed. Restriction enzymes were applied. Primers for the following fragments of genome were used: KRAS (exons 2, 3, 4), NRAS (exons 2, 3, 4), and BRAF exon 15 for PCR; KRAS, NRAS, BRAF, PIK3CA for RT-PCR. Proteins were extracted, purified and Western Blot was conducted. siRNA for Akt and specific PI3K inhibitors were used to silence PI3K-Akt activity.

**Results:** The analyzed material presented variable profiles of pathways activity. Interestingly, high expression of Ras protein was positively correlated with Akt protein level. In case of low level of Ras, Raf protein was dominating whereas Akt expression was significantly decreased.

**Conclusion:** Ras and Akt can simultaneously present a high level of expression. Thus, as PI3K-Akt is an alternative pathway to Ras-Raf-MAPK for EGFR signaling and its autonomic activity may affect the efficacy of anticancer treatment, it has a potential to be taken into consideration while planning a treatment and developing new anticancer agents.<sup>1,2</sup>

## References

1. Muzny DM, Bainbridge MN, et al. Comprehensive molecular characterization of human colon and rectal cancer. *Nature*. 2012;487:330–7.
2. Khwanraj K, Madlah S, et al. Comparative mRNA expression of eEF1A isoforms and a PI3K/Akt/mTOR pathway in a cellular model of parkinson's disease. Hindawi Publishing Corporation; 2016. p. 11. Article ID 8716016.

<http://dx.doi.org/10.1016/j.pbj.2017.07.097>

## PS124

### The role of the hypoxic tumor microenvironment on the macrophage-tumor cell interplay

F. Martins<sup>1,2,3,\*</sup>, F. Castro<sup>2,3,4</sup>, M.L. Pinto<sup>2,3,4</sup>, A.J. Silva<sup>2,3</sup>, B. Sousa<sup>2,5</sup>, M.J. Oliveira<sup>2,3,6</sup>, Â.M. Costa<sup>2,3</sup>

<sup>1</sup> Department of Biology, Faculty of Sciences, UPorto, Porto, Portugal

<sup>2</sup> i3S - Instituto de Investigação e Inovação em Saúde, UPorto, Porto, Portugal

<sup>3</sup> INEB - Institute of Biomedical Engineering, UPorto, Portugal

<sup>4</sup> ICBAS- Instituto de Ciências Biomédicas Abel Salazar, UPorto, Porto, Portugal

<sup>5</sup> IPATIMUP- Institute of Molecular Pathology and Immunology of the University of Porto, Portugal



<sup>6</sup> Department of Pathology and Oncology, Faculty of Medicine, UPorto, Porto, Portugal  
E-mail address: [flavia.martins@i3s.up.pt](mailto:flavia.martins@i3s.up.pt)  
(F. Martins).

**Aim:** The aim of this work is to unveil the role of the hypoxic microenvironment on macrophage-tumor cell interplay, using colorectal cancer (CRC) as a model.

**Introduction:** Microenvironment, in most cases hypoxic, is composed by cancer cells, extracellular matrix, stromal and immune cells, that cooperate and affect each other activities. Macrophages are one of the most abundant immune cells at the tumor microenvironment, acting as tumor suppressors or promoters. Previous research had shown that both hypoxia and immunosuppressive macrophages are associated with tumor progression. Nevertheless, these studies did not focus on the interplay between hypoxia and macrophage-cancer cell crosstalk.

**Methods:** To achieve our goal co-cultures of CRC cells and human macrophages, both in normoxia and hypoxia, were established. Macrophages were characterized functionally and phenotypically and their potential to induce cancer cell invasion was evaluated.

**Results:** Our results suggest that hypoxia, and the presence of cancer cells, decreases the cell surface expression of an anti-inflammatory marker (CD163), however the mRNA expression was not altered. Nevertheless, hypoxia induced an increase in the mRNA expression of the macrophage pro-inflammatory marker (CCR7).

Macrophages metabolic activity was not altered by hypoxia but decreased when co-cultured with cancer cells. In addition, lactate production decrease in co-culture while glucose consumption increased. Notably, macrophages in normoxia presented a more rounded morphology while in hypoxia are more elongated with evident cellular protrusions, suggesting dynamic alterations at the actin cytoskeleton organization. Interestingly, MMP-2 and MMP-9 activity profiles were not altered by the presence of cancer cells or hypoxia. Nevertheless, cancer cell invasion ability increased in the presence of macrophages, suggesting that other MMPs might be involved.

**Conclusion:** Findings in normoxia regarding macrophage potential to induce cancer cell invasion are consistent with those previously described by our group. Interestingly, we demonstrate now that hypoxia potentiates the invasive behavior of cancer cells and also macrophage pro-invasive ability.

<http://dx.doi.org/10.1016/j.pbj.2017.07.098>

## PS129

### Ethnopharmacological use of *Cymbopogon citratus* (DC.) Stapf and *Cymbopogon schoenanthus* (L.) Spreng.: Anti-inflammatory potential of phenol-rich extracts



Elisabete Gomes\*, João Bernardo, Mariana Barbosa, Paula B. Andrade, Patrícia Valentão, Graciliana Lopes

REQUIMTE/LAQV, Laboratório de Farmacognosia, Faculdade de Farmácia, Universidade do Porto, Rua Jorge Viterbo Ferreira, n° 228, Porto 4050-313, Portugal

E-mail address: [elisabetepatriciagomes@gmail.com](mailto:elisabetepatriciagomes@gmail.com)  
(E. Gomes).

**Aim:** The aim of this work consisted on expanding the knowledge on the chemical composition of different extracts from *Cymbopogon* spp., and on the evaluation of their anti-inflammatory potential in cell and cell-free systems.

**Introduction:** The ethnopharmacological use of *Cymbopogon* spp. dates back from ancient times. Traditionally used in tropical and semi-tropical countries for the repellent properties of their essential oil, the consumption of *Cymbopogon* spp. infusions is growing all over the world. This is not only due to the unique aroma, widely appreciated by the consumers, but also because of the antimicrobial, anti-inflammatory and sedative properties.<sup>1</sup>

**Methods:** The chemical characterization of infusions and ethanol:water (50:50, v/v) extracts from *Cymbopogon citratus* and *Cymbopogon schoenanthus* was achieved by HPLC-DAD. The anti-inflammatory potential of the extracts was assessed by cell and cell-free assays.

**Results:** HPLC-DAD analysis allowed the identification of several caffeic acid derivatives and flavonoids in the infusions and in the ethanol:water extracts of both species. The different extracts displayed scavenging activity against superoxide anion and nitric oxide (NO) radicals, and capacity to significantly reduce NO production by LPS-stimulated macrophages (RAW 264.7 cell line). In addition, the extracts were able to prevent hyaluronic acid degradation via inhibition of hyaluronidase, an enzyme recognized to participate in a number of physiological and pathological processes, including inflammation.<sup>2</sup> No toxicity was observed on human gastric adenocarcinoma and hepatocyte carcinoma cell lines, at a maximum concentration of 2.0 mg lyophilised extract/mL.

**Conclusion:** This study provided scientific evidence on the ethnopharmacological use of *Cymbopogon* species on inflammatory conditions, encouraging infusion consumption and future incorporation of *Cymbopogon* spp. extracts into nutraceuticals.

**Acknowledgements:** This work received financial support from National Funds (FCT/MEC) through project UID/QUI/50006/2013, co-financed by FEDER through COMPETE, under the Partnership Agreement PT2020, and from NORTE 2020, under the PORTUGAL 2020 Partnership Agreement, through ERDF (NORTE-01-0145-FEDER-000024). MB is indebted to FCT for the grant (SFRH/BD/95861/2013).

## References

- [1]. Negrelle RRB, Gomes EC. *Rev Bras Pl Med.* 2007;9:80–92.  
[2]. Ferreres F, et al. *Mar Drugs.* 2012;10:2766–81.

<http://dx.doi.org/10.1016/j.pbj.2017.07.099>

## PS140

### Cytotoxic effects of novel synthesized polyoxometalates on human neuroblastoma SH-SY5Y cell line



J. Isma\*, S. Jakovljević, A. Isaković

*Institute of Medical and Clinical Biochemistry,  
Faculty of medicine, University of Belgrade, Serbia  
E-mail address: isma.jovan@gmail.com  
(J. Isma).*

**Aim:** Investigation of cytotoxic effects of newly synthesized and untested polyoxometalates Pd1 and Pd2 on human neuroblastoma cells SH-SY5Y.

**Introduction:** Polyoxometalates (POMs) are transitional metal complexes, which are important in medicinal chemistry, as potent anticancer, antiviral and antibacterial agents. Inefficiently selective drugs and problems with dosing of usual chemotherapeutics directed the research towards investigation of new agents, such as POM.

**Methods:** Effects on viability rate of treated cells was tested using acid phosphatase assay. The mechanism of a cell death was examined using flow cytometry. JC-1, dihydroethidium, ApoStat,

propidium iodide and acridin orange stainings were conducted in order to elucidate mitochondrial depolarisation, production of superoxide anion, caspase activation, DNA fragmentation and intracellular acidity.

**Results:** Pd1 and Pd2 have shown dose and time dependent decrease in cell viability rate. Complexes induced mitochondrial depolarisation after 2 h of treatment, which was shown as increase in FL1/FL2 ratio from 1 to 1.3 (Pd1, 6  $\mu$ M) and from 1 to 1.7 (Pd2, 40  $\mu$ M). Superoxide anion production was increased after 5 h of treatment using Pd1 and 2 h of treatment using Pd2. Pd1 complex exhibits increase in percentage of cells with fragmented DNA (subG0) and activated caspases after 24 h treatment. Pd2 complex induced increase in SubG0 and S phase without caspase activation after 24 h treatment. POMs have shown intracellular acidification after 48 h (FL3/FL1 ratio: control 1, Pd1 2.3, Pd2 1.8).

**Conclusion:** POM complexes indicated cytotoxic effects on examined cell line. The mechanism by which these complexes exert those effects differ from one another. It was shown that both induce oxidative stress and mitochondrial depolarisation, accompanied by activation of caspases and DNA fragmentation in Pd1-treated cells, all indicative of apoptosis. In Pd2-treated group there was no increase in activation of caspases. Complexes have shown increase in intracellular acidification, which may suggest autophagy.

**Acknowledgements:** This research was a part of bilateral project between University of Belgrade, Belgrade, Serbia and Jacobs University, Bremen, Germany; 451-03-01038/2015-09/16.

<http://dx.doi.org/10.1016/j.pbj.2017.07.100>

## PS153

### HuR prevents c-fos mRNA degradation by proteasome-associated ribonuclease in vitro



E. Zhigalova<sup>1,\*</sup>, A. Mittenberg<sup>2</sup>

<sup>1</sup> *Skolkovo Institute of Science and Technology,  
Moscow, Russian Federation*

<sup>2</sup> *Institute of Cytology Russian Academy of Science,  
St. Petersburg, Russian Federation*

*E-mail address: katty\_zh@inbox.ru  
(E. Zhigalova).*

**Aim:** To estimate HuR protective activity against proteasome-associated ribonuclease for c-myc and c-fos mRNAs.

**Introduction:** Proteasome-associated proteins are attractive targets for multiple myeloma treatment. One of them is HuR protein known to selectively bind ARE-containing mRNAs and protect them from degradation. HuR is supposed to play a role in cancerogenesis since its expression is elevated in many cancer types and it stabilizes a lot of mRNAs encoding proteins involved in oncogenesis. Previously, it was shown that proteasome in addition to its main function – protein degradation – may act as a selective RNase. Moreover, HuR and proteasome have common targets – c-myc and c-fos protooncogene mRNAs.

**Methods:** HuR-GST fusion protein has been cloned, expressed and purified by affinity chromatography. Fragments of c-myc and c-fos were cloned and mRNAs have been transcribed in vitro. Proteasomes have been isolated from K562 cell line (human proerythrocytopenia) and Im-9 cells (human multiple myeloma). mRNAs were treated by proteasomes in presence and absence of HuR. The estimation of mRNA cleavage was held by gel-electrophoresis.

**Results:** GST-HuR has specifically bound ARE-containing fragments of c-myc and c-fos mRNAs. Proteasomes extracted from Im-9 and K562 cells cleaved target mRNAs in absence of HuR. It was shown that HuR prevents degradation of c-fos mRNA by proteasomal endoribonuclease, whereas c-myc mRNA was cleaved in the

same conditions. GST protein didn't bind with target mRNAs and didn't affect proteasome cleavage activity.

**Conclusion:** HuR protects c-fos mRNA from proteasome ribonuclease cleavage in vitro, but can't prevent c-myc mRNA degradation. HuR and proteasome compete with each other for manifestation of their opposite activities. Thus, a new mechanism of regulation of proto-oncogenes expression was observed. However, the functional role of this process in vivo should be evaluated in further studies.<sup>1–4</sup>

## References

1. Myer VE, Fan XC, Steitz JA. Identification of HuR as a protein implicated in AUUUA-mediated mRNA decay. *EMBO J.* 1997;16:2130–9.
2. Srikantan S, Gorospe M. HuR function in disease. *Front Biosci.* 2012;17:189–205.
3. Pouch MN, Petit F, Buri J, Briand Y, Schmid HP. Identification and initial characterization of a specific proteasome (prosome) associated RNase activity. *J Biol Chem.* 1995;270:22023–8.
4. Savant-Bhonsale S, Cleveland DW. Evidence for instability of mRNAs containing AUUUA motifs mediated through translation-dependent assembly of a > 20S degradation complex. *Genes Dev.* 1992;6:1927–39.

<http://dx.doi.org/10.1016/j.pbj.2017.07.101>

## PS156

### Metformin interferes with glucose cellular uptake by both estrogen and progesterone receptor-positive (MCF-7) and triple-negative (MDA-MB-231) breast cancer cell lines



I. Amaral\*, C. Silva, A. Correia-Branco, F. Martel

*i3S – FMUP, Portugal*

E-mail address: [inesamaral@ua.pt](mailto:inesamaral@ua.pt)

(I. Amaral).

**Aim:** Transport experiments with 3H-DG, culture growth and proliferation rate assays were performed. This work aimed to investigate the possible interference of metformin with glucose uptake by MCF-7 and MDA-MB-231 human breast adenocarcinoma cell lines as a mechanism contributing to its anticarcinogenic effect.

**Introduction:** Breast cancer, the most common cancer among women, remains one of the leading causes of mortality among women worldwide.<sup>1</sup> Metformin has been widely used as a treatment for type 2 diabetes for over 40 years.<sup>2</sup> The first report of a reduced risk of developing cancer for diabetic patients treated with metformin was published in 2005.<sup>3</sup> Several mechanisms of action of metformin appear to be implicated in this effect.<sup>2,4</sup>

**Methods:** Transport experiments with 3H-DG, culture growth and proliferation rate assays were performed.

**Results:** Acute (26 min) exposure of MCF-7 cells to metformin significantly inhibited uptake of 3H-deoxy-D-glucose (3H-DG) (maximal inhibition found with metformin 0.5 mM: 27 ± 2% reduction). Chronically (24 h), metformin induced a concentration-dependent increase in 3H-DG uptake (maximal increase observed with metformin 1 mM: 81 ± 15% increase). Acute (26 min) exposure of MDA-MB-231 cells to metformin slightly inhibited uptake of 3H-DG (maximal inhibition found with metformin 1 mM: 10 ± 3% reduction). Chronic (24 h) exposure to metformin significantly increased 3H-DG uptake by MDA-MB-231 cells (maximal increase observed with metformin 1 mM: 30 ± 8% increase).

Chronic (24 h) exposure of both cell lines to metformin (1 mM) decreased culture growth/cell mass; in contrast, it increased cell proliferation rates. Combination of metformin (1 mM) with the facilitative glucose transporter (GLUT) inhibitor kaempferol (30 μM) did not result in a more marked effect on culture growth and cell proliferation rates.

**Conclusion:** Summarizing, chronic exposure of MCF-7 and MDA-MB-231 cells to metformin induces a marked increase in glucose uptake, associated with an anticarcinogenic effect of the drug. We suggest that the increase in glucose uptake is a compensatory mechanism to cellular energy depletion induced by metformin.

**Acknowledgements:** This study was supported by Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Portugal (Plano estratégico UID/BIM/04293/2013).

## References

1. DeSantis, et al. *CA Cancer J Clin.* 2014;64:52–62.
2. Jara L-M. *Pharmacol Res.* 2015;101:102–8.
3. Evans, et al. *BMJ.* 2005;330:1304–5.
4. Daugan, et al. *Pharmacol Res.* 2016;113:675–85.

<http://dx.doi.org/10.1016/j.pbj.2017.07.102>

## PS162

### Endocannabinoids induce placental trophoblast reticulum stress



S.C.F. Pereira<sup>3,\*</sup>, M. Almada<sup>1,2</sup>, B.M. Fonseca<sup>1,2</sup>, L. Midão<sup>1,4</sup>, J. Maia<sup>1,2</sup>, N.A. Teixeira<sup>1,2</sup>, G. Correia-da-Silva<sup>1,2</sup>

<sup>1</sup> *Laboratório de Bioquímica, Faculdade de Farmácia Universidade do Porto, Porto, Portugal*

<sup>2</sup> *UCIBIO-REQUIMTE, Porto, Portugal*

<sup>3</sup> *Faculdade de Ciências e Instituto de Ciências*

*Biomédicas da Universidade do Porto, Porto, Portugal*

<sup>4</sup> *Departamento de Química, Universidade de Aveiro, Aveiro, Portugal*

E-mail address: [saracatarinapereira@gmail.com](mailto:saracatarinapereira@gmail.com)

(S.C.F. Pereira).

**Aim:** We aim to investigate in cytotrophoblasts whether these effects on cell viability loss are due to endoplasmic reticulum (ER) stress mediated apoptosis.

**Introduction:** Placental development relies on a balance between proliferation, differentiation and apoptosis of trophoblasts, a process tightly regulated by growth factors, cytokines and hormones. Endocannabinoids (eCB), such as 2-arachidonoylglycerol (2-AG) and anandamide (AEA) may play a role in these processes. We previously demonstrated that both eCB induced trophoblast cell death.<sup>1,2</sup> Here we investigated in cytotrophoblasts whether these effects on cell viability loss are due to endoplasmic reticulum (ER) stress mediated apoptosis.

ER stress is caused by the accumulation of unfolded proteins leading to an unfold protein response (UPR) triggered by transmembrane ER signaling proteins including: pancreatic ER kinase (PKR)-like ER kinase (PERK), inositol-requiring enzyme 1 (Ire1) and Activating transcription factor 6 (ATF6). The dissociation of Grp78 (BiP) from these sensors triggers a series of mechanisms that can restore homeostasis or lead to apoptosis. Placental stress has been implicated in the pathophysiology of pregnancy complications, including growth restriction and pre-eclampsia.

**Methods:** BeWo cells (ATCC, USA), an accepted model of cytotrophoblast stem cells were treated with AEA or 2-AG (10 micromolar) for 24 h. Through quantitative real time polymerase chain reaction (qPCR), we evaluated mRNA levels of ER stress markers: CHOP, Grp78, ATF4 and spliced mXBP1. Protein expression of CHOP was evaluated by western-blot.

**Results:** After 24 h of treatment with both eCB, we found an increase in mRNA levels of ER stress markers: CHOP, Grp78, ATF4 and spliced mXBP1. Protein expression of CHOP also increased in both cases.

**Conclusion:** These results suggest that cell viability loss promoted by 2-AG and AEA was associated with ER-stress since both PERK and IRE1 arms of UPR are activated. Prolonged ER-stress, contributes to the expression of pro-apoptotic proteins, such as CHOP.

These findings shed light to the impact of endocannabinoids induced-ER stress which may negatively affect trophoblast cell turnover and pregnancy outcomes.

**Acknowledgements:** This work received support from European Union (FEDER funds through COMPETE) and FCT through project PTDC/DTP-FTO/5651/2014-POCI-01-0145-FEDER-016562; FCT/MEC through national funds and co-financed by FEDER, under PT2020 (UID/01/0145/FERDER/007728) and CCDR-N/NORTE2020/Portugal 2020 (norte-01-0145-FEDER-000024).

## References

- Costa MA, Fonseca BM, Keating E, Teixeira NA, Correia-da-Silva G. 2-Arachidonoylglycerol effects in cytotrophoblasts: metabolic enzymes expression and apoptosis in BeWo cells. *Reproduction*. 2014;147:301–11.
- Costa MA, Fonseca BM, Teixeira NA, Correia-da-Silva G. The endocannabinoid anandamide induces apoptosis in cytotrophoblast cells: involvement of both mitochondrial and death receptor pathways. *Placenta*. 2015;36:69–76.

<http://dx.doi.org/10.1016/j.pbj.2017.07.103>

## PS163

### Analysis of imaging characteristics, incidence, and prognosis of brain metastases from thyroid cancer



Mafalda Sampaio Alves<sup>1,\*</sup>, Eduarda Carneiro<sup>4</sup>, Diana Ferreira<sup>4</sup>, Isabel Torres<sup>5</sup>, Susana Maria Silva<sup>2,3</sup>, Mavilde Arantes<sup>2,3,4</sup>

<sup>1</sup> Faculty of Medicine of the University of Porto, 4200-319 Porto

<sup>2</sup> Unit of Anatomy, Department of Biomedicine, Faculty of Medicine of the University of Porto, 4200-319 Porto

<sup>3</sup> Center for Health Technology and Services Research (CINTESIS), 4200-450 Porto, Portugal

<sup>4</sup> Division of Neuroradiology, Radiology Service, Portuguese Institute of Oncology, Porto, Portugal

<sup>5</sup> Endocrinology Service, Portuguese Institute of Oncology, Porto, Portugal

E-mail address: [sampaiolavesm@gmail.com](mailto:sampaiolavesm@gmail.com) (M.S. Alves).

**Aim:** The main objectives of this study were to evaluate the incidence, imaging characteristics, and prognosis of parenchymal brain metastases originating in thyroid cancer.

**Introduction:** While thyroid cancer is a relatively common type of cancer, it is usually highly curable.<sup>1</sup> Brain metastases from thyroid cancer are rare and their imaging appearance has not been well defined.<sup>2</sup>

**Methods:** Review of case records of thyroid cancer patients within the IPO Porto data base from 2005 to 2015 was conducted in order to identify the patients with thyroid cancer and evidence of brain metastases.

**Results:** We identified 3175 patients with thyroid cancer, with only five having evidence of brain metastases (two from papillary thyroid cancer, two from follicular thyroid cancer and one from poorly differentiated thyroid cancer). At the time of brain metastases detection, 100% of the patients had concurrent lymph node metastases, 80% lung metastases and 60% osseous metastases. Of those brain metastases, 60% were multifocal and 40% presented as partially cystic/necrotic. Of the two cases in which the patients

died, the median overall survival after brain metastasis detection was less than one year.

**Conclusion:** Brain metastasis from thyroid cancer remains a rare phenomenon that most frequently occurs in the setting of widely disseminated lymph node disease. The imaging appearance is highly variable and the prognosis is poor.

## References

- Hjiyiannakis P, Jefferies S, Harmer CL. Brain metastases in patients with differentiated thyroid carcinoma. *Clin Oncol*. 1996;8:327–30.
- Tsuda K, Tsurushima H, Takano S, Tsuboi K, Matsumura A. Brain metastasis from papillary thyroid carcinomas. *Molec Clin Oncol*. 2013;1:817–9.

<http://dx.doi.org/10.1016/j.pbj.2017.07.104>

## PS166

### The association of GSTP1 genotype with the risk and survival in ccRCC patients with advanced tumor stage



S. Mihailovic<sup>1,\*</sup>, T. Radic<sup>1,2</sup>, M. Pljesa Ercegovac<sup>1,2</sup>, V. Coric<sup>1,2</sup>

<sup>1</sup> Faculty of Medicine, University of Belgrade, Serbia

<sup>2</sup> Institute of Medical and Clinical Biochemistry, Faculty of Medicine, University of Belgrade, Serbia

E-mail address: [smiljanamihailovic@gmail.com](mailto:smiljanamihailovic@gmail.com) (S. Mihailovic).

**Aim:** The aim of this study was to evaluate specific role of glutathione S-transferase P1 (GSTP1) gene variants as determinants of ccRCC risk in patients with advanced tumor stage (pT3 and pT4). Furthermore, we evaluated the effect of GSTP1 gene variants on postoperative prognosis in these patients.

**Introduction:** Renal cell carcinoma (RCC) accounts for up to 90% of malignant kidney tumors with clear renal cell carcinoma (ccRCC) being the most frequent and the most aggressive subtype of sporadic RCC in adults. Unfortunately, most RCCs are asymptomatic in early stages, whereas symptomatic RCC correlates with aggressive histology and advanced disease. Aside from known risk factors for RCC, evidence suggest that the development of RCC can be partially explained by genetic variations among the populations. Highly polymorphic cytosolic glutathione S-transferases are known to be involved in both the development and the progression of renal cell carcinoma.

**Methods:** GSTP1 genotype was determined in 99 ccRCC patients and 326 matched-controls by qPCR method, using TaqMan<sup>®</sup> SNP Genotyping Assay. The risk for disease was computed by odds ratios (OR) and 95% confidence intervals (CI) using logistic regression analysis. Furthermore, overall survival was analyzed as well by Kaplan-Meier method and Cox proportional hazard regression model.

**Results:** GSTP1-variant genotype was associated with 5-fold increased risk for ccRCC in comparison with GSTP1-wild type genotype ( $p < 0.001$ ). Moreover, survival analysis clearly indicated shorter overall survival in ccRCC patients with GSTP1-variant genotype, however without reaching statistical significance ( $p = 0.166$ ). Additionally, ccRCC patients with GSTP1-variant genotype had a 7-fold higher hazard ratio ( $p = 0.177$ ), compared to the carriers of GSTP1-wild type genotype.

**Conclusion:** GSTP1-variant genotype contributed independently towards the risk of ccRCC in our patients. Moreover, GSTP1-variant genotype is associated with poor postoperative prognosis in ccRCC.

<http://dx.doi.org/10.1016/j.pbj.2017.07.105>

## PS171

**Neuroimaging analysis of rare brain metastases from prostate cancer**

Juliana Macedo<sup>1,\*</sup>, Eduarda Carneiro<sup>4</sup>,  
Diana Ferreira<sup>4</sup>, António Verdelho<sup>5</sup>,  
Luís Pedro Afonso<sup>6</sup>, Joaquina Maurício<sup>7</sup>,  
Susana Maria Silva<sup>2,3</sup>, Mavilde Arantes<sup>2,3,4</sup>

<sup>1</sup> Faculty of Medicine of the University of Porto,  
4200-319 Porto, Portugal

<sup>2</sup> Unit of Anatomy, Department of Biomedicine,  
Faculty of Medicine of the University of Porto,  
4200-319 Porto, Portugal

<sup>3</sup> Center for Health Technology and Services Research  
(CINTESIS), 4200-450, Portugal

<sup>4</sup> Division of Neuroradiology, Radiology Service,  
Portuguese Institute of Oncology, Porto, Portugal

<sup>5</sup> Neurosurgery Service, Portuguese Institute of  
Oncology, Porto, Portugal

<sup>6</sup> Pathological Anatomy Service, Portuguese Institute  
of Oncology, Porto, Portugal

<sup>7</sup> Medical Oncology Service, Portuguese Institute of  
Oncology, Porto, Portugal

E-mail address: [ju.p.macedo18@gmail.com](mailto:ju.p.macedo18@gmail.com)

(J. Macedo).

**Aim:** Our main study focus was to evaluate the incidence, imaging characteristics, and prognosis of parenchymal brain metastases originating from prostatic tumors.

**Introduction:** Prostate cancer is considered the second most commonly diagnosed cancer.<sup>1</sup> In addition, it is considered the fifth leading cause of cancer death amongst males.<sup>2</sup> A small percentage (2%) of patients with prostate cancer are found to be castrate-resistant and to develop brain metastasis, a rare complication which is associated to an advanced systemic state of the disease when the tumor has already spread to other sites.<sup>3</sup> Although, there is not much evidence on optimal management of these patients.<sup>4</sup>

**Methods:** A review of case records of prostate cancer patients within the IPO Porto data base from 2013 to 2015 was conducted in order to identify the patients with prostate cancer and evidence of brain metastases. As criteria of exclusion, cases transferred to other hospitals without follow up and cases that were incorrectly categorized were excluded.

**Results:** We screened 2194 patients with prostate cancer, with only one having evidence of brain metastasis. Additionally, one case was identified with bilateral orbital metastatic lesions. The patient with evidence of brain metastasis aged 48 years old. Magnetic resonance imaging showed six metastatic lesions, three infratentorial and three supratentorial. The largest lesion was found in the parieto-occipital region. These brain metastases were detected 42 months after the initial diagnosis of prostate adenocarcinoma. In addition, by the time of brain metastasis detection, the patient already had bone metastatic lesion.

**Conclusion:** Brain metastases from prostate cancer are rare, with only a few cases described in the literature. Variable magnetic resonance imaging characteristics are described. Brain metastases are also associated with a poor prognosis, with a mean survival of 1–7.6 months.

**References**

1. Barakat T, Agarwal A, McDonald R, Ganesh V, Vuong S, Borean M, et al. Solitary brain metastasis from prostate cancer: a case report. *Ann Palliat Med.* 2016;5:227–32.
2. Craig J, Woulfe J, Sinclair J, Malone S. Isolated brain metastases as first site of recurrence in prostate cancer: case report and review of the literature. *Curr Oncol.* 2015;22:e493–7.

3. Taddei G, Marzi S, Coletti G, De Paulis D, Ricci A, Galzio RJ. Brain metastasis from prostate adenocarcinoma: case report and review of literature. *World J Oncol.* 2012;3:83–6.
4. Gizewska A, Witkowska-Patena E, Stembrowicz-Nowakoska Z, Buraczewska A, Dziuk M. Brain metastases in patient with prostate cancer found in 18F-choline PET/CT. *Nuclear Med Rev.* 2015;18:39–41.

<http://dx.doi.org/10.1016/j.pbj.2017.07.106>

## PS177

**Studies towards the synthesis of dicarboxylic acid metabolite of mitoxantrone**

Ivanna Hrynchak<sup>1</sup>, Emília Sousa<sup>1,2,\*</sup>,  
Maria de Lourdes Bastos<sup>1,2,3</sup>, Madalena Pinto<sup>1,2</sup>,  
Vera Marisa Costa<sup>3</sup>

<sup>1</sup> Laboratório de Química Orgânica e Farmacêutica,  
Departamento de Ciências Químicas, Faculdade de  
Farmácia, Universidade do Porto, Portugal

<sup>2</sup> Centro Interdisciplinar de Investigação Marinha e

Ambiental (CIIMAR), Universidade do Porto, Portugal

<sup>3</sup> UCIBIO, REQUIMTE (Rede de Química e Tecnologia),

Laboratório de Toxicologia, Departamento de

Ciências Biológicas, Faculdade de Farmácia,

Universidade do Porto

E-mail address: [esousa@ff.up.pt](mailto:esousa@ff.up.pt)

(E. Sousa).

**Aim:** The objective of this work was to synthesize the dicarboxylic acid metabolite of mitoxantrone (MTX) to further investigate its cardiotoxicity in H9c2 differentiated cells.

**Introduction:** Drug metabolism can result in active or toxic metabolites that can lead to side effects, namely cardiotoxicity.<sup>1</sup> MTX, an antineoplastic that belongs to the synthetic anthracenediones,<sup>2</sup> is used to treat breast cancer, acute leukaemias, and acute lymphomas in adults.<sup>3</sup> Nowadays, it is also used to treat aggressive multiple sclerosis. One of the most frequent and relevant MTX side effect is cardiotoxicity. Previously, we found the MTX-naphthoquinoline metabolite (NAPHT) to be less cardiotoxic than MTX.<sup>4</sup> One of the main human MTX metabolites was identified as the dicarboxylic acid resulting from the oxidation of the terminal hydroxyl groups of the side chains.<sup>5</sup> However, its putative cardiotoxicity was not yet assessed. Herein, the synthesis and structure elucidation of the dicarboxylic acid metabolite will be presented.

**Methods:** The total synthesis of the metabolite involved five steps, starting from chrysazin. The enzymatic reaction was accomplished through the horseradish peroxidase (HRP)-catalysed H<sub>2</sub>O<sub>2</sub> and the studies of the oxidative reactions involved sodium tungstate, chromium trioxide, sodium nitrite, and potassium permanganate.

**Results:** In order to obtain the carboxylic derivative, several approaches were undertaken, namely, total synthesis from a commercial available anthraquinone, as well as enzymatic and oxidative reactions from MTX. Different derivatives were obtained. The structure elucidation of the intermediates was established by spectroscopic techniques and the characterization of the dicarboxylic acid metabolite is ongoing.

**Conclusion:** The synthesis of the dicarboxylic acid metabolite was only achieved by the multistep approach. Future work will involve cardiotoxicity studies of this metabolite in H9c2 differentiated cells.

**Acknowledgements:** We thank FCT/MCTES and ERDF through the COMPETE-POFC programme, under the Strategic Funding UID/Multi/04423/2013, the project PTDC/MAR-BIO/4694/2014 (POCI-01-0145-FEDER-016790; 3599-PPCDT) and PTDC/DTP-FTO/1489/2014 (POCI-01-0145-FEDER-016537) in the framework

of PT2020, to INNOVMAR (NORTE-01-0145-FEDER-000035, NOV-ELMAR), supported by NORTE 2020, under PORTUGAL 2020, through ERDF. To Dr. S Cravo for technical assistance.

## References

1. Reis-Mendes AF, et al. *Curr Drug Metab.* 2015;17:75–90.
2. Ehninger G, et al. *Clin Pharmacokinet.* 1990;18:365–80.
3. Shenkenberg TD, et al. *Ann Intern Med.* 1986;105:67–81.
4. Reis-Mendes A, et al. *Arch Toxicol.* 2017;91:1871–90.
5. Chiccarelli FS, et al. *Cancer Res.* 1986;46:4858–61.

<http://dx.doi.org/10.1016/j.pbj.2017.07.107>

## PS179

### Human papillomavirus in the etiology of oropharyngeal carcinoma



T. Kovacic<sup>1,\*</sup>, P. Stefanicka<sup>2</sup>

<sup>1</sup> Faculty of Medicine, Comenius University in Bratislava, Slovakia

<sup>2</sup> Department of Otorhinolaryngology – Head and Neck Surgery, Faculty of Medicine and University Hospital in Bratislava, Slovakia

E-mail address: [tomas.kovacic93@gmail.com](mailto:tomas.kovacic93@gmail.com)

(T. Kovacic).

**Aim:** The aim of this retrospective survey was to investigate the association of HPV status in patients with oropharyngeal cancer with tumour staging and other clinical features.

**Introduction:** Infection by human papillomavirus (HPV) stands for the most frequent viral carcinogenesis in the world. Over-expression of cell oncoprotein p16 is routinely diagnosed by immunohistochemistry (IHC) as the surrogate marker of viral activity.

**Methods:** Records from the oropharyngeal cancer patients treated in the Department of Otorhinolaryngology-Head and Neck Surgery in Bratislava from January 2013 to December 2016 were retrospectively analysed. Patients were divided, according to IHC results on oncoprotein p16, into p16 positive, considered HPV-positive, and p16 negative as HPV-negative. The incidence of oropharyngeal carcinoma, location, T and N staging, age, gender of patients was compared based on HPV status.

**Results:** From 129 oropharyngeal cancer patients with p16 examination were 52 (40%) considered as HPV positive. HPV positive group consisted of 45 (86.5%) men and 7 (13.5%) women. The primary tumour in HPV-positive patients originated from the palatine tonsil and base of the tongue in 96% of cases. The peak of occurrence of HPV-associated carcinoma was found between 50 and 59 years of age. HPV positive tumours were diagnosed in early T stage (T1/2) in 52%. Both HPV positive and negative patients were predominantly diagnosed with advanced-stage cancer, 90.4% in HPV-positive and 87% in HPV-negative group.

**Conclusion:** Early T stage in HPV positive carcinomas was approved, as well as more advanced regional spreading and prevalence of men and non-smokers. Wide variations in numbers of diagnosed patients during years of study may be caused by relatively small size of the studied group. Survey is focusing at HPV status as the most important prognostic factor in oropharyngeal cancer and systematized introduction of HPV status examination as progressive approach to effective and targeted therapy.

**Acknowledgements:** I am thankful to my tutor MUDr. Patrik Stefanicka, PhD. from Department of Otorhinolaryngology - Head and Neck Surgery, Faculty of Medicine and University Hospital in Bratislava.<sup>1,2</sup>

## References

1. Gillison ML, Castellsaqué X, Chaturvedi A. Eurogin Roadmap: comparative epidemiology of HPV infection and associated cancers of the head and neck and cervix. *Int J Cancer.* 2014;134:497–507.
2. Dahlstrom K, Bell D, Hanby D. Socioeconomic characteristics of patients with oropharyngeal carcinoma according to tumor HPV status, patient smoking status, and sexual behavior. *Oral Oncol.* 2015;51:832–8.

<http://dx.doi.org/10.1016/j.pbj.2017.07.108>

## PS181

### The effects of cannabinoids in exemestane-resistant breast cancer cells



C. Almeida<sup>1,2,3,\*</sup>, T. Augusto<sup>3</sup>,  
G. Correia-da-Silva<sup>3,4</sup>, N. Teixeira<sup>4</sup>, C. Amaral<sup>4</sup>

<sup>1</sup> Faculdade de Ciências, Universidade do Porto, Portugal

<sup>2</sup> Instituto de Ciências Biomédicas Abel Salazar, Universidade do Porto, Portugal

<sup>3</sup> Laboratório de Bioquímica, Departamento de Ciências Biológicas, Faculdade de Farmácia, Universidade do Porto, Portugal

<sup>4</sup> UCIBIO, REQUIMTE, Laboratório de Bioquímica, Departamento de Ciências Biológicas, Faculdade de Farmácia, Universidade do Porto

E-mail address: [cristina-almeida96@hotmail.com](mailto:cristina-almeida96@hotmail.com)

(C. Almeida).

**Aim:** Considering that the development of resistance is the main reason for endocrine treatment failure, our group decided to explore the ability of three cannabinoids,  $\Delta^9$ -tetrahydrocannabinol (THC), cannabidiol (CBD) and anandamide (AEA), to reverse resistance to exemestane. The THC and CBD are phytocannabinoids derived from the plant *Cannabis sativa* (marijuana) whereas AEA is an endocannabinoid. For that, it was used LTEDaro cells, a long-term estrogen deprived ER+ breast cancer cell line that mimics resistance to exemestane.<sup>1</sup> These cells were treated with exemestane in combination with two phytocannabinoids, CBD and THC, and the endocannabinoid AEA.

**Introduction:** Exemestane is one of the aromatase inhibitors (AI) used as first line treatment for estrogen-receptor positive breast cancer in post-menopausal women. Exemestane acts by inhibiting aromatase, the enzyme responsible for the conversion of androgens to estrogens<sup>2</sup> and also by promoting apoptosis of breast cancer cells.<sup>3</sup> Nevertheless, despite its therapeutic success, this AI, after prolonged treatment, can induce acquired resistance, which causes tumor relapse. Therefore, it is important to find new strategies to overcome resistance in order to improve breast cancer treatment.

**Methods:** The presence of CB1 and CB2 in LTEDaro cells was confirmed by Western blot analysis and the effects of the combination of cannabinoids with exemestane were evaluated by MTT and LDH assays. Cell morphology was analyzed by Giemsa and Hoechst staining.

**Results:** Our results demonstrate that all the cannabinoids induce a decrease in viability of exemestane-resistant cells, in a dose- and time-dependent manner, without LDH release. These results indicate that the studied cannabinoids, mainly THC and AEA, revert the resistance to exemestane, probably by inducing apoptosis, as observed in Giemsa/Hoechst stain by the presence of typical morphological features of apoptosis.

**Conclusion:** This study highlights the efficacy of using cannabinoids as a potential adjuvant treatment to revert resistance to AIs.

**Acknowledgements:** Amaral C. grant (SFRH/BPD/98304/2013), Augusto T. (BD/128333/2017) and (UID/MULTI/04378/

2013–POCI/01/0145/FERDER/007728); Prof. Shiuan Chen (Beckman Research Institute, USA) for LTEDaro cells.

## References

1. Chen S. An “omics” approach to determine the mechanisms of acquired aromatase inhibitor resistance. *OMICS*. 2011;15:347–52.
2. Sobral AF, Amaral C, Correia-da-Silva G, Teixeira N. Unravelling exemestane: from biology to clinical prospects. *J Steroid Biochem Mol Biol*. 2016;163:1–11.
3. Amaral C, Borges M, Melo S, da Silva ET, Correia-da-Silva G, Teixeira N. Apoptosis and autophagy in breast cancer cells following exemestane treatment. *PLoS ONE*. 2012;7:e42398.

<http://dx.doi.org/10.1016/j.pbj.2017.07.109>

## PS184

### Quantitative structure-property relationship (QSPR) of thiazolidin-4-one derivatives as RTIs of HIV virus



N. Moussavi Harami\*, A. Almasi Rad

*Department of Medicinal Chemistry, Faculty of Pharmacy, Islamic Azad University, Iran*

*E-mail address: [moussavi.nasim@gmail.com](mailto:moussavi.nasim@gmail.com)*

(N. Moussavi Harami).

**Aim:** The aim of this study is to build a quantitative structure-property relationship (QSPR) of 66 thiazolidin-4-one derivatives in order to predict their logP.

**Introduction:** Performing computational drug design is an important step for their synthesis and properties characterizations. In this work, quantitative structure-property relationship (QSPR) of 66 thiazolidin-4-one derivatives was examined in order to predict their logP which is the most commonly used measure of lipophilicity in chemical molecules. These group of compounds act as non-nucleoside reversed transcriptase inhibitors of HIV.

**Methods:** Two different quantum mechanics approaches including HF and DFT were applied for energy minimization of structures and different classes of molecular descriptors including quantum chemical descriptors were generated for prediction of their LogP. Numbers of descriptors which showed high correlation with each other were removed by MATLAB software. The model between structures and their LogP was built for both methods with performing Multiple Linear Regression (MLR) in Spss package

**Results:** Statistical results and application of developed model to the test set demonstrates that the DFT model is reliable with good predictive accuracy. ( $R^2_{cal} = 0.90$ ,  $R^2_{cv} = 0.88$ ) The lack of significant difference between the original and modeled values of logP reveals the validity of the built model which was built with 2D and 3D descriptors. The coefficients of model are statistically significant.

**Conclusion:** QSPR models can be used to predict molecular properties such as LogP. That will be beneficial in drug design processes. In this research, MLR model was built in order to correlate structure of 66 compounds with their LogP. Molecules that were optimized by DFT method showed better correlation than HF method that indicates the accuracy of the built model with 2D and 3D descriptors.

<http://dx.doi.org/10.1016/j.pbj.2017.07.110>

## PS185

### Lung branching morphogenesis, in the chicken model, is accompanied by temporal metabolic changes



H. Fernandes-Silva<sup>1,2,\*</sup>, M.G. Alves<sup>3,4</sup>, J. Correia-Pinto<sup>1,2,5</sup>, P.F. Oliveira<sup>3,4,6,7</sup>, R.S. Moura<sup>1,2</sup>

<sup>1</sup> *Life and Health Sciences Research Institute (ICVS), School of Medicine, University of Minho, 4710-057 Braga, Portugal*

<sup>2</sup> *ICVS/3B's – PT Government Associate Laboratory, 4710-057 Braga/Guimarães, Portugal*

<sup>3</sup> *Unit for Multidisciplinary Research in Biomedicine (UMIB), Institute of Biomedical Sciences Abel Salazar (ICBAS), University of Porto, 4050-313 Porto, Portugal*

<sup>4</sup> *Department of Microscopy, Institute of Biomedical Sciences Abel Salazar (ICBAS), University of Porto, 4050-313 Porto, Portugal*

<sup>5</sup> *Department of Pediatric Surgery, Hospital de Braga, 4710-243 Braga, Portugal*

<sup>6</sup> *i3S – Instituto de Inovação e Investigação em Saúde, University of Porto, 4050-313 Porto, Portugal*

<sup>7</sup> *Department of Genetics, Faculty of Medicine (FMUP), University of Porto*

*E-mail address: [hugomiguelfsilva@gmail.com](mailto:hugomiguelfsilva@gmail.com)*

(H. Fernandes-Silva).

**Aim:** In this work, we characterized, for the first time, the metabolic profile of chick lung branching in early stages of development: b1, b2 and b3 (1, 2 or 3 secondary bronchi, respectively).

**Introduction:** Pulmonary development is a complex process that depends on the activation of conserved signaling pathways that regulate cellular processes such as proliferation, differentiation and migration.<sup>1–3</sup> These cellular processes require high amounts of energy and nutrients to form new biomass.<sup>4,5</sup> However, the metabolic changes that occur during lung branching morphogenesis have not been described so far.

**Methods:** Ex vivo lung explant culture was performed and the medium collected to analyze the production/consumption of metabolic intermediates associated with glucose catabolism (lactate, acetate, alanine), by 1H-NMR. qPCR was performed to assess the expression levels of key enzymes and transporters from the correspondent metabolic pathways.

**Results:** The results showed that the major variations occur from stage b1 to stage b3. In b3 there is an increase in lactate and acetate production. Still, glucose consumption is maintained from b1 to b3 stage, with a concurrent decrease of glucose transporter 3 (glut3) transcript levels. Hexokinase 1 (hk1) levels also decrease in b3 stage (as compared to b2). This phenomenon suggests an increase in the glycolytic efficiency and a shift to lactic acid production (in detriment of mitochondrial respiration). In fact, we observed a decrease on pyruvate dehydrogenase B (pdhB) and an increase in lactate dehydrogenase A (ldhA) expression levels in b3 stage (as compared to b2), while lactate dehydrogenase B (ldhB) levels decrease.

**Conclusion:** This study describes, for the first time, the temporal metabolic changes associated with chick pulmonary branching. It seems that glycolytic efficiency is increased and Krebs cycle metabolism shifts to lactate production along development. Furthermore, acetate and lactate are potentially seen as metabolic biomarkers of lung development.

**Acknowledgements:** This work has been funded by FEDER funds, through the Competitiveness Factors Operational Programme (COMPETE), and by National funds, through the Foundation for Science and Technology (FCT), under the scope of the project POCI-01-0145-FEDER-007038; and by the project NORTE-01-0145-FEDER-000013, supported by the Northern Portugal Regional Operational Programme (NORTE 2020), under the Portugal 2020 Partnership Agreement, through the European Regional Development Fund (FEDER). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## References

- Ornitz DM, et al. Cold Spring Harb Perspect Biol. 2012;4, <http://dx.doi.org/10.1101/cshperspect.a008318>, pii:a008318.
- Moura RS, et al. PLoS ONE. 2014;9:e112388, <http://dx.doi.org/10.1371/journal.pone.0112388>.
- Moura RS, et al. Histochem Cell Biol. 2016;146:457–66, <http://dx.doi.org/10.1007/s00418-016-1448-1>.
- Ito T, et al. Histol Histopathol. 1999;14:895–904.
- Oliveira PF, et al. Med Res Rev. 2015;35:126–51, <http://dx.doi.org/10.1002/med.21325>.

<http://dx.doi.org/10.1016/j.pbj.2017.07.111>

## PS186

### Epigenetic modifications as targets to new therapies for Chronic Lymphocytic leukaemia – A preliminary study



B. Ribau<sup>1,2,\*</sup>, J. Jorge<sup>2,4</sup>, R. Alves<sup>2,3</sup>, P.I. Ribeiro<sup>4,5</sup>, A.C. Gonçalves<sup>2,3,4</sup>, I.M. Carreira<sup>4,5</sup>, A.B. Sarmento-Ribeiro<sup>2,3,4,6</sup>

<sup>1</sup> Department of Chemistry, University of Aveiro, Portugal

<sup>2</sup> Laboratory of Oncobiology and Hematology (LOH), University Clinic of Hematology and Applied Molecular Biology, FMUC, Portugal

<sup>3</sup> Center for Neuroscience and Cell Biology, IBILI (CNC.IBILI), University of Coimbra, Portugal

<sup>4</sup> CIMAGO - Center of Investigation on Environment Genetics and Oncobiology, Faculty of Medicine, University of Coimbra, Portugal

<sup>5</sup> Laboratory of Cytogenetics and Genomics (LCG), Faculty of Medicine, University of Coimbra, Portugal

<sup>6</sup> Clinical Hematology Service, University Hospital of Coimbra, Portugal

E-mail address: [beatriz.ribau@ua.pt](mailto:beatriz.ribau@ua.pt) (B. Ribau).

**Aim:** This study aimed to clarify the involvement of epigenetic modifications in chronic lymphocytic leukaemia development and analyse the therapeutic potential of epigenetic modulators.

**Introduction:** CLL is the most common type of leukaemia found in adults and is an extremely variable and heterogeneous disease. The CLL aetiology is unknown and its natural history is heterogeneous. However, epigenetic modifications may play an important role in CLL.

**Methods:** This study enrolled 18 CLL and 7 controls. To perform primary CLL cultures, peripheral blood mononuclear cells from CLL patients were isolated using Ficoll gradient and incubated with the hypomethylants, Azacytidine and Decitabine, and deacetylase inhibitors, Panobinostat and Vorinostat, in monotherapy (single dose and daily administration) and in combination for 24 h/48 h. The cytotoxic/cytostatic effect of drugs was evaluated by fluorometric microculture cytotoxicity assay (FMCA). Cell death

and cell cycle were determined by flow cytometry using Annexin V and PI/RNase, respectively. CD5 and CD19 antibodies were used to identify normal (CD5<sup>-</sup>/CD19<sup>+</sup>) and neoplastic cells (CD5<sup>+</sup>/CD19<sup>+</sup>). Methylation pattern was determined by MS-MLPA. Data were analysed using univariate approaches.

**Results:** Preliminary results show that patients appear to be more sensitive to Azacytidine and Vorinostat than Decitabine and Panobinostat, on single dose administration. Combination of Panobinostat with Azacytidine and Decitabine induced higher cytotoxicity than single dose. For all drugs, daily administration schedule reduced more effectively cell viability/proliferation than the same doses in single administration. These drugs induced cell death mainly by apoptosis with specificity to neoplastic cells. Moreover, CLL patients had a significant higher methylation frequency of PAX5 (70%), KLLN (80%), WT1 (100%), THBS1 (90%) and GATA5 (90%) gene promoters when compared with controls (all genes demethylated, except MSH6). Furthermore, all CLL patients had at least one methylated gene.

**Conclusion:** The preliminary results suggest that methylation of tumour suppressor genes is a common event in CLL patients and that epigenetic modulators induce a cytotoxic effect, reducing cell viability/proliferation, in a time- and dose-dependent manner. Therefore, these results are promising and encourage further studies in CLL.

<http://dx.doi.org/10.1016/j.pbj.2017.07.112>

## PS191

### Imaging features of brain metastases from testicular cancer



Ana Filipa Pinto<sup>1,\*</sup>, Susana Maria Silva<sup>2,3</sup>, Eduarda Carneiro<sup>4</sup>, Diana Ferreira<sup>4</sup>, Joaquina Maurício<sup>5</sup>, Mavilde Arantes<sup>2,3,4</sup>

<sup>1</sup> Faculty of Medicine of the University of Porto, 4200-319 Porto, Portugal

<sup>2</sup> Unit of Anatomy, Department of Biomedicine, Faculty of Medicine of the University of Porto, 4200-319 Porto, Portugal

<sup>3</sup> Center for Health Technology and Services Research (CINTESIS), 4200-450 Porto, Portugal

<sup>4</sup> Division of Neuroradiology, Radiology Service, Portuguese Institute of Oncology, Porto, Portugal

<sup>5</sup> Medical Oncology Service, Portuguese Institute of Oncology, Porto, Portugal

E-mail address: [anafilipapinto95@gmail.com](mailto:anafilipapinto95@gmail.com) (A.F. Pinto).

**Aim:** Our study evaluated the incidence, imaging characteristics, and prognosis of brain metastases originating from primary testicular tumors.

**Introduction:** Approximately 95% of testicular tumors are testicular germ cell tumors (TGCT).<sup>1</sup> Sertoli cell tumors are rare non-germ cell origin tumors and account for less than 1% of testicular cancer.<sup>2</sup> Brain metastases from germ cell tumors are very uncommon, occurring in less than 2–3% of patients.<sup>3</sup> In testicular cell cancer, it is estimated that the incidence of brain metastases is 1–2% in all TGCT, whereas in advanced stages of TGCT the incidence rises to about 10–15%<sup>4–9</sup>

**Methods:** Case records of testicular tumors patients within the IPO Porto data base from 2006 to 2015 were reviewed to identify patients with testicular tumors and evidence of brain metastases.

**Results:** 368 patients with testicular tumors were identified, with only four having evidence of brain metastases.

Histopathological evaluation revealed that one of the patients had a non-germ cell tumor, a Sertoli cell tumor, while others had mixed germ cell tumors. Half of them had only a single right frontoparietal lesion (21 mm) or right occipital (42 mm), both were heterogeneous in T1WI and T2WI, and with intense and heterogeneous enhancement with gadolinium. The other two patients had multiple lesions. One of them had left frontoparietal (2.2 mm, hyperintense in T1) and right occipital (1.8 mm, hypointense in T1) lesions, both heterogeneous and predominantly hypointense in T2 and T1WI with no enhancement. The other had right temporal (5 mm) and left occipital (11 mm) lesions, both isointense in T1WI and T2WI with intense and homogeneous enhancement. There was no diffusion restriction in all three cases and all four cases were hypointense in T2\*.

**Conclusion:** Although the imaging features of brain metastases differ in some aspects, they all have a hemorrhagic component and a very low survival rate after diagnosis.

## References

- Huyghe E, Matsuda T, Thonneay P. Increasing incidence of testicular cancer worldwide: a review. *J Urol.* 2003;170:5–11.
- Sesterhenn IA, Chevillet J, Woodward PJ, et al. Sex cord/gonadal stromal tumours. In: Eble JN, Sauter G, Epstein JI, Sesterhenn IA, editors. *Pathology and genetics of tumours of the urinary system and male genital organs.* Lyon, France: IARC Press; 2004. p. 250–5.
- Raj S, Parkinson C, Williams M, Mazhar D. Management of brain metastases from germ cell tumors: do we know what we are doing? *Future Oncol.* 2008;4:1–4.
- Raina V, Singh SP, Kamble N, et al. Brain metastasis as the site of relapse in germ cell tumor of testis. *Cancer.* 1993;72:2182–5.
- Bower M, Newlands ES, Holden L, Rustin GJ, Begent RH. Treatment of men with metastatic non-seminomatous germ cell tumours with cyclical POMB/ACE chemotherapy. *Ann Oncol.* 1997;8:477–83.
- Bokemeyer C, Nowak P, Haupt A, et al. Treatment of brain metastases in patients with testicular cancer. *J Clin Oncol.* 1997;15:1449–54.
- Fossa SD, Bokemeyer C, Gerl A, et al. Treatment outcome of patients with brain metastases from malignant germ cell tumors. *Cancer.* 1999;85:988–97.
- Williams SD, Einhorn LH. Brain metastases in disseminated germinal neoplasms: incidence and clinical course. *Cancer.* 1979;44:1514–6.
- Kaye SB, Bagshawe KD, McElwain TJ, Peckham MJ. Brain metastases in malignant teratoma: a review of four years' experience and an assessment of the role of tumour markers. *Br J Cancer.* 1979;39:217–23.

<http://dx.doi.org/10.1016/j.pbj.2017.07.113>

## PS196

### Synthesis and tumor cell growth inhibitory effects of the marine product analogues of fiscalin B

N. Lopes<sup>1,\*,#</sup>, S. Long<sup>1,#</sup>, D. Resende<sup>1,2</sup>,  
A. Kijjoo<sup>2,3</sup>, A. Silva<sup>4</sup>, A. Pina<sup>3,5,6,7</sup>,  
T. Fernández-Marcelo<sup>5,6</sup>, M.H. Vasconcelos<sup>5,6,8</sup>,  
M. Pinto<sup>1,2</sup>, E. Sousa<sup>1,2</sup>

<sup>1</sup> *Laboratório de Química Orgânica e Farmacêutica, Departamento de Ciências Químicas, Faculdade de Farmácia, Universidade do Porto, Portugal*

<sup>2</sup> *Instituto de Ciências Biomédicas Abel Salazar (ICBAS), Universidade do Porto, Portugal*

<sup>3</sup> *CIIMAR – Centro Interdisciplinar de Investigação Marinha e Ambiental, Matosinhos, Portugal*

<sup>4</sup> *Organic Chemistry Group, QOPNA, Department of Chemistry, University of Aveiro, 3810-193, Aveiro, Portugal*

<sup>5</sup> *i3S - Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Porto, Portugal*

<sup>6</sup> *Cancer Drug Resistance Group, IPATIMUP - Institute of Molecular Pathology and Immunology of the University of Porto*

<sup>7</sup> *FCUP – Faculty of Sciences of the University of Porto*

<sup>8</sup> *Department of Biological Sciences, FFUP - Faculty of Pharmacy of the University of Porto*  
E-mail address: [natalia.lopes17@gmail.com](mailto:natalia.lopes17@gmail.com)  
(N. Lopes).

**Aim:** The aim of this work was to synthesize fiscalin B, to pursue the development of a library of derivatives and to investigate the derivatives for their potential antitumor activity.

**Introduction:** Marine organisms provided numerous novel compounds with sensational multiple pharmacological properties. The necessity of novel therapeutics has gain more importance especially because of the resistance associated to the current therapeutics and the inexistent treatments for incurable diseases. Fiscalin B is a fungal metabolite with a pyrazino[2,1-b]quinazoline-3,6-dione system reported to have significant biological activities.

**Methods:** Two methods were studied for synthesis – double cyclization and microwave assisted procedures. First method started with coupling reactions to form tripeptide of tryptophan methyl ester linked to N-Fmoc-valine via anthranilic acid. Then, the dehydrative cyclization was performed using formamide to form the intermediate oxazine. The coupling reaction to form the fiscalin B were achieved after deprotection.<sup>1</sup> The second method is the coupling of anthranilic acid with N-Boc-valine to form Boc-protected benzoxazin-4-one by thermal heating conditions. Then, the addition of tryptophan methyl ester hydrochloride led to 4-quinazoline-3,6-dione scaffold by microwave irradiation.<sup>2</sup> The cell growth inhibitory effect was investigated by the Sulforhodamine B assay.

**Results:** The use of amino acids with different configurations and different side chains or even the derivatization of the existing functional groups were enable the application of this synthetic methodology for a library of fiscalin B analogues. The formation yields of fiscalin B analogues were low, ranging from 3 to 16%. Eight derivatives were tested on non-small cell lung cancer (H460), colon adenocarcinoma (HCT15) and breast cancer (MCF7) human cell lines and showed moderate cytotoxic effects, with GI50 concentrations ranging from 30 to 80  $\mu$ M.

**Conclusion:** Significant differences were obtained between enantiomeric pairs.

**Acknowledgements:** To national funds provided by FCT, ERDF, and COMPETE under the projects PEst-C/MAR/LA0015/2013, QOPNA (FCT UID/QUI/00062/2013), PTDC/MAR-BIO/4694/2014 (POCI-01-0145-FEDER-016790), PTDC/AAG-TEC/0739/2014 (POCI-01-0145-FEDER-016793), and INNOVMAR, reference NORTE-01-0145-FEDER-000035, Research Line NOVELMAR and grant reference NOVELMAR/BPD\_2/2016-019. To University of Aveiro and FCT/MEC for the financial support to the QOPNA research project (UID/QUI/00062/2013) financed by national funds and co-financed by FEDER under the PT2020, and to the Portuguese NMR Network. S.L. thanks Erasmus Mundus Action 2 (LOTUS+, LP15DF0205) for full PhD scholarship.

## References

- Wang H, Ganesan A. Total synthesis of the fumiquinazoline alkaloids: solution-phase studies. *J Organ Chem.* 2000;65:1022–30.
- Liu JF, Ye P, Zhang B, Bi G, Sargent K, Yu L, et al. Three-component one-pot total syntheses of gyantrypine, fumiquinazoline F, and fiscalin B promoted by microwave irradiation. *J Organ Chem.* 2005;70:6339–45.

<http://dx.doi.org/10.1016/j.pbj.2017.07.114>

\* Both authors contributed equally.



## PS207

**Heterocyclic chalcone derivatives: Synthesis and biological activity evaluation**

C. Machado<sup>1,\*</sup>, P. Pinto<sup>2</sup>, P. Silva<sup>3,4,5</sup>,  
D. Almeida<sup>5</sup>, J. Moreira<sup>1,6</sup>, M. Pinto<sup>1,2,3</sup>,  
H. Bousbaa<sup>5,6</sup>, H. Cidade<sup>1,6</sup>

<sup>1</sup> Laboratório de Química Orgânica e Farmacêutica, Departamento de Ciências Químicas, Faculdade de Farmácia, Universidade do Porto, Portugal

<sup>2</sup> Laboratório de Química Farmacêutica, Faculdade de Farmácia, Universidade de Coimbra, Portugal

<sup>3</sup> Center for Biomedical Research, CBMR, University of Algarve, Faro 8005-139, Portugal

<sup>4</sup> Departamento Ciências Biomédicas e Medicina, University of Algarve, Faro, Portugal

<sup>5</sup> CESPU, Instituto de Investigação e Formação Avançada em Ciências e Tecnologias da Saúde, IINFACTS, 4585-116 Gandra PRD, Portugal

<sup>6</sup> Centro Interdisciplinar de Investigação Marinha e Ambiental (CIIMAR/CIMAR), Universidade do Porto, Portugal

E-mail address: [cmmariana3@hotmail.com](mailto:cmmariana3@hotmail.com) (C. Machado).

**Aim:** Synthesis of new heterocyclic chalcone derivatives with promising antitumor activity.

**Introduction:** Natural chalcones have been intensively studied for their wide range of biological activities, namely antitumor.<sup>1</sup> Possessing two electrophilic reactive centers at  $\alpha,\beta$ -unsaturated ketone group, chalcone derivatives can participate in addition reactions leading to the synthesis of promising bioactive compounds with a more rigid structure, like isoxazoles and pyrazoles.<sup>2</sup>

**Methods:** Chalcones were synthesized by base catalysed Claisen Schmidt condensation via microwave assisted organic synthesis (MAOS). The antiproliferative activity was assessed using sulforhodamine B assay.<sup>3</sup>

**Results:** Seventeen chalcone derivatives were synthesized and identified as having in vitro growth inhibitory activity on three human tumor cell lines from breast, lung and melanoma (MCF-7, NCI-H460, and A375-C5).

**Conclusion:** Most of the synthesized chalcones revealed to be promising growth inhibitors of human tumor cell lines. The molecular mechanisms involved in their antiproliferative effect are being evaluated.

**Acknowledgements:** This research was partially supported through national funds provided by FCT, ERDF and COMPETE under the Strategic Funding UID/Multi/04423/2013, and the projects PTDC/MAR-BIO/4694/2014 (POCI-01-0145-FEDER-016790), PTDC/DTPFTO/1981/2014 (POCI-01-0145-FEDER-016581), and PTDC/AAGTEC/0739/2014 (POCI-01-0145-FEDER-016793) in the framework of the programme PT2020 as well as by the project INNOVMAR-Innovation and Sustainability in the Management and Exploitation of Marine Resources (NORTE-01-0145-FEDER-000035, within Research Line NOVELMAR), supported by NORTE 2020, under the PORTUGAL 2020 Partnership Agreement, through the ERDF. Patrícia M.A. Silva is a PhD fellowship holder from FCT (SFRH/BD/90744/2012).

**References**

1. Mahapatra DK, et al. *Eur J Med Chem.* 2015;98:69–114.
2. Albuquerque HMT, et al. *Curr Organic Chem.* 2014;18:2750–75.
3. Neves M, et al. *Bioorganic Med Chem.* 2012;20:25–33.

<http://dx.doi.org/10.1016/j.pbj.2017.07.115>

## PS209

**A posttranslational modification in histones as prognostic/predictive marker in Estrogen-Positive Breast Cancer**

S. Lobo<sup>1,2,\*</sup>, M. Fontes-Sousa<sup>2,3</sup>, S. Salta<sup>2</sup>,  
P. Lopes<sup>2,4</sup>, J. Lobo<sup>2,4</sup>, S. Sousa<sup>3</sup>, R. Henrique<sup>2,4,5</sup>,  
C. Jerónimo<sup>2,5</sup>

<sup>1</sup> Faculty of Science – University of Porto (FCUP-UP), Porto, Portugal

<sup>2</sup> Cancer Biology and Epigenetics Group, IPO Porto Research Center (CI-IPOP), Portuguese Oncology Institute of Porto (IPO Porto), Porto, Portugal

<sup>3</sup> Department of Medical Oncology; Portuguese Oncology Institute of Porto, Portugal

<sup>4</sup> Department of Pathology, Portuguese Oncology Institute of Porto, Porto, Portugal

<sup>5</sup> Department of Pathology and Molecular Immunology, Institute of Biomedical Sciences Abel Salazar - University of Porto (ICBAS-UP), Porto, Portugal

E-mail address: [silvana.lobo.sousa@live.com.pt](mailto:silvana.lobo.sousa@live.com.pt) (S. Lobo).

**Aim:** This work aims to evaluate H3K27me3 expression in luminal-like breast tumors, using immunohistochemistry assay, to assess the prognostic value of this epigenetic alterations in estrogen positive breast cancer (BrC).

**Introduction:** BrC is the second most incident cancer worldwide. In Portugal, in 2012, BrC was simultaneously the leading cancer in incidence and mortality in women.<sup>1</sup> Around 70% of all BrC are hormone-receptor positive, that is the major part of breast tumors is luminal-like.<sup>2</sup> H3K27m3 is a gene repression marker<sup>3,4</sup> and is associated with gene silencing, playing a crucial role in cell proliferation and differentiation.<sup>3</sup> H3K27me3 may have some clinical value in several types of cancer since it can be used as a biomarker. This histone modification has been associated with poor prognosis of some BrC subtypes.<sup>5</sup>

**Methods:** It was used a cohort of BrC patients of the Portuguese Oncology Institution of Porto (IPO-Porto), diagnosed between 1994 and 2002. A total of 102 luminal-like tumor cases were assessed by immunohistochemistry, to H3K27me3 expression. To verify the prognostic value of H3K27me3 levels, Cox regression with a log rank test was performed for both disease-specific and disease-free survival.

**Results:** Through the result analysis, it was established that only tumor grade ( $p=0.021$ ) was significant associated with disease-specific survival. Nevertheless, both luminal subtype ( $p=0.016$ ) and H3K27me3 expression ( $p=0.012$ ) were significantly associated with disease-free survival. Indeed, H3K27me3 high expression is associated with higher recurrence risk, especially in Luminal A.

**Conclusion:** We could confirm the prognostic value of H3K27me3 expression in luminal A subtype BrC patients. Therefore, higher H3K27me3 expression in luminal A tumors is associated with a greater probability of recurrence.

However, studies in larger cohorts are mandatory to validate its clinical utility.

**Acknowledgements:** This study was funded by a grant of the Research Centre of Portuguese Oncology Institute of Porto (CI-IPOP-74-2016).

**References**

1. Ferlay JSI, Ervik M, Dikshit R, Eser S, Mathers C, Rebelo M, et al. GLOBOCAN 2012 v1.0, cancer incidence and mortality worldwide: IARC CancerBase No. 11. Lyon, France: International Agency for Research on Cancer; 2013.

- Senkus E, Kyriakides S, Ohno S, Penault-Llorca F, Poortmans P, Rutgers E, et al. Primary breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2015;26 Suppl. 5, v8-30.
- Chase A, Cross NC. Aberrations of EZH2 in cancer. *Clin Cancer Res*. 2011;17:2613–8.
- Yoo KH, Hennighausen L. EZH2 methyltransferase and H3K27 methylation in breast cancer. *Int J Biol Sci*. 2012;8:59–65.
- Hayashi A, Yamauchi N, Shibahara J, Kimura H, Morikawa T, Ishikawa S, et al. Concurrent activation of acetylation and tri-methylation of H3K27 in a subset of hepatocellular carcinoma with aggressive behavior. *PLoS ONE*. 2014;9:e91330.

<http://dx.doi.org/10.1016/j.pbj.2017.07.116>

## PS212

### Is P-glycoprotein relevant for the release of microvesicles by tumor cells?

I. Castro<sup>1,2,3,\*</sup>, C.P.R. Xavier<sup>1,2</sup>, M.H. Vasconcelos<sup>1,2,4</sup>

<sup>1</sup> i3S – Instituto de Investigação e Inovação em Saúde, Universidade do Porto, 4200-135 Porto, Portugal

<sup>2</sup> Cancer Drug Resistance Group, IPATIMUP – Institute of Molecular Pathology and Immunology of the University of Porto, 4200-465 Porto, Portugal

<sup>3</sup> FMUP – Faculty of Medicine of the University of Porto, 4200-319 Porto, Portugal

<sup>4</sup> FFUP – Faculty of Pharmacy of the University of Porto, 4050-313 Porto, Portugal

E-mail address: [msilva@ipatimup.pt](mailto:msilva@ipatimup.pt)

(I. Castro).

**Aim:** In this study, we aimed to verify if MDR cells without expression of P-gp also produced more microvesicles and less exosomes than their DS counterpart cells.

**Introduction:** Cancer multidrug resistance (MDR) is a major cause of chemotherapy failure and is highly associated with overexpression of drug-efflux pumps such as P-glycoprotein (P-gp). The identification of mechanisms specific of P-gp overexpressing cells may contribute to the identification of biomarkers of MDR.

It was recently discovered that a drug-resistant phenotype may be horizontally transferred from drug-resistant (DR) to drug-sensitive (DS) cells, mediated by the cargo of extracellular vesicles (EVs) released by DR cells and captured by DS cells. These EVs include smaller exosomes and larger microvesicles. Our previous work showed that MDR cells with overexpression of P-gp released more microvesicles than exosomes, unlike their DS counterpart cells. However, it is not known if this phenomenon is restricted to MDR cells with overexpression of P-gp or if it is extensive to all DR cells (with other mechanism of drug resistance).

**Methods:** Drug-response curves of MDR and DS counterpart cells were obtained, using resazurin and trypan blue assays, to confirm the resistant or sensitive phenotype of the cell lines. Confirmation of their P-gp status was possible by Western-Blot. EVs released by both DS and MDR cells were isolated by ultracentrifugation and characterized by transmission electron microscopy, dynamic light scattering, nanoparticle tracking analysis and Western blot analysis.

**Results:** We confirmed that MDR cells without expression of P-gp release EVs with similar sizes to the ones released by their DS counterparts.

**Conclusion:** So, P-gp may be associated with the release of larger EVs by MDR cells. These results will be further confirmed by characterizing the EVs released by P-gp overexpressing MDR cell lines following downregulation of P-gp expression and the EVs released by DS cell lines following transfection of P-gp.<sup>1-4</sup>



## References

- Chen Z, et al. Mammalian drug efflux transporters of the ATP binding cassette (ABC) family in multidrug resistance: a review of the past decade. *Cancer Lett*. 2016;370:153–64.
- Sousa D, Lima RT, Vasconcelos MH. Intercellular transfer of cancer drug resistance traits by extracellular vesicles. *Trends Mol Med*. 2015;21:595–608.
- Lopes-Rodrigues V, et al. Data supporting the shedding of larger extracellular vesicles by multidrug resistant tumour cells. *Data Brief*. 2016;6:1023–7.
- Lopes-Rodrigues V, et al. Multidrug resistant tumour cells shed more microvesicle-like EVs and less exosomes than their drug-sensitive counterpart cells. *Biochim Biophys Acta*. 2016;1860:618–27.

<http://dx.doi.org/10.1016/j.pbj.2017.07.117>

## PS215

### Uterine protein oxidative modifications may condition trophoblast function



S. Mendes<sup>1,2,\*</sup>, A.I. Soares<sup>1,2</sup>, S. Silveira<sup>1</sup>, L. Matos<sup>1,2</sup>, L. Guedes-Martins<sup>2,3</sup>, J. Saraiva<sup>2,3,4</sup>, H. Almeida<sup>1,2,5</sup>, E. Silva<sup>1,2</sup>

<sup>1</sup> I3s; IBMC and FMUP, Portugal

<sup>2</sup> FMDUP, Portugal

<sup>3</sup> Centro Hospitalar do Porto EPE, Portugal

<sup>4</sup> Private Hospital Trofa, Portugal

<sup>5</sup> Obstetrics-Gynecology, Hospital-CUF Porto, Portugal

E-mail address: [saramendes313@gmail.com](mailto:saramendes313@gmail.com)

(S. Mendes).

**Aim:** Evaluate whether protein carbonylation resulting from uterine altered redox imbalance interferes with extravillous trophoblast viability.

**Introduction:** Local redox homeostasis is believed to have a pivot role in uterine transformation necessary for blastocyst implantation and placenta development. By contrast, redox status imbalance plays a role in deficient placentation and the development of pregnancy-related complications (e.g. preeclampsia or gestational diabetes) with increased incidence in older women. Thus, it was hypothesized that at an older reproductive age, loss of redox homeostasis is a contributor to disruption of foetal/placental interactions and the development of such complications.

**Methods:** Uterine human samples were collected at delivery by elective caesarean section. The protocol was approved by the ethical committee of “Centro Materno-Infantil do Porto”, volunteers gave written consent to be included in the study. Total protein carbonylation was detected by oxyblot and protein expression was quantified by western blotting. Specific protein carbonylation was verified by immunoprecipitation. Albumin was carbonylated using hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), followed by dialysis, and western blotting to confirm carbonylated albumin. HST-8SV neo extravillous trophoblasts were treated with carbonylated/non-carbonylated albumin, followed by cell viability assay. A *P* value less than 0.05 was assumed to denote significant difference.

**Results:** At the placental site, carbonylated albumin normalized to total albumin expression showed a positive and significant association with maternal age. ( $r=0.6909$ ,  $P=0.0021$ ) In vitro, carbonylated albumin displayed a cytotoxic effect, at concentrations ranging from 10 to 100 µg/ml. Lower concentrations did not affect trophoblast viability.

**Conclusion:** Uterine aging is accompanied by selective albumin oxidative modifications, which appears to interfere with trophoblast ability to invade and transform the maternal placental site.

<http://dx.doi.org/10.1016/j.pbj.2017.07.118>

## PS223

### Advances on photodynamic therapy through new pyridine-fused diphenylchlorins as photosensitizers for melanoma treatment



J. Dias-Ferreira<sup>1,2,\*</sup>, Nelson A.M. Pereira<sup>3</sup>,  
Mafalda Laranjo<sup>1,4,5</sup>, Marta Pineiro<sup>3</sup>,  
João Casalta-Lopes<sup>1,4,6</sup>,  
Ana Margarida Abrantes<sup>1,4,5</sup>,  
Teresa M.V.D. Pinho e Melo<sup>3</sup>,  
Maria Filomena Botelho<sup>1,4,5</sup>

<sup>1</sup> Biophysics Unit, Faculty of Medicine of University of Coimbra, Azinhaga de Santa Comba, Celas 3004-548, Coimbra, Portugal

<sup>2</sup> Faculty of Pharmacy of University of Coimbra, Azinhaga de Santa Comba, Celas 3004-548, Coimbra, Portugal

<sup>3</sup> CQC and Department of Chemistry, University of Coimbra, 3004-535 Coimbra, Portugal

<sup>4</sup> CIMAGO –Center of Investigation in Environment, Genetics and Oncobiology, Faculty of Medicine of University of Coimbra, Azinhaga de Santa Comba, Celas, 3004-548 Coimbra, Portugal

<sup>5</sup> CNC.IBILI, University of Coimbra, 3004-535 Coimbra, Portugal

<sup>6</sup> Serviço de Radioterapia, Centro Hospitalar e Universitário de Coimbra, Praceta Mota Pinto, 3000-075 Coimbra, Portugal

E-mail address: [j.dias.ferreira@outlook.pt](mailto:j.dias.ferreira@outlook.pt) (J. Dias-Ferreira).

**Aim:** Assessment of cytotoxicity of four new photosensitizers intended for photodynamic therapy (PDT) against melanoma cells (A375 cells).

**Introduction:** Melanoma is the rarest form of skin cancer. PDT combines a photosensitizer with light culminating in the production of reactive oxygen species leading to cellular death. A new type of stable 4,5,6,7-tetrahydropyrazolo[1,5-a]pyridine-fused tetraphenylchlorins<sup>1</sup>, proved to be very active as photodynamic agents. Thus, looking for a new generation photosensitizers with optimized properties for PDT we synthesized new diphenylchlorins.

**Methods:** The human melanoma cell line A375 was seeded in 48 well plates. The photosensitizers NAMP103A, NAMP103B (the tetraphenylchlorins monoester), NAMP263A and NAMP263B (the tetraphenylchlorins alcohol) were administered ranging 5 nM to 10 μM. Irradiation was performed after 24 h ( $\lambda < 560$  nm). MTT and SRB assays as well as flow-cytometry were performed 24 h after the PDT.

**Results:** MTT assay results allowed to obtain dose-response curves and to calculate the concentration that inhibits the cultures by 50% (IC<sub>50</sub>). Phototoxicity (10J) was dependent on the chlorines concentration. Moreover, NAMP263B was significantly more cytotoxic than NAMP103A ( $p=0.037$ ) and NAMP103B ( $p=0.042$ ). From SRB assay we verified that with a 125 nM concentration the NAMP103A, NAMP103B, NAMP263A and NAMP263B produce a cellular viability of 36.9%; 33.2%; 18.3% and 18.8%, respectively. Flow cytometry studies confirmed the decrease of viability associated with cell death by apoptosis and necrosis. Loss of mitochondrial membrane potential, apoptosis hallmark, was also observed. An imbalance of ROS, namely superoxide anion and peroxides, was observed for all photosensitizers studied with an exhaustion of antioxidative intracellular defenses (GSH).

At low PS concentrations (5 nM), metabolic activity was variable with light energy (5J, 10J and 20J) with lower values for higher

fluence. Dark toxicity studies revealed photosensitizer dependence of irradiation.

**Conclusion:** We hereby conclude that the photosensitizers are indeed very promising, which rouses plans for following proceedings to verify in vivo outcome.

**Acknowledgements:** FCT Portugal: Project PTDC/QEQ-MED/0262/2014, Strategic Project Pest CNC.IBILI UID/NEU/04539/2013 and Strategic Project Coimbra Chemistry Centre UID/QUI/00313/2013, COMPETE-FEDER.<sup>1,2</sup>

### References

1. Eur J Org Chem. 2011:3970–9.
2. Eur J Med Chem. 2015:374–80.

<http://dx.doi.org/10.1016/j.pbj.2017.07.119>

## PS226

### Angiogenesis and inflammation at the crossroads between diabetes and cancer



R. Rocha<sup>1,2,\*</sup>, I. Rodrigues<sup>1</sup>, I. Gullo<sup>3,4,5,6</sup>,  
G. Gonçalves<sup>3,4</sup>, J. Pedro<sup>7</sup>, D. Carvalho<sup>4,7</sup>,  
F. Carneiro<sup>3,4,5,6</sup>, R. Soares<sup>1,4</sup>, S. Andrade<sup>1,3,4</sup>

<sup>1</sup> Unit of Biochemistry, Department of Biomedicine, Faculty of Medicine, University of Porto, Portugal

<sup>2</sup> Instituto de Ciências Biomédicas Abel Salazar, University of Porto, Portugal

<sup>3</sup> Institute of Molecular Pathology and Immunology at the University of Porto (IPATIMUP), Porto, Portugal

<sup>4</sup> Instituto de Investigação e Inovação em Saúde (i3S), University of Porto, Porto, Portugal

<sup>5</sup> Department of Pathology, Centro Hospitalar de São João, Porto, Portugal

<sup>6</sup> Department of Pathology, Faculty of Medicine of the University of Porto (FMUP), Porto, Portugal

<sup>7</sup> Department of Endocrinology, Centro Hospitalar de São João, Porto, Portugal

E-mail address: [anaritarocha@ua.pt](mailto:anaritarocha@ua.pt) (R. Rocha).

**Aim:** To study fibrosis, angiogenesis, oxidative stress and inflammation markers in diabetic and non-diabetic patients with gastric cancer (GC).

**Introduction:** Type 2 Diabetes mellitus (DM2) is a major health problem, with 415 million people diagnosed worldwide.<sup>1</sup> Evidence regarding its association with various types of cancer has been reported, including GC.<sup>2</sup> Some hypotheses have been suggested to explain how DM2 could enhance the risk of cancer development, such as hyperglycemia, hyperinsulinemia, oxidative stress, vascular disturbances and a chronic low inflammation state.<sup>3–5</sup>

Gastric cancer (GC) is the fifth most common cancer worldwide and ranks as the third leading cause of cancer-related death.<sup>6</sup> GC is frequently associated with infection by *Helicobacter pylori* and inflammation plays a central role in the carcinogenic process. Such chronic inflammatory state, linked with angiogenesis imbalance, oxidative stress and metabolic signaling, suggests that also DM2 might be a major risk factor in initiation and progression of GC, demanding further investigation.

**Methods:** A series of GC from DM2 ( $n=22$ ) and nonDM2 ( $n=21$ ) patients were studied. Immunohistochemistry (IHC) using antibodies against CD31 and 3-Nitrotyrosine was performed, to assess density of vessels and oxidative stress status. Histochemical staining with Sirius red was performed to determine the percentage of fibrosis in the tumor and non-neoplastic adjacent mucosa. Based on assessment of tumor inflammatory cell infiltrate and tumor stroma

percentage, a semi-quantitative evaluation of Glasgow Microenvironment Score<sup>7</sup> was performed. Also, Glasgow Prognostic Score, that is widely known as a systemic inflammatory-based marker, was determined for each patient.<sup>8</sup>

**Results:** Diabetic patients presented a significant higher glycaemia than the control patients ( $190.1 \pm 13.6$  mg/dL vs  $98.2 \pm 3.6$  mg/dL,  $p < 0.001$ , respectively). Decreased survival rates were observed in diabetic patients ( $611.5$  vs  $916.0$ ,  $p = ns$ ). Tumours exhibited increased fibrosis relatively to the adjacent mucosa in both groups and diabetic patients ( $N: 9.362 \pm 1.337$ ;  $T: 12.29 \pm 1.407$ ) presented higher fibrosis levels than the non-diabetic patients ( $N: 7.165 \pm 1.017$ ;  $T: 10.97 \pm 1.076$ ).

**Conclusion:** Expected results: Identifying the distinct features that characterize GC of DM2 patients compared to nondiabetic patients (namely fibrosis, angiogenesis, inflammation, and oxidative stress biomarkers) will enable to study this subset of GC patients and unravel key mechanisms behind the relationship between DM2 and GC.

**Acknowledgements:** Funding: This work was supported by the project Diabetes & obesity at the crossroads between Oncological and Cardiovascular diseases – a system analysis NETWORK towards precision medicine (DOCnet) – A multi-omics approach to decipher diabetes-related molecular targets in cancer: a step towards precision medicine. NORTE2020 – “Programa Operacional Regional do Norte” (NORTE-01-0145-FEDER-000003) (Jan 2016–Dez2018).

## References

- Ogurtsova K, et al. IDF Diabetes Atlas: Global estimates for the prevalence of diabetes for 2015 and 2040. *Diabetes Res Clin Pract.* 2017;128:40–50.
- Vigneri R. Diabetes: diabetes therapy and cancer risk. *Nat Rev Endocrinol.* 2009;5:651–2.
- Sekikawa A, et al. Diabetes mellitus increases the risk of early gastric cancer development. *Eur J Cancer.* 2014;50:2065–71.
- Ikeda F, Kiyohara Y. Helicobacter pylori infection and Hyperglycemia/Diabetes are associated with an increased risk of Gastric Cancer. *Gan To Kagaku Ryoho.* 2015;42:529–33.
- Shen Z, et al. Glycemic changes after gastrectomy in non-morbidly obese patients with gastric cancer and diabetes. *Hepatogastroenterology.* 2015;62:245–50.
- Ferlay JSI, Ervik M, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and mortality worldwide: IARC CancerBase No. 11. GLOBOCAN 2012 v1.0; 2013. Available from: <http://globocan.iarc.fr> [cited 24.01.17].
- Zhou ZH, et al. The prognostic value and pathobiological significance of Glasgow microenvironment score in gastric cancer. *J Cancer Res Clin Oncol.* 2017;143:883–94.
- Zhang CX, et al. Association between pretreatment Glasgow prognostic score and gastric cancer survival and clinicopathological features: a meta-analysis. *Oncol Targets Ther.* 2016;9:3883–91.

<http://dx.doi.org/10.1016/j.pbj.2017.07.120>

## PS229

### Circulating EVs for AML minimal residual disease biomarkers detection

P.C. Nunes<sup>1,\*</sup>, H.R. Caires<sup>2</sup>,  
M.A. Sobrinho-Simões<sup>3</sup>, M.H. Vasconcelos<sup>4</sup>

<sup>1</sup> Cancer Drug Resistance Group, IPATIMUP – Institute of Molecular Pathology and Immunology of the University of Porto, Portugal; i3S – Instituto de Investigação e Inovação em Saúde, University of Porto, Portugal; ICBAS-UP – Institute of Biomedical Sciences Abel Salazar of the University of Porto, Portugal

<sup>2</sup> Cancer Drug Resistance Group, IPATIMUP – Institute of Molecular Pathology and Immunology of the University of Porto, Portugal; i3S – Instituto de Investigação e Inovação em Saúde, University of Porto, Portugal



<sup>3</sup> Cancer Drug Resistance Group, IPATIMUP – Institute of Molecular Pathology and Immunology of the University of Porto, Portugal; i3S – Instituto de Investigação e Inovação em Saúde, University of Porto, Portugal; FMUP – Faculty of Medicine of the University of Porto, Portugal; HSJ – Hospital de São João, Porto, Portugal

<sup>4</sup> Cancer Drug Resistance Group, IPATIMUP – Institute of Molecular Pathology and Immunology of the University of Porto, Portugal; i3S – Instituto de Investigação e Inovação em Saúde, University of Porto, Portugal; FFUP – Faculty of Pharmacy of the University of Porto, Portugal  
E-mail address: [pnunes@ipatimup.pt](mailto:pnunes@ipatimup.pt)  
(P.C. Nunes).

**Aim:** We propose to evaluate the feasibility of a peripheral blood EV-based liquid biopsy method for AML disease monitoring in real time with molecular precision.

**Introduction:** Acute myeloid leukemia (AML) is a hematopoietic stem cell disorder with high mortality rate mainly due to the high frequency of post-treatment relapse. Minimal residual disease (MRD) determination in AML patients receiving treatment is useful to assess chemotherapy response and predict relapse. One approach to upgrade the current invasive MRD monitoring (traditionally based on bone marrow aspirates/biopsies) is to use methods that identify cancer-associated biomarkers in patients' blood. Recently, extracellular vesicles (EVs) have been increasingly recognized as a potential source of biomarkers, since the levels of EVs are markedly increased in cancer patients' blood and those EVs potentially carry molecular signatures associated with specific cancer phenotypes.

**Methods:** The profile of EVs isolated from AML patients' blood plasma collected from paired AML diagnostic and complete remission samples is being compared and correlated with clinical data. A size-exclusion chromatography (SEC) method was optimized to isolate the plasmatic EVs. The EVs profile is then characterized according to their size, plasmatic concentration, morphology and protein content.

**Results:** EVs with decreasing size were successfully isolated between SEC fractions 3 to 6, with a size ranging from 300 nm to 30 nm, respectively. Fraction 7 presented the smaller EVs, although mixed with some plasmatic protein contaminants. The expression of EVs markers such as CD63, HSP70 or Syntenin-1 was confirmed and allow to distinguish EV subpopulations between fractions 3 to 7. The expression of leukemia-specific markers is currently being studied in the EVs isolated from the paired AML blood samples.

**Conclusion:** The presented EV-based liquid biopsy proposed method for AML monitoring could unravel biomarkers for diagnostic and prognostic purposes in AML patients.

<http://dx.doi.org/10.1016/j.pbj.2017.07.121>

## PS232

### The association of Generalized Epilepsy with Febrile Seizures plus (GEFS+) with FEB1 gene: A new insight to the etiology of GEFS+

Ali Rafati<sup>1,\*</sup>, Shahram Teimourian<sup>2</sup>

<sup>1</sup> Student Research Committee, School of Medicine, Iran University of Medical Sciences, Tehran, Iran  
<sup>2</sup> Department of Medical Genetics, Iran University of Medical Sciences Tehran Iran  
E-mail address: [rafatali1995@gmail.com](mailto:rafatali1995@gmail.com)  
(A. Rafati).



**Aim:** We aimed to carry out a linkage analysis in 6 Iranian families to find an association between the FEB1 and GEFS+.

**Introduction:** Generalized Epilepsy with Febrile Seizures plus (GEFS+), is a group of genetic epilepsy syndromes, likely to commence in the first year of life, in which, patient presents with febrile and tonic-clonic seizures. GEFS+ is associated with an autosomal dominant pattern and is caused by mutations in SCN1B which encodes the beta 1 subunit of sodium channels.

**Methods:** We conducted a case–control study in January 2017, with 6 families, with a total of 35 members entering the study with convenience sampling method, within which, 12 members were as the case group, diagnosed with autosomal dominant GEFS+, hospitalized in Ali Asghar Children's hospital, Iran University of Medical Sciences. 23 family members with no diagnosed GEFS+ were as the control group. Written consent was obtained from all family members according to the protocols of the ethics committee of the university. Afterwards, using D8S533 marker for FEB1 gene, with a Logarithms of Odds (LOD) of 3.16, two-point linkage analysis and haplotype reconstruction was carried out using MLINK program and Simwalk2 respectively.

**Results:** Haplotype reconstruction analysis in the case group revealed a haplotype associated with GEFS+. However, in the control group, not such an haplotype was seen and the difference between two groups was significant ( $p < 0.05$ ).

**Conclusion:** In this study we reported a strong linkage between GEFS+ and FEB1 gene. This may clarify the etiology underlying GEFS+ and gives us chances in GEFS+ screening using FEB1.

<http://dx.doi.org/10.1016/j.pbj.2017.07.122>

Physiology & Immunology Poster Session  
Thursday, September 14th, 16h00

#### PS037

##### Genetical variability of VP1 gene of BK virus in HIV-infected patients



O. Lijeskić<sup>1,\*</sup>, S. Leštarević<sup>1</sup>, D. Karalić<sup>2</sup>

<sup>1</sup> School of Medicine, University of Belgrade, Portugal

<sup>2</sup> Institute of Microbiology and Immunology, School of Medicine, University of Belgrade, Portugal

E-mail address: [oljalol2@gmail.com](mailto:oljalol2@gmail.com)

(O. Lijeskić).

**Aim:** The aims of this study were: to determine the prevalence of BK viruria in HIV-infected patients, to determine the distribution of BKV subtypes and the presence of nucleotide substitutions and mutations in the VP1 gene of BKV isolates.

**Introduction:** A broad range of diseases associated with BK virus (BKV) such as nephritis, haemorrhagic cystitis, encephalitis, retinitis and pneumonia have been reported in HIV-infected patients over the last few years. However, these diseases do not occur in all HIV-infected patients, suggesting that other factors, such as genetic variability of BKV, can contribute to their occurrence. Mutations in the BC loop of the VP1 gene may lead to selection of more aggressive variants of BKV.

**Methods:** The study included 50 HIV-infected patients. Sem-nested PCR was used for amplification of 290-nt fragment within the VP1 gene and all the positive PCR products were then directly sequenced. The sequence analysis was performed by using the appropriate bioinformatics tools.

**Results:** The frequency of BK viruria in HIV-infected patients was 56%. The predominant BKV subtype was I, followed by subtype IV. The majority of mutations were located within BC loop of VP1. The most frequent mutation was E82D.

**Conclusion:** The increased levels of BKV replication are associated with a higher incidence of mutations in the BC loop of VP1, and mutations in this domain may lead to changed tropism and the selection of more aggressive variants of BKV. Further studies are needed in order to select the patients with a higher risk of developing BKV associated-diseases.

<http://dx.doi.org/10.1016/j.pbj.2017.07.123>

#### PS182

##### Cellular interaction in central and peripheral immune organs due to chronic light stress



Bocharova Tetiana

Kharkiv National Medical University, Ukraine

E-mail address: [bochata@ukr.net](mailto:bochata@ukr.net).

**Aim:** Study cellular interaction in central and peripheral immune organs at prolonged all-day illumination in an experiment on rabbits.

**Introduction:** Prolonged all-day illumination is considered nowadays as one of the stress-factors for the living organism and causes malfunctions of the neuroendocrine system and may initial immune dysfunction.

**Methods:** Experimental rabbits ( $n = 10$ ) were in artificial lighting in the day and electric lighting at night during 12 months. Control animals ( $n = 5$ ) were kept in natural day and night lighting conditions. Cell density in immune organs (thymus, bone marrow, spleen) were measured in surface area which was determined by a rectangle  $100 \times 100 \mu\text{m}$ . The results were processed with standard statistical methods and reported as mean  $\pm$  standard deviation (SD).

**Results:** The cell density in the thymus and the bone marrow was decreased: in the cortex of the thymus was  $359.6 \pm 2.9$ , in the medullar part –  $250.8 \pm 2.9$ , in the bone marrow –  $176.4 \pm 2.9$  (cells in  $100 \times 100 \mu\text{m}$ ). An intensified formation of the connective tissue, an increasing of involutive processes and degenerative changes of lymphocytes were microscopically found in the spleen and the thymus. The cell density in the spleen was decreased too: in T zone –  $235.8 \pm 3.7$ , in B-zone –  $159.5 \pm 1.9$  (cells in  $100 \mu\text{m} \times 100 \mu\text{m}$ ). The causes of these changes, probably, may be decrease of the differentiation and migration of lymphocytes as result negative influence of the prolonged light on central immune organs.

**Conclusion:** These changes in organs of the immune system indicate both a premature aging of the spleen and the thymus and probably of all the immune system. Significant reduction in cell density in the immune organs associate with negative effects of the chronic light stress and leads to expressed immune dysfunction.

<http://dx.doi.org/10.1016/j.pbj.2017.07.124>

#### PS217

##### Intermittent low-level lead exposure causes anxiety and cardiorespiratory impairment



L. Shvachiy<sup>1,\*</sup>, V. Gerales<sup>1,2</sup>, Â. Amaro-Leal<sup>1</sup>, I. Rocha<sup>1,2</sup>

<sup>1</sup> Centro Cardiovascular da Universidade de Lisboa, Faculdade de Medicina, Universidade de Lisboa, Portugal

<sup>2</sup> Instituto de Fisiologia, Faculdade de Medicina, Universidade de Lisboa, Avenida Professor Egas Moniz, 1649-028 Lisboa, Portugal

E-mail address: [shvachiy.liana@gmail.com](mailto:shvachiy.liana@gmail.com)

(L. Shvachiy).

**Aim:** To characterize behavioural and cardiorespiratory changes in a new, intermittent low-level lead exposure animal model.

**Introduction:** Lead (Pb) is a cumulative toxic metal affecting all body systems that are particularly vulnerable during developmental phase. Permanent lead exposure has been defined as a cause of behavioural changes, cognitive impairment, sympathoexcitation, tachycardia, hypertension and autonomic dysfunction. However, no studies have been performed to describe a new, intermittent low-level lead exposure profile, that has been increased in the past years.

**Methods:** Foetuses were intermittently (PbI) exposed to water containing lead acetate (0.2%, w/v) throughout life until adulthood (28 weeks of age). A control group (without exposure, CTL), matching in age and sex was used. At 26 weeks, behavioural tests were performed for anxiety (Elevated Plus Maze Test) and locomotor activity (Open Field Test) assessment. Blood pressure (BP), electrocardiogram (ECG), heart rate (HR) and respiratory frequency (RF) rates were recorded at 28 weeks of age. Baroreflex gain (BRG) and chemoreflex sensitivity (ChS) were calculated. Student's *T*-test was used (significance  $p < 0.05$ ) for statistical analysis.

**Results:** An intermittent lead exposure causes hypertension (increased diastolic and mean BP), increased RF, decreased baroreflex function and increased ChS, without significant changes in HR, when compared to CTL group. Regarding behavioral changes, the intermittent lead exposure model showed an anxiety-like behaviour without changes in locomotor activity.

**Conclusion:** Intermittent low-level lead exposure induces changes on the cardiorespiratory profile characterized by hypertension, carotid chemosensitivity and baroreflex impairment. According to behavioural tests results, this study also shows that the exposure to lead during developmental phases causes anxiety in adult animals without locomotor activity impairment.

In summary, this study brings new insights on the environmental factors that influence nervous and cardiovascular systems during development, which can help creating public policy strategies to prevent and control the adverse effects of Pb toxicity.

<http://dx.doi.org/10.1016/j.pbj.2017.07.125>

## PS120

### Antihypertensive effects of two novel dihydropyridine derivatives

M. khoramjouy<sup>1,\*</sup>, A. Feizi<sup>2</sup>, M. Mahmoudian<sup>3</sup>, M. Faizi<sup>4</sup>

<sup>1</sup> Department of Pharmacology and Toxicology, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>2</sup> Pharmaceutical Sciences Branch, Islamic Azad University, Tehran, Iran

<sup>3</sup> Department of Pharmacology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

<sup>4</sup> Department of Pharmacology and Toxicology, school of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran

E-mail address: [khoramjouy.mona@gmail.com](mailto:khoramjouy.mona@gmail.com) (M. khoramjouy).

**Aim:** Treatment of hypertension.

**Introduction:** Mebudipine and dibudipine are two novel derivations of dihydropyridine (DHP) Ca<sup>2+</sup> channel blockers. Previous studies have shown that these two compounds have relaxant effects on vascular smooth muscles. In addition, DHPs are able to reduce contraction force of cardiac muscle in rat. In this study we decided to evaluate the antihypertensive effects of these two novel DHPs in hypertensive rat.

**Methods:** Male Sprague-Dawley rats were used in the study (8–10 weeks old). The rats were randomly divided to 4 groups of 10 rats (one control and 3 test groups). Blood pressure was measured by Tail cuff method. Left kidneys of the rats were removed by nephrectomy and sodium chloride 1% was added to the drinking water of animals and desoxycorticosterone acetate 20 mg/kg (SC) were injected twice a week. During and after 4 weeks, blood pressure of animals was evaluated to confirm the hypertension. Blood pressure of the animals was measured before i.p. injection of mebudipine and dibudipine (1–8 μmole/kg) and 2 min after the drug administration.

**Results:** Mebudipine and dibudipine significantly reduced the systolic blood pressure. Mebudipine was more effective than dibudipine and nifedipine in hypertensive animals and has significant results.

**Conclusion:** Previous studies showed that i.p. injection and oral usage of mebudipine and dibudipine decrease systolic hypertension in normotensive animals, on the other hand vasodilation effects of DHPs have been proved on aorta. Both novel drugs showed significant reduction in systolic blood pressure in hypertensive animals and mebudipine was more potent than dibudipine and nifedipine (as a standard drug uses). It is remarkable that, two new DHPs have similar efficacy and safety profile, but have higher efficacy compared to nifedipine in present study. The brilliant point is that DHPs as calcium channel blockers are more effective in hypertensive animals compared to normotensive animals.

<http://dx.doi.org/10.1016/j.pbj.2017.07.126>

## PS062

### Biological processes of polyphenols in the cardiovascular system: A bioinformatics approach

Augusto Rachão<sup>1,\*</sup>, Ana Filipa Silva<sup>2,3,a</sup>, Rita Nogueira-Ferreira<sup>2</sup>, Fábio Trindade<sup>2,3,4</sup>, Rui Vitorino<sup>2,3,4</sup>, Adelino Leite-Moreira<sup>2,3</sup>, Daniel Moreira-Gonçalves<sup>3,5</sup>, Tiago Henriques-Coelho<sup>2,6</sup>, Rita Negrão<sup>1,7</sup>

<sup>1</sup> Departamento de Biomedicina – Unidade de Bioquímica, Faculdade de Medicina da Universidade do Porto, Portugal

<sup>2</sup> Unidade de Investigação Cardiovascular, Faculdade de Medicina da Universidade do Porto, Portugal

<sup>3</sup> Departamento de Cirurgia e Fisiologia, Faculdade de Medicina da Universidade do Porto, Portugal

<sup>4</sup> Instituto de Biomedicina – iBiMED, Departamento de Ciências Médicas, Universidade de Aveiro, Portugal

<sup>5</sup> Centro de Investigação em Atividade Física, Saúde e Lazer, Faculdade de Desporto da Universidade do Porto, Portugal

<sup>6</sup> Departamento de Ginecologia, Obstetrícia e Pediatria, Faculdade de Medicina da Universidade do Porto, Portugal

<sup>7</sup> I3S-Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Portugal

E-mail address: [augusto-rachao\\_3@hotmail.com](mailto:augusto-rachao_3@hotmail.com) (A. Rachão).

**Aim:** In this study, we aimed to evaluate the cardiovascular system-related biological processes (BP) modulated by polyphenols in rodents and humans, and to verify which of them are specie-

<sup>a</sup> Augusto Rachão and Ana Filipa Silva contributed equally to this work.



specific, in order to understand which outcomes for cardiovascular diseases (CVD) could be translated from animal to human studies.

**Introduction:** CVD stand as a great cause of morbi-mortality worldwide and polyphenol-rich diets have been associated with improved cardiovascular risk profiles. Although rodent models have been a resourceful means of understanding the CVD mechanisms and possible outcomes of the use of polyphenols in that context, most experimental models do not fully reproduce human CVD.

**Methods:** Database searching was carried out on PubMed and Google Scholar using specific keywords concerning CVD, retrieving close to 300 publications. After excluding irrelevant results, proteome data was organized in Excel® spreadsheets and the Cytoscape platform, ClueGo + CluePedia and Venny 2.1.0 were used to explore the biological processes influenced by flavonoids in the approached CVD.

**Results:** This study was mainly focused in the species *Rattus norvegicus* and *Homo sapiens* and in flavonoids, a polyphenol subgroup. Only about 5% of the BP influenced by flavonoids were common to both species and they were mostly related to the maintenance of blood pressure and the fatty acid metabolic process. Nevertheless, these effects were accomplished through different proteins/pathways and different subgroups of flavonoids.

**Conclusion:** Our research highlights the need for a careful translation of the flavonoids' effects observed in rat models to clinical trials, since different proteins and subgroups of flavonoids mediated the observed actions. Though this type of studies can provide insights to help choosing the most adequate polyphenols as preventive approaches or therapies for human CVD, further investigation should be performed to clarify the described effects. Besides, pharmacokinetic aspects of the flavonoids' action should also be considered when planning clinical trials.

**Acknowledgements:** This work was supported by Portuguese Foundation for Science and Technology grants PEst-OE/SAU/UI0038/2014; UID/BIM/04293/2013, UID/IC/00051/2013 (financed by Fundo Europeu do Desenvolvimento Regional through COMPETE 2020 – Programa Operacional Competitividade e Internacionalização) and The European Foundation for Alcohol Research (ERAB) (EA 14 23).

<http://dx.doi.org/10.1016/j.pbj.2017.07.127>

## PS167

### Affinity of *Listeria sp.* proteins to cAMP and role in virulence

M. Fidalgo<sup>1,2,3,\*</sup>, J. Moscoso<sup>2,3</sup>, S. Sousa<sup>2,3</sup>, D. Cabanes<sup>2,3</sup>

<sup>1</sup> Universidade de Trás-os-Montes e Alto Douro (UTAD), Portugal

<sup>2</sup> Instituto de Biologia Molecular e Celular (IBMC), Portugal

<sup>3</sup> Instituto de Investigação e Inovação em Saúde (I3S), Portugal

E-mail address: [martafilipafidalgo@gmail.com](mailto:martafilipafidalgo@gmail.com) (M. Fidalgo).

**Aim:** The aim of this study was thus to identify Lm proteins capable to bind cAMP.

**Introduction:** Infectious diseases are still a major cause of death worldwide. To infect a host and survive the environment, bacteria have to sense their surrounding and adjust their behaviour. In this adaptation process, cAMP (cyclic adenosine monophosphate) is known to be an important player in pathogens such as *Pseudomonas spp.*, *Vibrio spp.* or *Mycobacterium spp.* The small molecule

cAMP is a cyclic nucleotide that relays information from receptors to one or more effector proteins within a bacterial cell, functioning as a second messenger. To mediate a response, cAMP allosterically interacts with cAMP-binding proteins. Understanding how this happens is fundamental to predict how bacteria will adapt/act to/in a given context.

**Methods:** We recently showed that the human foodborne pathogen *Listeria monocytogenes* (Lm) produces cAMP. The aim of this study was thus to identify Lm proteins capable to bind cAMP. To do this, four candidate proteins selected by bioinformatics analyses were expressed, purified and studied biochemically. Three approaches were used: cAMP affinity chromatography; competitive cAMP affinity chromatography; and isothermal titration calorimetry (ITC).

**Results:** Among the four tested proteins, CbpA displayed cAMP-binding ability on the three approaches used.

**Conclusion:** Hence, our preliminary results showed that CbpA binds to cAMP. It is now mandatory to understand the relation between cAMP and CbpA, to determine the function of the protein itself and in complex with cAMP, and to understand the importance of this signalling system for virulence.

<http://dx.doi.org/10.1016/j.pbj.2017.07.128>

Public Health & Medical Informatics Poster Session  
Saturday, September 16th, 10h00

## PS044

### Assessment of ECG interpretation skills among Polish medical students, nursing, emergency medicine and English Division medical students



Marek Stopa\*, Harison Sevenathan,  
Mateusz Bogusławski, Izabela Pałasz

Students Scientific Group of Interventional  
Cardiology at the 2nd Department of Cardiology and  
Cardiovascular Interventions

E-mail address: [mrk.stopa@gmail.com](mailto:mrk.stopa@gmail.com)

(M. Stopa).

**Aim:** The aim of the study was to evaluate ECG interpretation skills among study population, and analyze factors determining their score.

**Introduction:** The electrocardiogram examination is one of the most frequently performed diagnostic test. Correct interpretation of the ECG, particularly in life-threatening scenarios (LTS) may influence the decisions on appropriate actions and consequently have an impact on the lives and health of patients. It is important for medical, nursing and emergency medicine students to acquire this skill.

**Methods:** ECG interpretation skills were assessed by self-prepared questionnaire including questions about demographic data and 20 ECG problems with 17 cases. In 6 cases there were LTS. Three questions evaluated basic knowledge about rhythm, heart rate and axis. The survey was conducted via Internet. Study population consist of 551 medical, nursing and emergency medicine students.

**Results:** The overall score among Polish medical students is 46% which is higher than nursing and emergency medicine students (22% and 37% respectively;  $p < 0.001$  in both). English division students scored almost similarly (49%;  $p = 0.27$ ). Polish medical students scored better in LTS than the nursing students (37% vs 23%;  $p < 0.001$ ). Among Polish medical students: Students in year “4–6” scored higher than those from year “1–3” (overall score: 51% vs. 31%;  $p < 0.001$ , LTS: 41% vs 25%;  $p < 0,001$ ). In addition, members



of cardiology scientific groups scored higher than the rest (57% vs 43%;  $p < 0.001$ ).

**Conclusion:** There is low level of ECG interpretation among medical students and quality of ECG training should be improved. Various factors influences ECG interpretation knowledge among students.

<http://dx.doi.org/10.1016/j.pbj.2017.07.129>

#### PS126

##### Assessment of safe injection practice among nurses in Port Said General Hospital



Karim Farag<sup>1,\*</sup>, Ahmed El-kiki<sup>1</sup>, Ahmed Emam<sup>1</sup>, Ahmed Mourad<sup>1</sup>, Alaa Abdelrahman<sup>1</sup>, Amira Fekry<sup>1</sup>, Asmaa Eita<sup>1</sup>, Asmaa Galal<sup>1</sup>, Asmaa Ghanem<sup>1</sup>, Eslam El-shourbagy<sup>1,2</sup>, Esraa Hawas<sup>1,2</sup>, Nermeen Gmal<sup>1</sup>, Rawan Ghaly<sup>1</sup>, Sara El-salou<sup>1</sup>, Ayat Tawfik<sup>2</sup>, Ei<sup>2</sup>

<sup>1</sup> Students at Faculty of Medicine Port Said University, Egypt

<sup>2</sup> Department of of Public Health and Preventive Medicine Faculty of Medicine Port Said University, Egypt

E-mail address: [king\\_kemoo.2010@yahoo.com](mailto:king_kemoo.2010@yahoo.com) (K. Farag).

**Aim:** Improving safe injection practice in Port-Said General Hospital.

**Introduction:** A safe injection is one that, “does not harm the recipient, does not expose the provider to any avoidable risk and does not result in waste that is dangerous for the community”. In developing countries, about 16 billion injections are administered each year.

**Methods:** Through-out March 2015, a cross-sectional, descriptive study was conducted to assess safe injection practice among 150 nurses in Port-Said General Hospital. Data collected by observational CDC Checklist and another checklist for unit evaluation.

**Results:** Regarding needle disposal 77% of nurses got rid of the needle in safety box, 1% threw it in the pin while 22% threw it in a barrel. Regarding hand washing 41% of nurses washed their hands before preparing medication, while 23% of nurses washed their hands before touching patients and 51% of them washed their hands after touching patients. 57% of nurses wore gloves while 43% didn't. We found 52% of nurses didn't have HBV vaccine. We found also 77% of nurses were trained on safe injection while 23% weren't trained.

Regarding observation, 73% of medication areas were cleaned while 27% weren't. 83% of nurses used single dose vials, ampoules or bottles of intravenous solution for only one patient while 17% didn't. Regarding to hospital unites, only 44% of unites had written policies or procedures for safe injection.

**Conclusion:** Our evaluation results are good regarding clean medication area, needles for one patient, new needles and syringes, using single dose Vial and using medical connectors for one patient, while are poor regarding disinfecting rubber septum of vial, dating multi dose vials for 28 when opened and keeping multi dose vial in a centralized medication area and not to enter it in the immediate patient area, these poor results may be due to some untrained nurses.

<http://dx.doi.org/10.1016/j.pbj.2017.07.130>

#### PS199

##### Frequency of analgesic drugs use and patients' awareness of their possible interactions with antiplatelet therapy in coronary heart disease



I. Pałasz\*, Ł. Reczek, M. Schonborn, S. Janiec, M. Cebenko

Students' Scientific Group at the 1st Department of Cardiology, Interventional Electrophysiology and Hypertension, The Netherlands

E-mail address: [isanula@gmail.com](mailto:isanula@gmail.com)

(I. Pałasz).

**Aim:** The aim of the study was to assess the prevalence and frequency of analgesic drug use in patients with coronary heart disease, as well as to assess patients knowledge about possible interactions of these drugs with conventional cardiac therapy.

**Introduction:** Nonsteroidal antiinflammatory drugs (NSAIDs) are commonly used in the management of pain in a variety of conditions. Available data clearly indicate that the NSAIDs use is associated with a number of adverse effects especially in patients with cardiovascular disease.

**Methods:** The study group consists 93 patients hospitalized in the tertiary cardiology center (mean age  $67 \pm 11$  years, 30% females). Thirty nine subjects were hospitalized with diagnosis of acute coronary syndrome and 54 underwent elective coronary intervention. Self-prepared questionnaire was used to collect data.

**Results:** In the examined group 56 patients (60%) declared the use of analgesic drugs with regular use (defined as at least 3 times per week) reported by 25 subjects (27%). The most frequently used analgesics were NSAIDs ( $n = 37$ ), paracetamol ( $n = 36$ ), less commonly patients reported the use of tramadol ( $n = 6$ ) or metamizol ( $n = 9$ ). The majority of patients using analgesic are not aware about possible interactions with antiplatelet therapy (the answer “yes” for the question about knowledge of possible interactions of analgesic with cardiac treatment gave only 21% of responders, while majority answered “do not know” (72%)). Only 20% of patients admitted that they received the information about analgesics from their doctor. Majority of patients do not consult the use of analgesics with the physician (72%).

**Conclusion:** The regular use of NSAIDs/analgesic by 27% of hospitalized patients with coronary heart disease is a significant concern. Patients with coronary heart disease should be provided with detailed information and recommendation about safe analgesic therapy and alternatives for NSAIDs.

**Acknowledgements:** The authors thank Prof. Danuta Czarnecka, head of the 1st Department of Cardiology, Interventional Electrophysiology and Hypertension Collegium Medicum Jagiellonian University and Agnieszka Olszanecka Ph.D. for substantive supervision over the study.

<http://dx.doi.org/10.1016/j.pbj.2017.07.131>

#### PS094

##### A new route for Medical Education: Rethinking Anatomy's learning strategies



S. Tsisar<sup>1,\*</sup>, J.M. Diniz<sup>1</sup>, B. Viana<sup>1</sup>, M. Sousa<sup>1</sup>, B. Afonso<sup>1</sup>, R. Santos<sup>1</sup>, J.F. Silva<sup>1</sup>, B. Guimarães<sup>1,2</sup>

<sup>1</sup> Department of Public Health, Forensic Sciences and Medical Education. Unit of Medical Education and Simulation. Faculty of Medicine, University of Porto, Porto, Portugal

<sup>2</sup> Center for Research in Health Technologies and Information Systems. Faculty of Medicine. University of Porto, Porto, Portugal  
E-mail address: [stastv94@gmail.com](mailto:stastv94@gmail.com)  
(S. Tsisar).

**Aim:** The aim of this work is to analyze the advantages of the introduction and diversification of pedagogical strategies in Anatomy Education, as a comprehensive model of Medical Education.

**Introduction:** Medical Education has suffered a paradigmatic shift that led to curricular reforms. Due to scientific and technological development, Medical curriculum has been adopting a vertical integration model, in which basic and clinical sciences coexist during medical instruction. This context favours the introduction of new complementary technology-based pedagogical approaches. Thus, even traditional core fields of medical curriculum, like Anatomy, are refocusing their teaching/learning standards.

**Methods:** This work presents the main conclusions of a bibliographic review that reflected on Medical Education's current pedagogical trend, by analyzing the advantages of the introduction and diversification of pedagogical approaches in Anatomy Education.

**Results:** Anatomy Education's status quo is characterized by less available teaching time, increasing demands of 2D perspective of human anatomy from radiology and endoscopy imaging and other invasive and non-invasive medical techniques, increasing number of medical students and other logistical restraints. The traditional learning approach, mainly based in the cadaveric dissection, is drifting to complementary newer technologies as 3D models or 2D/3D digital imaging to examine the human anatomy. Also, knowledge transference is taking different channels, as learning management systems, social networks and computer-assisted learning and assessment are assuming relevant roles.

**Conclusion:** The future holds promising approaches for education models. Artificial Intelligence, Virtual Reality and Learning Analytics may provide analytic tools towards a real-time and personalized learning process.

A reflection on Anatomy Education, as a comprehensive model, allows us to understand Medical Education's complexity. Therefore, the present Medical Education context favours a blended learning approach, based on multi-modality pedagogical strategies.

<http://dx.doi.org/10.1016/j.pbj.2017.07.132>

#### PS174

### Human *Dirofilaria* (Nochtiella) *repens* infection in Serbia



J. Savkić<sup>1,\*</sup>, A. Džamić<sup>2</sup>

<sup>1</sup> School of Medicine, University of Belgrade, Belgrade, Serbia

<sup>2</sup> Department of Parasitology-Mycology, Institute of Microbiology and Immunology, School of Medicine, University of Belgrade, Belgrade, Serbia  
E-mail address: [savkic.jana@gmail.com](mailto:savkic.jana@gmail.com)  
(J. Savkić).

**Aim:** The aim of this study was to present and describe cases of *D. repens* infection in Serbia from 2013 to 2016. The cases were reported in National Reference Laboratory for Parasitic Zoonoses in Belgrade.

**Introduction:** Dirofilariasis is a vector-borne parasitic zoonosis caused by *Dirofilaria* genus which uses female mosquitoes as vectors. One of the important species is *Dirofilaria repens*. Dogs are main reservoir hosts for *D. repens*, whilst humans can be accidental

hosts. The most common site of infection is ocular region. Human dirofilariasis is a rare infection, however, the number of reported cases is increasing in Europe and Serbia.

**Methods:** Retrospective study was performed including data from National Reference Laboratory for Parasitic Zoonoses in Belgrade. The results were processed and selected parameters were described: sex and age of the patients, locality, location of the lesion, clinical signs and characteristics of parasites. Diagnostic and therapeutic procedures, previous diagnosis were also described.

**Results:** Seven cases were reported during selected period. The mean age of patients was 44.9 years. Six patients were females (85.7%) and five patients were from Belgrade (71.4%). The most frequent site of infection was ocular region in four patients – 57.5%, while the other sites were skin of abdomen and limbs. All patients had nonspecific clinical signs. The most common cause of the infection was immature female worm in five cases (71.4%).

**Conclusion:** The number of cases of human *D. repens* infection is increasing in Serbia, and the most common site of the infection is ocular region. Considering the fact that cases are mainly misdiagnosed, it is important to point out the significance of this infection in differential diagnosis of different diseases.<sup>1–4</sup>

#### References

- Genchi C, Mortarino M, Rinaldi L, Cringoli G, Traldi G, Genchi M. Changing climate and changing vector-borne disease distribution: the example of *Dirofilaria* in Europe. *Vet Parasitol.* 2011;176:295–9.
- Simon F, Siles-Lucas M, Morchon R, et al. Human and animal dirofilariasis: the emergence of a zoonotic mosaic. *Clin Microbiol Rev.* 2012;25:507–44.
- Džamić AM, Čolović IV, Arsić-Arsenijević VS, et al. Human *Dirofilaria repens* infection in Serbia. *J Helminthol.* 2009;83:129–37.
- Tasić-Otašević SA, Trenkić Božinović MS, Gabrielli SV, Genchi C. Canine and human *Dirofilaria* infections in the Balkan Peninsula. *Vet Parasitol.* 2015;209: 151–6.

<http://dx.doi.org/10.1016/j.pbj.2017.07.133>

#### PS104

### The multidimensional approach to suicide done through self-mutilation with an overview of wounds



Ciuk Katarzyna\*, Ciuk Szymon, Dadański Emil, Chukwu Ositadima, Bociąga Marta, Burghardt Wiktoria

Jagiellonian University Medical College, Poland  
E-mail address: [kataryna962@gmail.com](mailto:kataryna962@gmail.com)  
(C. Katarzyna).

**Aim:** The aim of the study was to evaluate the methods and wounds of suicide done by self-injury.

**Introduction:** Hanging and drug overdose are the most common ways of suicide. However, there are also more painful methods of dying. This study considers: stabbing, cutting with a knife, ingestion of sharp foreign body, self-shooting, self-arson, crushing.

**Methods:** There were 65 recorded cases ( $M=56$ ,  $F=9$ , mean age:  $49.96 \pm 15.78$ ) of self-mutilation as a way of death in archives of the Department of Forensic Medicine of Jagiellonian University Medical College in Cracow in years 2011–2016. All of them were studied in terms of the method, trial and mortal wounds (number, area, type), condition of clothing, prior psychiatric treatment, prior suicide attempts. All calculations were done with the usage of Statistica software.

**Results:** The most common methods of suicide were self-shooting (38.46%), cutting (26.15%), stabbing (16.92%). There was 1 case of foreign body ingestion and 1 of head crushing in a blacksmith machine. There were 6 cases of self-arson. Trial wounds were observed in 29.23% cases, all of them were recorded in

cases of either stabbing or cutting. The places of mortal wounds: 43.4% head, 20.8% thorax, 18.9% upper limb, 9.4% neck, 3.8% lower limb, 1.9% abdomen, 1.9% digestive tract. The majority of patients (72.3%) had no previous mental treatment and prior suicide attempt (84.61%). In 57/65 cases the place of the wound was exposed.

**Conclusion:** People in their fifties commit suicide with self-injury. It happens most often with either a gun or a knife. The trial wounds were observed in cases of stabbing or cutting. 4 of 6 cases of self-arsony were accompanied by previous psychiatric treatment. Females commit suicides through self-mutilation more rarely than males.

**Acknowledgements:** Jagiellonian University Medical College.

<http://dx.doi.org/10.1016/j.pbj.2017.07.134>

## PS160

### Intestinal colonization of residents of long-term care facilities and nursing homes in Braga area with Multidrug-resistant Gram-negatives



G. Duarte<sup>1,\*</sup>, R. Mota<sup>1</sup>, D. Gonçalves<sup>1,2,3</sup>,  
H. Ferreira<sup>1,2</sup>

<sup>1</sup> Microbiology, Department of Biological Sciences,  
Faculty of Pharmacy, University of Porto, Portugal

<sup>2</sup> 2UCIBIO, University of Porto, Portugal

<sup>3</sup> Superior Institute of Health of Alto Ave, Portugal

E-mail address:

[gracinda.duarte.bluestar@hotmail.com](mailto:gracinda.duarte.bluestar@hotmail.com)

(G. Duarte).

**Aim:** The aim of our work was the detection of Enterobacteriaceae isolates producing extended-spectrum beta lactamases (ESBL) and isolates with reduced susceptibility to carbapenems, in the intestinal flora of institutionalized-residents in extra-hospital-health-care facilities in Braga region.

**Introduction:** Care of aging population has been a growing challenge to public-health and health-care providers. Due to the disabilities of older people, there is a growing need for long-term care facilities (LTCF) and nursing homes (NH). This brings a new paradigm for the spread of bacteria showing multidrug-resistance (MDR) to antibiotics.

**Methods:** Fecal samples of 27 residents of these institutions were collected (September-to-December, 2016). One gram of each sample was suspended in 10 mL of saline and 100L of the suspension was spread on MacConkey agar with ampicillin(100 mg/L)/cefotaxime(2 mg/L)/meropenem(1 mg/L). Susceptibility to antibiotics was determined by disk-diffusion methods, according to CLSI. ESBL-producers were detected by the double-disk-synergy-test and/or clavulanic-acid addition and PCR was performed for detection of blaTEM, blaOXA, blaSHV, blaCTX-M-group-1, blaCTX-M-group-2, blaCTX-M-group-8, blaCTX-M-group-9, blaCTX-M-group-25, tetA, tetB, aac(3)-II, sul1, aac(6)-Ib and qnrB genes.

**Results:** The study revealed 6 ESBL-producing Enterobacteriaceae colonizing 2 residents in LTCF (2-Escherichia coli/1-Klebsiella, Enterobacter, Serratia and Citrobacter (KESCgroup)) and 3 residents in NH (2-Escherichia coli/1-KESCgroup). Isolates showed positive for blaCTX-M-group-1, blaCTX-M-group-9, blaTEM, blaSHV, blaOXA, tetA, tetB, aac(3)-II, sul1 and aac(6)-Ib. These isolates showed resistance to non-beta-lactam antibiotics, namely to tetracycline, ciprofloxacin, trimethoprim-sulfamethoxazole, gentamicin and amikacin. We detected 6 MDR-bacteria isolates and 1 isolate with reduced susceptibility to carbapenems.

**Conclusion:** Our results show the dissemination of ESBL-producing-Enterobacteriaceae in intestinal colonization of LTCF/NH patients, who may act as vehicles of MDR-bacteria within the health-care-facilities and community.

<http://dx.doi.org/10.1016/j.pbj.2017.07.135>

## PS202

### Social desirability in medical school admission: Differences between students from regular contingent and graduate admission



Ana Catarina Silva\*, Ana Filipa Pinto, Joana Abreu,  
Juliana Couras, Sofia Calçada

Faculty of Medicine of University of Porto, Portugal

E-mail address: [catarina.silva12@live.com.pt](mailto:catarina.silva12@live.com.pt)

(A.C. Silva).

**Aim:** Our work targeted the degree of social desirable answers among two different contingents with the objective of providing useful statistically insight about how this variable may be responsible for the differences among both groups. With the new evidence found in this study we hope to provide useful insight to help improving the selection process of the applicants.

**Introduction:** The relation between social desirability and medical school applicants may be of interest when analyzing the results of questionnaires in medical school admission. Our study analyzes the difference in social desirability between graduates admitted to a medical school, and students enrolling in the school by the regular admission process.

**Methods:** We used a resumed version of the Marlow-Crowne Desirability Scale to compare the social desirability between students from the regular and graduate admission. After collecting 181 questionnaires between 2005 and 2006, all the data was analyzed using R software. The sample was described by performing *t*-test between regular and graduate admission in the following variables: gender, marital status, childbearing, working status, previous residence and age.

**Results:** Statistically significant differences were found in marital and working status as well as within. The main outcome – social desirability – was also statistically different among both groups, indicating that students with graduate admission had given more desirable answers. A analysis of subgroups according to marital and working status was made, being the results also significant.

**Conclusion:** Our study suggests that graduate admission students have more social desirability than regular students. However, more research is needed in order to find in what way other factors, as age or working status, for example, might have influenced these results.

**Acknowledgements:** Milton Severo.

<http://dx.doi.org/10.1016/j.pbj.2017.07.136>

## PS105

### Headache among medical students in Bukovina Region of Ukraine



I. Yaremchuk\*, O. Yaremchuk

Department of Nervous Diseases, Psychiatry and  
Medical Psychology

E-mail address: [yaremchuk.cv@gmail.com](mailto:yaremchuk.cv@gmail.com)

(I. Yaremchuk).

**Aim:** To study the prevalence of headache among medical students in Bukovina region of Ukraine.

**Introduction:** Headaches are the most prevalent neurological disorder and among the most frequent symptoms among medical students. Headache disorder is a major public health issue and is a great burden for a person, health care system, and a society. Identifying of headache risk factors is necessary for treatment and effective prevention.

**Methods:** A cross sectional study has been conducted in duration from 30th of January to the end of March 2017. 146 students of Bukovinian State Medical University aged 19–26 years were interviewed by using specially designed questionnaires. All students with a headache who completed questionnaires were examined by neurologist. The type of headache was determined according to the diagnostic criteria of the classification of the International Headache Society, 2003.

**Results:** Our study has found that headaches bother 121 (82.8%) students. Among respondents periodic headache was observed in 71.9% (105 students). 16 (10.1%) students experienced chronic headaches that bothered them more than 15 days a month. Among girls the prevalence of headache was significantly higher than among men – 62.1% and 38.9% respectively. Among the students the most frequent headache (58.7%) was tension headache, 9.1% of respondents had migraine headache, 32.2% had other types of headaches. According to the students' responses the most prevalent causes of headache were stress – 36%, sleep disturbance – 20%, and weather changes – 14%. The other causes included skipping meals – 11%, tea or coffee overdrink – 8%, alcohol consumption – 6%, and menstrual cycle disturbances in 2% students.

**Conclusion:** By means of the research there was found a high prevalence of headaches among medical students. There was determined the prevalence of primary cephalgias, namely, migraine without aura and episodic tension-type headaches. There was shown a lack of awareness of students about the causes of headache reflected in inadequate symptomatic treatment in most cases.

<http://dx.doi.org/10.1016/j.pbj.2017.07.137>

## PS194

### New health problems: Assessment of nutritional and metabolic profile from indigenous citizens in the reserve park of Xingu



André Alencastro Curado Filho<sup>1,\*</sup>,  
Patricia Gardiman Arruda<sup>1</sup>,  
Pedro Ernesto Carvalho de Cillo<sup>1</sup>,  
Douglas Rodrigues<sup>2</sup>

<sup>1</sup> Medicine undergraduation student of Escola Paulista de Medicina (EPM), Universidade Federal de São Paulo (UNIFESP)

<sup>2</sup> MD, PhD in Epidemiology - Researcher in the Area of Indigenous Health and Tradicional Communities  
E-mail address: [andrealecastro79@gmail.com](mailto:andrealecastro79@gmail.com)  
(A.A.C. Filho).

**Aim:** To identify the prevalence of metabolic syndrome, dyslipidemia, systemic arterial hypertension and type 2 diabetes mellitus in the adult population.

**Introduction:** In the last 20 years, there have been many changes in the way of life of the Indians of the Xingu Indigenous Park (PIX), resulting from their contact with our society. Factors such as the increasing consumption of processed foods in substitution of the traditional diet based on natural products, the reduction of physical activity by the incorporation of technology and the monetarization of the economy have produced changes in the epidemiological profile of these populations. Previously non-existent diseases like type 2 diabetes mellitus, systemic arterial

hypertension, obesity and dyslipidemias have been increasingly observed. In this work, preliminary results of data collected between February and March 2017 are presented.

**Methods:** Individuals over 18 years old of both sexes undergoing physical examination (clinical and anthropometric), bioimpedance test to evaluate the percentage of body fat and blood collection by digital puncture for lipid profile and fasting glucose of 8–12 h.

**Results:** The metabolic profile of 188 indigenous people of the Pavuru polo showed: High index of overweight and dyslipidemia in both populations; high central obesity in females; Increased number of hypertensive and fasting blood glucose altered among men; 25% have intermediate or high cardiovascular risk in the next 10 years among men, almost twice as many women.

**Conclusion:** Based on the data obtained, we can conclude that the process of socio-cultural transition to which the residents of the 4 villages studied are inserted has a direct impact on their health, making it imperative to indetificate and delineate the magnitude of the problem, seeking to develop strategies for the primary and secondary prevention of diseases related to nutritional and metabolic alterations and their consequences among the people living in the PIX.<sup>1–5</sup>

## References

- [WHO] World Health Organization. Physical Status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. Geneva: WHO; 1995. 457 p. [Technical Report Series n. 854].
- 7a Diretrizes Brasileiras de Hipertensão. Sociedade Brasileira de Cardiologia. Arq Bras Cardiol. 2016;107 Suppl. 3:11–2.
- Simão AF, Prêcoma DB, Andrade JP, Correa Filho H, Saraiva JFK, Oliveira GMM, et al. Sociedade Brasileira de Cardiologia. I Diretriz Brasileira de Prevenção Cardiovascular. Arq Bras Cardiol. 2013;101 Suppl. 2:1–63.
- Diretriz. Brasileira sobre Dislipidemias e Prevenção da Aterosclerose, V – Departamento de Aterosclerose da Sociedade Brasileira de Cardiologia. Arq Bras Cardiol. 2013;88(S1):2–19.
- Diretrizes da Sociedade Brasileira de Diabetes (2015–2016), Milech A. Organização José Egídio Paulo de Oliveira. Sérgio Vencio – São Paulo: A.C. Farmacêutica; 2016.

<http://dx.doi.org/10.1016/j.pbj.2017.07.138>

## PS149

### Late diagnostics of Alzheimer's disease and other dementias, retrospective research in Vilnius (Lithuania)



Roberta Vaikutytė\*, Lina Lekečinskaitė

Vilnius University, Faculty of Medicine, Lithuania

E-mail address: [roberta.vaikutyte@gmail.com](mailto:roberta.vaikutyte@gmail.com)

(R. Vaikutytė).

**Aim:** To gain a deeper understanding around the prolonged diagnosis time for Alzheimer's disease.

**Introduction:** The world population is getting older. Correspondingly the number of old age sicknesses (like Alzheimer's disease (AD), dementias) is rising in the populations. These patients require custody, medicine and that takes a lot of money and resources from the country.

**Methods:** There were two multiple choice/short answer surveys done. One for doctors (psychiatrists, neurologists, general practitioners) and the other for the patients care givers. The surveys were administered in the hospitals in Vilnius. The data were evaluated by SPSS program ( $\alpha=0.05$ ).

**Results:** 50 doctors completed the surveys. Doctors emphasize that just about 14% of patients' care givers are aware of the possibility for the genetic screening for AD and are interested in pursuing it. Moreover, according to the surveys, patients with cognitive disorders seek help when they are about 65–75 years old and 70% of

the doctors believe that this is too late. 68% of the doctors agree that the society does not have enough knowledge about dementias.

50 patients' caregivers completed the surveys: 26% – AD, 28% – vascular dementia, 46% – non-defined. The first contact with their doctors varied greatly among different dementia patients: AD – waited for 2.8 y, vascular dementia – 1.46y, non-defined – 0.87 y. Even though patients' caregivers indicated that they received enough information from their doctor, they admitted that they had no or not enough knowledge concerning the disease before. They also highlighted that it was hard to find information in Lithuanian language.

**Conclusion:** The research showed that people do not have enough knowledge about dementias and have limited access to information. This is one of the main reasons why the diagnostics is late. Considering the mental health is a stigmatic topic in the world, the fact that people do not know about the dementias might be the problem in the whole world.<sup>1</sup>

## Reference

1. <https://www.nia.nih.gov/newsroom/2016/03/worlds-older-population-grows-dramatically>

<http://dx.doi.org/10.1016/j.pbj.2017.07.139>

## PS010

### Obsessive-compulsive behaviour tendencies among medical students in Poland



Zuzanna Goetz\*, Ewa Żelnic

Medical University of Warsaw, Outpatient Psychiatric Association for medical students at the NZOZ Centrum Terapii DIALOG, Poland  
E-mail address: [zuzannagoetz@gmail.com](mailto:zuzannagoetz@gmail.com)  
(Z. Goetz).

**Aim:** Determination of a need for educational program, focused at students' knowledge and awareness of OCD.

**Introduction:** As medical students, we are familiar with terms such as professional burnout and workaholism on an every day basis. However other psychiatric disorders seem to be still stigmatized and not well perceived for this profession. It does not mean that they do not exist among future physicians.

Gabbard and Mayers believed that perfectionism is one of the most common personality traits for medics. Moreover, they assumed that perfectionistic physician might perform more often "obsessive triad". Namely: self-doubt, guilt feelings, exaggerated sense of responsibility. Those personal features can lead to obsessive-compulsive behaviour. Authors also suggest that those traits are already widely pre-existing for medical students. However no studies supporting this thesis were conducted.

**Methods:** Anonymous self-completion questionnaire, completed by students from different faculties who study at the Polish universities. Questions based on structuralized clinical questioner designed by the Polish psychiatrists (dr Bryńska and Wolańczyk, 2005).

**Results:** The questionnaire was completed by 855 students out of whom: 393 medical students (46%). 53% (454) students declared performing specific acts, although not purposely or despite their will (e.g. checking if the door are locked, cleaning hands, counting. Only 49.6% (194) of medical students answered positively, in respect to 56% (259) non-medical students. Persistent thoughts and fantasies, which are not wanted but occur very often and persistently return – admitted 53.5% (457) questioned. Among the future doctors only 43.5% (170) confirmed to have those thoughts. In contrary to 63% (287) non-medical students.

**Conclusion:** At first glance it seems that medical students less frequently admit to perform compulsive behaviour or to have obsessive thoughts. But what is the cause? Are they less prone to those behaviours? Or, as Gabbard and Mayers suggest, they have to be seen as "perfect" in front of themselves or others? To answer those questions further research is needed.

<http://dx.doi.org/10.1016/j.pbj.2017.07.140>

## PS132

### Assessing the prevalence of HBV and HCV infections in children under going hemodialysis and the related risk factors in a children's Medical Center



Seyyed Mostafa Ahmadi, Neda Raeessi\*

Tehran University of Medical Sciences, Research Center, Tehran, Iran

E-mail address: [Raeessi.f@gmail.com](mailto:Raeessi.f@gmail.com)

(N. Raeessi).

**Aim:** Assessing the prevalence of HBV and HCV infections in children under going hemodialysis.

**Introduction:** Chronic hemodialysis is a life saving process in patient with end stage renal disease. Hemodialysis patients are at high risk for viral hepatitis infections due to the high number of blood transfusion sessions, prolonged vascular access and the potential for exposure to infected patients and contaminated equipments. Approximately 8% and 20% of hemodialysis patients have B hepatitis and C hepatitis respectively and this data varies from country to country. Hepatitis B virus (HBV) and hepatitis C virus(HCV) infections are important causes of morbidity and mortality in hemodialysis patients.

**Methods:** Based on the information gathered from the 149 hemodialysis children files, some special questionnaires were filled in. the obtained data was assessed and analyzed in SPSS software.

**Results:** A total of 149 hemodialysis patients with mean age 8.8(range: 0.24–16.74) years were enrolled in the study. Out of the total 149 patients, 74 were male and 75 were female. The majority of the patients (51 people) were in the 7–10 years age range. After glomeropathies (34 cases – 22.8%),reflux nephropathies(24 cases – 16.10%) were the main reasons in charge of renal impairment in our study population.

The results of our study in hemodialysis patients referring to the children's Medical Center of Iran from 1991 to 2009 suggests that prevalence of B and C hepatitis were both 2.04%and the prevalence of the concurrent infections (B and C hepatitis) were 2.72%.

**Conclusion:** This study confirms that the prevalence of B and C hepatitis among hemodialysis children referred to children's medical center are much lower than the adult hemodialysis patients in Iran and worldwide. This might indicate the higher health standards and the absence of intravenous drug abuse and unsafe sex among our study population. Screening donated bloods, treating anemia with erythropoietin, avoidance of dialyzer reuse, assignation of dedicated dialysis rooms, machines, and staff for infected patients, new disinfection methods, screening the patients before entering dialysis program and vaccination of susceptible patients and staff all have been referred to as means of limiting hepatitis transmission within our dialysis unit.

The results of this study can be used in health programming and budget allocating for this group.

<http://dx.doi.org/10.1016/j.pbj.2017.07.141>

## PS078

**Evaluation of differences in attitudes of service users about private and public health system of Serbia**

Stefan Jerotic, Emilija Ivancajic\*

*School of Medicine, University of Belgrade, Serbia**E-mail address: [emilijaivancajic@gmail.com](mailto:emilijaivancajic@gmail.com)*

(E. Ivancajic).

**Aim:** To evaluate attitudes of service users towards public and private health system of Serbia and how they influence users.**Introduction:** It has been noticed that there are differences in attitudes towards public and private health system in Serbia. This attitudes influence service users in the process of choosing their medical treatment.**Methods:** 400 users of the public and private health system (50% male and 50% female users) in Serbia filed up online questioner that consisted of 9 questions.**Results:** 49% of users had more trust in doctors in public sector and 51% of users had more trust in doctors in private sector. For users that had more trust in private health sector, a perfect doctor was a male (for 14% of users) under age of 35 (for 50.2% of users), if they had a health problem 54.8% of them would go to a doctor, 44% of this users thought that private health system in Serbia is too expensive, and 31.3 of them did not understand their diagnosis after the visit to the doctors. For respondents that had more trust in public health sector, a perfect doctor was a male (for 10% of users) under age of 35 (for 54.3% of users), if they had a health problem 41.3% of them would look it up on the internet, 32% of this users thought that private health system in Serbia is too expensive, and 31.2% of hem did not understand their diagnosis after the visit to the doctors.**Conclusion:** The biggest differences among users of the private and public health system are seen in the manner of what would be the best source of information for their health problems, other similarities can be explained by the fact that same doctor worked both in public and private health sectors.<http://dx.doi.org/10.1016/j.pbj.2017.07.142>

## PS106

**Non-consumable alcohol poisoning in the material of the Forensic Medicine Department in Cracow in 2007–2016**

K. Hapkiewicz\*, V. Bezshapkin, R. Jastrzębski, A. Koszman, M. Niemirowska

*Jagiellonian University Medical College, Poland**E-mail address: [kamhap96@gmail.com](mailto:kamhap96@gmail.com)*

(K. Hapkiewicz).

**Aim:** To analyse cases, find correlation and compare results with other Departments in the future.**Introduction:** Non-consumable alcohols – ethylene glycol, methanol, isopropanol and acetone – are the products of everyday use, though cases of lethal poisoning are not very frequent. They may be found in mixture with comestible ethanol.**Methods:** The subject of the study was 50 lethal cases (8 women), which had been caused by direct or indirect poisoning connected with consumption substances in the period 2007–2016 in the material of the Department of Forensic Medicine in Cracow.**Results:** The mean of cases for study period is 5,1 cases/year, with the notable rise of incidents in 2012 (16). Methanol poisonings were reported in 23 cases (46%), with the prevalence of men (78.3%). The blood concentration in both sexes ranged from 0.45 mg/l to

5340 mg/l. In period 24.10.2012–07.01.2013 the highest frequency of methanol poisoning was observed. It is linked to 2012 Czech methanol poisoning scandal, and influence of this event on Polish alcohol market. Ethylene glycol poisonings are reported in 9 cases (1 woman). The concentrations of alcohol in blood of males varied from 8 mg/l to 5710 mg/l; in female was 96.7 mg/l. The average age of victims was 45.6 years. Isopropanol and acetone poisonings were observed in 17 cases. The concentration of isopropanol ranged from 160 to 5589 mg/l. It is worth noticing, that acetone may partially be a product of metabolic transformation of isopropanol. The average age of victims is 45.6 years.

**Conclusion:** The vast majority of lethal non-consumable alcohol poisonings of victims were adult men with chronic overconsumption of alcohol. Nevertheless, some mean blood alcohol concentrations in women are comparable to those in men. There is an obvious time correlation between lethal alcohol poisonings and information about non-consumable alcohol appearance on the market.<http://dx.doi.org/10.1016/j.pbj.2017.07.143>

## PS026

**Scientific support of the most widespread and socially significant diseases in 2016**A. Zakrutko<sup>1,\*</sup>, L. Zakrutko<sup>2</sup><sup>1</sup> Bukovinian State Medical University, Ukraine<sup>2</sup> Ukrainian Centre of Scientific Medical Information

and Patent License Provision, Ukraine

*E-mail address: [lzakrutko@ukr.net](mailto:lzakrutko@ukr.net)*

(A. Zakrutko).

**Aim:** Analysis of scientific support of the most widespread and socially significant disease in 2016.**Introduction:** Scientific and technical activity is carried out by academic programs, scientific directions, scientific problems and research tasks, which are defined by perspective, practice and necessities.**Methods:** The regulatory and legal materials in healthcare of Ukraine are analyzed. The methods of systems analysis, statistical, structural and logical analysis are applied.**Results:** In 2016 year 177 scientific-research works were being done in 36 establishments (institutions) of Ministry of Health of Ukraine; including 85.4% applied researches and 14.6% fundamental ones. Among the total number of establishments 19 scientific and research institutions provided 83 research projects (46.9%), including 77 applied and 6 fundamental researches. In 17 medical educational establishments 94 research projects (53.1%) were being done, including 74 applied and 20 fundamental researches.

In 2016 year 69 scientific-research works were mainly focused on the development and improvement of existing methods of diagnostics, treatment and especially prevention of the most widespread and socially significant diseases, among them 27 works were devoted to researching of cardiovascular diseases, 24 – to cancer pathology, 3 – to tuberculosis, 4 – to HIV/AIDS, 4 – to diabetes, 7 were devoted to issues of scientific evidence of medical and rehabilitation provision of soldiers who took part in anti-terrorist operation, especially those, who were injured and became disabled.

**Conclusion:** Thus, the results of scientific researches of leading scientists will have real impact on the quality of medical care, improvement of indicators and health care, accelerate the reform of the industry through the scientifically based measures.<http://dx.doi.org/10.1016/j.pbj.2017.07.144>

## PS228

**Is more stressful to become a physician or a pharmacist? A study on medical and pharmacy students' psychological state**R. Silva<sup>1,\*</sup>, M. Figueiredo-Braga<sup>1,2</sup><sup>1</sup> Medical Psychology Unit, Clinical Neurosciences and Mental Health Department, Faculty of Medicine, University of Porto, Portugal<sup>2</sup> Instituto de Inovação e Investigação em Saúde I3S, Porto, PortugalE-mail address: [rtg.silva@msn.com](mailto:rtg.silva@msn.com)

(R. Silva).

**Aim:** The objective of this study was to evaluate stress, anxiety, depression and happiness in medical and pharmacy students and to explore similarities and differences between them.**Introduction:** Higher levels of depression, anxiety and stress have been found in medical and pharmacy students when compared to general population,<sup>1,2</sup> varying across year in school and gender. Well-being during school years conversely may decrease depressive symptoms, boost happiness and life satisfaction, and contributes to resilience to stressful academic experiences. Students awareness of symptoms and consequences of distress may foster the search for psychoeducation and psychotherapy which offer effective strategies to improve mental health and academic performance.<sup>3,4</sup>**Methods:** A cross-sectional study included 420 students of Faculty of Medicine of the University of Porto (FMUP) and 200 students of Faculty of Pharmacy of the University of Porto (FFUP). Assessment included sociodemographic characterization, screening for anxiety and depressive symptoms - Hospital Anxiety and Depression Scale (HADS), stress - Perceived Stress Scale (PSS) and subjective wellbeing - Subjective Happiness Scale (SHS). One-way analysis of variance (ANOVA) and the independent paired t-test were applied to compare demographic and psychological characteristics from within each group.**Results:** Statistically significantly higher number of anxiety and depressive symptoms were found in medical students ( $p < 0.001$ ), and pharmacy students presented significantly higher PSS scores ( $p < 0.001$ ). Interestingly, medical students showed statistically significantly higher SHS scores than pharmacy students. Female students revealed significantly higher levels of anxiety, depression and stress in pharmacy school, but in medical school female students presented uniquely higher stress levels.**Conclusion:** Attending a faculty degree is a challenging experience which involves life changing experiences and poses different personal and academic problems according each specific school. These findings demonstrate the need to better understand the balance between students' stressful experiences and happiness to identify students at risk in both schools.**References**

- Hope V, Henderson M. Medical student depression, anxiety and distress outside North America: a systematic review. *Med Educ*. 2014;48:963–79.
- Gallagher CT, et al. Perceived stress levels among undergraduate pharmacy students in the UK. *Curr Pharm Teach Learn*. 2014;6:437–41.
- Duan W. The benefits of personal strengths in mental health of stressed students: a longitudinal investigation. *Qual Life Res*. 2016;1–10.
- Rajiah K, Saravanan C. The effectiveness of psychoeducation and systematic desensitization to reduce test anxiety among first-year pharmacy students. *Am J Pharm Educ*. 2014;78:163.

<http://dx.doi.org/10.1016/j.pbj.2017.07.145>

## PS234

**The relationship between socio-economic determinants and incidence of most common types of cancer in Poland**

Katarzyna Orlewska

Medical University of Warsaw, Poland

E-mail address: [korlewska@gmail.com](mailto:korlewska@gmail.com).**Aim:** To establish the link between incidence rates of cancer and selected socio-economic variables.**Introduction:** Geographical analysis of cancer incidence rates shows significant regional diversity and can be viewed as an approximation of the actual risk of particular types of cancer.**Methods:** The absolute numbers of new registered cases of lung, breast and colon cancer in Poland in 2014 by voivodeships (Polish provinces) were obtained from the Polish National Cancer Registry. The situation in individual voivodeships in terms of social isolation, social capital, religious activity and poverty was assessed based on the results of the Polish Social Cohesion Survey for 2015. The Spearman's rank correlation coefficient (rS) was used to test the association between incidence rates of types of cancer (number of cases/100 inhabitants) and social variables. The significance level was set at  $p < 0.05$  (2-tailed tests).**Results:** Spearman's correlation analysis showed a statistically significant strong positive correlation between lung cancer risk and: social isolation ( $rS = 0.73$ ;  $p < 0.0013$ ), living conditions poverty ( $rS = 0.55$ ;  $p = 0.028$ ), poverty resulting from the lack of budget balance ( $rS = 0.72$ ;  $p = 0.0015$ ), and low/no involvement in religious activity ( $rS = 0.7$ ;  $p = 0.003$ ). Strong negative correlation with  $rS = -0.64$  and  $p < 0.008$  exists between lung cancer risk and high level of association-based social capital. In colon cancer, only negative correlation between colon cancer risk and high level of friend- and neighbour-based social capital ( $rS = -0.56$ ;  $p = 0.020$ ) was statistically significant. Breast cancer risk was statistically significant for strong negative correlation with high level of friend- and neighbour-based social capital ( $rS = -0.74$ ;  $p = 0.0009$ ) and for a fairly strong positive correlation with low/no involvement in religious activity ( $rS = 0.53$ ;  $p = 0.04$ ).**Conclusion:** Our findings provide important evidence for the link between social and economic environment and the risk of most common cancer sites in Poland, and highlight the need to address these determinants as part of national cancer preventive programs.<http://dx.doi.org/10.1016/j.pbj.2017.07.146>

Surgery Poster Session

Friday, September 15th, 10h00

## PS093

**Multimodal analgesia after total knee joint arthroplasty surgery: Intrathecal morphine vs. local infiltration with ropivacaine**G. Bruzyte<sup>1,\*</sup>, G. Bukelyte<sup>1</sup>, E. Kontrimavičiute<sup>2</sup>, T. Strainys<sup>2</sup><sup>1</sup> Vilnius University, Faculty of Medicine, Vilnius, Lithuania<sup>2</sup> Clinic of Anaesthesiology and Intensive Care, Vilnius University, Faculty of Medicine, Vilnius, LithuaniaE-mail address: [gretbuckis@gmail.com](mailto:gretbuckis@gmail.com)

(G. Bruzyte).

**Aim:** To assess and compare effectiveness and side effects of postoperative anesthesia methods, using intrathecal morphine and

local infiltration of ropivacaine, a day after knee joint arthroplasty operations with spinal anesthesia.

**Introduction:** Inadequately chosen postoperative anesthesia method after knee joint arthroplasty surgery might cause prolonged hospitalization period, readmissions due to pain and overall increased cost of care.

**Methods:** In 2016 a prospective research was conducted in Vilnius University Hospital Santaros Clinics. 25 patients undergoing knee joint arthroplasty surgery with spinal anesthesia were enrolled in the study. Group1 – local soft tissue ropivacaine infiltration anesthesia around the knee ( $n = 13$ ; dose 300 mg); Group2 – intrathecal morphine sulfate analgesia ( $n = 12$ ; dose 0.1–0.2 mg). Pain intensity (using VAS) at rest and in motion, patient's satisfaction and side effects – nausea, vomiting, itch, urinary retention – were assessed at time intervals – 1, 2, 4, 6, 12, 18, 24 h postoperatively.

**Results:** In the first 12 h mean values of VAS were  $1.8 \pm 2.6/1.4 \pm 1.7$  in Group1 and Group2 accordingly. After 12 h period a downtrend occurred and values were  $1.7 \pm 1.1/1.1 \pm 1.5$ , respectively ( $p > 0.05$ ). Examining pain in motion 12 h after the surgery pain intensity values were  $2.5 \pm 2.7/3.3 \pm 2.7$  and after 24 h in both groups pain intensity was  $3.2 \pm 1.5/3.6 \pm 2.1$ , resp. ( $p > 0.05$ ). Zero episodes of nausea/vomiting were registered in Group1, while 58.3% ( $n = 7$ ) of Group2 patients experienced nausea and 5 of them also vomited. Even 66.7% ( $n = 8$ ) patients in Group2 had itch while none patients of Group1 indicated this side effect. It was difficult to assess urinary retention as 30.8% ( $n = 4$ ) Group1 and 66.7% ( $n = 8$ ) Group2 patients were catheterized prior surgery. Finally, satisfaction level of both groups were evaluated very similarly:  $8.2 \pm 1.7/8.2 \pm 1.3$  ( $p > 0.05$ ).

**Conclusion:** VAS values at rest were very similar in both groups, but pain relief efficiency compared to the intensity of pain during movement was better with local ropivacaine infiltration, also patients with ropivacaine analgesia experienced no side effects.

<http://dx.doi.org/10.1016/j.pbj.2017.07.147>

#### PS134

##### The role of cerebroplacental ratio in prediction of neonatal outcomes and route of delivery



P. Janas\*, A. Staroń, G. Wilczyńska,  
M. Brzozowska

Jagiellonian University Medical College, Cracow,  
Poland

E-mail address: [przemyslaw.janas@gmail.com](mailto:przemyslaw.janas@gmail.com)  
(P. Janas).

**Aim:** The aim of our study was to check the appropriability of cerebroplacental ratio (CPR) measured within 48 h before delivery in prediction of route of delivery and adverse neonatal outcomes.

**Introduction:** The cerebroplacental ratio is an important obstetric ultrasound tool used for assessment of foetal oxygenation. It is also a valuable predictor of adverse pregnancy outcomes. CPR is calculated by dividing the Doppler pulsatile indices of the middle cerebral artery (MCA) and the umbilical artery (UA).

**Methods:** The retrospective study included 1328 pregnant women who gave birth in Department of Obstetrics and Perinatology Jagiellonian University Medical College, Cracow, Poland. Main inclusion criteria were: singleton pregnancy and the interval between ultrasound examination and delivery within 48 h. Exclusion criteria consisted: active labour, multiple pregnancy, preeclampsia, foetal growth restriction and evidence of intrauterine infection. CPR value lower than 1.08 was classified as pathological. Participants were divided into 2 groups: control

( $CPR \geq 1.08$ ,  $n = 1228$ ) and study ( $CPR < 1.08$ ,  $n = 100$ ). The differences in socio-demographic factors between control and study group were not statistically significant. Data were analysed using chi-squared test, independent sample 2-tailed T-test and logistic regression.  $p$  value  $< 0.05$  was statistically significant.

**Results:** In study group was observed statistically significant increased risk of delivery provided by cesarean section ( $OR = 1.8$ ;  $p = 0.015$ ), preterm delivery ( $OR = 2.91$ ;  $p = 0.0001$ ), birth weight  $< 2500$  g ( $OR = 5.87$ ;  $p < 0.00001$ ) and APGAR score  $< 7$  in 1st ( $OR = 6.56$ ;  $p < 0.00001$ ), 3rd ( $OR = 7.04$ ;  $p < 0.00001$ ) and 5th ( $OR = 5.4$ ;  $p = 0.017$ ) minute after delivery, compared to control group. Moreover, low CPR was associated with lower incidence of foetus birth weight within normal limits ( $OR = 0.37$ ;  $p < 0.00001$ ) and on-term delivery ( $OR = 0.61$ ;  $p < 0.00001$ ).

**Conclusion:** Detection of low value of CPR in every case should be alarming signal for obstetrician. Normal CPR appears to suggest better foetal tolerance to the stress of labour. CPR may be used to stratify the risk of pregnancy before labour.

<http://dx.doi.org/10.1016/j.pbj.2017.07.148>

#### PS072

##### Chronic subdural hematoma in aging population – How the age influence the outcome after surgical treatment



Uladzislau Ulasavets\*, Ewelina Grzywna

Uniwersytet Jagielloński – Collegium Medicum,  
Poland

E-mail address: [vladvlasgeo@icloud.com](mailto:vladvlasgeo@icloud.com)  
(U. Ulasavets).

**Aim:** The aim of our work is to examine how the age influence the outcome after surgical treatment of chronic subdural hematoma.

**Introduction:** Chronic subdural hematoma (CSDH) is a common condition, characterized by the collection of hemolyzed blood between dura and arachnoid mater of the brain surrounded by two pathological hematoma membranes – internal and external. The number of CSDH incidence increases with age and it is why more attention should be directed for surgical treatment in elder patients group.

**Methods:** Data on management and outcomes for patients with CSDH were collected retrospectively from years 2014–2017 and investigated using statistic methods. The study group was divided into two subgroups according to the age:  $< 75$  years and  $\geq 75$  years old. Age, gender, comorbidities, neurological status on admission and at discharge, pre-/postoperative epilepsy, surgical technique were investigated.

**Results:** We analyzed 257 patients with a diagnosis CSDH. Analyzed subgroups have not differ significantly except the gender and concomitant diseases according to the Chi2 and exact Fisher tests. We found craniotomy in patients  $\geq 75$  years old increases the risk of postoperative epilepsy compering to the bur-hole (logistic regression analysis: 9.8 [95% CI: 1.9–49.8],  $p = .006$ ), same as the internal hematoma membrane removal during surgery (logistic regression analysis: 10.3 [95% CI: 2.0–52.15],  $p = .005$ ). These dependencies do not occur in the younger age group. Type of treatment have not influenced the mRS in patients younger than 75 years old. In elder patients reoperation and removal of the internal membrane of the hematoma worsened outcome measured in mRS (logistic regression analysis: 5.5 [95% CI: 1.4–20.90],  $p = .013$  and 3.1 [95% CI: 1.4–7.2],  $p = .007$ ).

**Conclusion:** Craniotomy and internal membrane removal increase the risk of epilepsy in elder CSDH patients. Reoperation

and hematoma internal membrane removal are the risk factors of unfavorable outcome in patients  $\geq 75$  years old.

<http://dx.doi.org/10.1016/j.pbj.2017.07.149>

### PS103

#### The frequency of MINS (Myocardial Injury after Noncardiac Surgery) and others postoperative complications in different age groups of elderly patients who underwent endovascular aortic repair because of abdominal aortic aneurysm



Gajdosz Anna\*, Kaszuba Aleksandra

Student Scientific Society at the Second Department of Internal Diseases UJ CM, Poland

E-mail address: [anna.gajdosz13@gmail.com](mailto:anna.gajdosz13@gmail.com)

(G. Anna).

**Aim:** The aim of our study was to estimate frequency of MINS and others complications after endovascular aorta repair because of AAA in different age groups.

**Introduction:** Nowadays, endovascular aneurysm repair (EVAR) is the most common technique for repair of abdominal aorta aneurysm (AAA). This procedure involves less complications than open surgery, nevertheless they still occurs.

**Methods:** The study group consisted of 143 patients (85.3% men), aged  $76.8 \pm 7.7$  with AAA who had endovascular aneurysm repair between January 2015 and May 2017 in the Department of Vascular Surgery and Angiology. Patients were divided into two groups depending on age: group I  $\leq 75$  years (60 patients, aged  $69.3 \pm 4.5$ ), group II  $> 75$  yrs (83 patients, aged  $82.2 \pm 4.2$ ). We considered coexistent diseases, some laboratory tests and Revised Cardiac Risk Index for Pre-Operative Risk (Lee index). Statistical analysis was performed with U Mann-Whitney and Chi2 tests.

**Results:** The study groups were comparable regarding the coexistent diseases and preoperative risk. Older patients had higher mean creatine level on admission than younger patients (group I:  $103.29 \pm 84.10$  vs. group II:  $118.5 \pm 61.9$   $\mu\text{mol/l}$ ,  $p < 0.005$ ) and lower eGFR ( $80.6 \pm 27.6$  vs.  $61.61 \pm 21.9$  vs.  $1.73$   $\text{ml/min/m}^2$ ,  $p < 0.001$ ). The mean concentration of haemoglobin and amount of white blood cells were also lower in elderly patient ( $13.19 \pm 1.93$  vs.  $12.9 \pm 18.3$   $\text{g/dl}$ ,  $p = 0.01$ ;  $8.23 \pm 2.74$  vs.  $7.49 \pm 2.7 \times 10^3/\mu\text{l}$ ,  $p = 0.04$ ).

Frequency of some complications such as acute kidney injury, pneumonia, sepsis, stroke or intrahospital mortality were similar in both groups.

However, we observed a statistically significant difference in the frequency of MINS (26.67% vs. 45.78%;  $p = 0.04$ ). Older patients also needed red blood cells concentrate transfusion after surgery more often than younger (6.67% vs. 19.28%;  $p = 0.03$ ).

**Conclusion:** MINS is the most common complication after EVAR. Age seems to be a significant feature which increases the frequency of MINS in compared groups despite similar coexistent diseases and preoperative risk assessment determined by Lee index.<sup>1-3</sup>

### References

- Sessler DI, Devereaux PJ. Perioperative troponin screening. *Anesth Analg*. 2016;123:359–60.
- Devereaux PJ, Sessler DI. Cardiac complications in patients undergoing major noncardiac surgery. *N Engl J Med*. 2015;373:2258–69.
- Jennifer A, Heller, et al. Two decades of abdominal aortic aneurysm repair: have we made any progress? *J Vasc Surg*. 2000;32:1091–100.

<http://dx.doi.org/10.1016/j.pbj.2017.07.150>

### PS210

#### Evaluation of clinical characteristics as indicators for shunt procedure in patients with medulloblastoma



A. Paunović\*, F. Milisavljević, J. Bošković

School of Medicine, University of Belgrade, Serbia

E-mail address: [aleksandra.paunovic92@gmail.com](mailto:aleksandra.paunovic92@gmail.com)

(A. Paunović).

**Aim:** Determining clinical characteristics and parameters reliable as predictors of the need for the shunt installation and their relation to the treatment outcome.

**Introduction:** Medulloblastoma represents the most common pediatric tumor, that most frequently involves posterior cranial fossa and often manifests as hydrocephalus. Current therapy involves tumor excision and posterior cranial fossa decompression, with or without temporary external drainage of cerebrospinal fluid, endoscopic ventriculocisternostomy and ventriculoperitoneal (VP) shunt placement.

**Methods:** This retrospective study included 36 patients treated in the period from January 1st 2007 to December 31st 2015 in the Clinic of Neurosurgery, Clinical Center of Serbia. Basic demographic data, symptoms and signs at admission, degree of tumor resection and disease outcome information were analyzed.

**Results:** 22 patients (61.1%) were male and 14 (38.9%) were female, most of them 4–14 years old (58.3). Sex and age showed no significant correlation with VP shunt installation, or timing of shunt installation. VP shunt was installed in 92% of patients, in 33.3% prior to and in remaining after surgery. The most frequently observed symptoms on admission were cerebellar symptomatology (91.2%), headache (75.7%) and vomiting (68.5%), which showed no significant correlation with the VP shunt installation and shunt installation timing. In 83% of patients total resection was achieved. The degree of tumor resectability and VP shunt installation were significantly related ( $p < 0.001$ ). Correlation among shunt installation and treatment outcomes, as well as the shunt installation timing and outcome showed a statistical significance ( $p < 0.001$ ).

**Conclusion:** No clinical characteristics reliable as prognostic parameter for VP shunt installation in medulloblastoma patients have been found. Shunt placement is recommended in all cases of incomplete tumor resection, unless already placed preoperatively. Patients with a shunt placed prior to surgery have had significantly better outcome.<sup>1-35</sup>

### References

- Lachi PK, Syed FAJD, Moinca I, et al. Medulloblastoma: a common pediatric tumor: prognostic factors and predictors of outcome. *Asian J Neurosurg*. 2015;10:50.
- Koeller KK, Rushing EJ. From the archives of the AFIP: medulloblastoma: a comprehensive review with radiologic-pathologic correlation. *Radiographics*. 2003;23:1613–37.
- Taylor MD, Northcott PA, Korshunov A, et al. Molecular subgroups of medulloblastoma: the current consensus. *Acta Neuropathol*. 2012;123:465–72.
- David NL, Arie P, Guido R, et al. The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol*. 2016;131:803–20.
- Albright AL, Pollack IF, Adelson PD. Principles and practice of pediatric neurosurgery. 3rd ed. New York: Thieme; 2014.
- Thompson EM, Hielscher T, Bouffet E, et al. Prognostic value of medulloblastoma extent of resection after accounting for molecular subgroup: a retrospective integrated clinical and molecular analysis. *Lancet Oncol*. 2016;17:484–95.
- Nikitovic MR, Golubicic IV, Borojevic ND, et al. Pediatric brain tumors—diagnostic and treatment. *Acta Chir Jugosl*. 2009;56:19–24.
- Nikitovic MR, Golubicic IV. Treatment options for childhood medulloblastoma. *Vojnosanit Pregl*. 2013;70:773–7.
- Fatema M, Ira JD. Treatment approach for recurrent medulloblastoma. *Clin Insights: Opt Therapy Pediatr Medulloblastoma*. 2015:59–73.
- Pilkington G, Walker D. Guidelines on the diagnosis and management of Adult PNETs. *Br Neuro-Oncol Soc/NCAT Rare Tumour Guidelines*. 2011.

11. Handler MH, Callahan B. Laparoscopic placement of distal ventriculoperitoneal shunt catheters. *J Neurosurg Pediatr.* 2008;2:282–5.
12. Naftel RP, Argo JL, Shannon CN, et al. Laparoscopic versus open insertion of the peritoneal catheter in ventriculoperitoneal shunt placement: review of 810 consecutive cases. *J Neurosurg.* 2011;115:151–8.
13. Roth J, Sagie B, Szold A, Elran H. Laparoscopic versus non-laparoscopic-assisted ventriculoperitoneal shunt placement in adults. A retrospective analysis. *Surg Neurol.* 2007;68:177–84.
14. Pettersson D, Schmitz KR, Jeffrey M, et al. Medulloblastoma: seeding of VP shunt tract and peritoneum. *Clin Pract.* 2012;2:e37.
15. Jennett B, Bond M. Assessment of outcome after severe brain damage. *Lancet.* 1975;7905:480–4.
16. Culley DJ, Berger MS, Shaw D, et al. An analysis of factors determining the need for ventriculoperitoneal shunts after posterior fossa tumor surgery in children. *Neurosurgery.* 1994;34:402–7 [discussion 407–8].
17. Riva-Cambrin J, Detsky AS, Lamberti-Pasculli M, et al. Predicting postresection hydrocephalus in pediatric patients with posterior fossa tumors. *J Neurosurg Pediatr.* 2009;3:378–85.
18. Albright L, Reigel DH. Management of hydrocephalus secondary to posterior fossa tumors. *J Neurosurg.* 1977;46:52–5.
19. Raimondi AJ, Tomita T. Hydrocephalus and infratentorial tumors. Incidence, clinical picture and treatment. *J Neurosurg.* 1981;55:174–82.
20. Hekmatpanah J, Mullan S. Ventriculo-caval shunt in the management of posterior fossa tumors. *J Neurosurg.* 1967;26:609–13.
21. Schneider C, Ramaswamy V, Kulkarni AV, et al. Clinical implication of medulloblastoma subgroups: incidence of CSF diversion surgery. *J Neurosurg Pediatr.* 2014;1–7.
22. Foreman P, McClugage S, Naftel R, et al. Validation and modification of a predictive model of postresection hydrocephalus in pediatric patients with posterior fossa tumors. *J Neurosurg Pediatr.* 2013;12:220–6.
23. Morelli D, Pirote B, Lubansu A, et al. Persistent hydrocephalus after early surgical management of posterior fossa tumors in children: is routine preoperative endoscopic third ventriculostomy justified? *J Neurosurg.* 2005;103:247–52.
24. Gnanalingham KK, Lafuente J, Thompson D, et al. The natural history of ventriculomegaly and tonsillar herniation in children with posterior fossa tumours—an MRI study. *Pediatr Neurosurg.* 2003;39:246–53.
25. Sainte-Rose C, Cinalli G, Roux FE, et al. Management of hydrocephalus in pediatric patients with posterior fossa tumors: the role of endoscopic third ventriculostomy. *J Neurosurg.* 2001;95:791–7.
26. Tamburrini G, Pettorini BL, Massimi L, et al. Endoscopic third ventriculostomy: the best option in the treatment of persistent hydrocephalus after posterior cranial fossa tumour removal? *Childs Nerv Syst.* 2008;24:1405–12.
27. Tuli S, Tuli J, Drake J, et al. Predictors of death in pediatric patients requiring cerebrospinal fluid shunts. *J Neurosurg.* 2004;100:442–6.
28. Drake JM, Kestle JRW, Milner R, et al. Randomized trial of cerebrospinal fluid shunt valve design in pediatric hydrocephalus. *Neurosurgery.* 1998;43:294.
29. Cochrane DD, Kestle JRW. The influence of surgical operative experience on the duration of first ventriculoperitoneal shunt function and infection. *Pediatr Neurosurg.* 2003;38:295–301.
30. McLaurin RL. Disadvantages of the preoperative shunt in posterior fossa tumors. *Clin Neurosurg.* 1983;30:286–92.
31. Taylor WA, Todd NV, Leighton SE. CSF drainage in patients with posterior fossa tumours. *Acta Neurochir(Wien).* 1992;117:1–6.
32. Di Rocco F, Juca CE, Zerah M, et al. Endoscopic third ventriculostomy and posterior fossa tumors. *WNEU.* 2013;79:S18.e15–9.
33. Bouras T, Sgouros S. Complications of endoscopic third ventriculostomy. *J Neurosurg Pediatr.* 2011;7:643–9.
34. Kulkarni AV, Drake JM, Mallucci CL, et al. Endoscopic third ventriculostomy in the treatment of childhood hydrocephalus. *J Pediatr.* 2009;155:254–9.
35. Drake JM. Canadian Pediatric Neurosurgery Study Group. Endoscopic third ventriculostomy in pediatric patients: the Canadian experience. *Neurosurgery.* 2007;60:881–6.

<http://dx.doi.org/10.1016/j.pbj.2017.07.151>

## PS009

### Percutaneous cholecystostomy in the management of acute cholecystitis

Sara Gomes-Rodrigues<sup>1,\*</sup>, Telma Vale-Fonseca<sup>1,2</sup>, Rui Mendes Costa<sup>1,2</sup>

<sup>1</sup> Faculty of Medicine of University of Porto, Porto, Portugal

<sup>2</sup> Department of General Surgery. Hospital of São João, Porto, Portugal

E-mail address: [sara.gomes.rodriques@gmail.com](mailto:sara.gomes.rodriques@gmail.com) (S. Gomes-Rodrigues).



**Aim:** The aim of this study is to clarify the role of percutaneous cholecystostomy in calculous acute cholecystitis treatment and to elucidate about its association with the surgical treatment.

**Introduction:** Laparoscopic cholecystectomy is the gold-standard treatment in acute cholecystitis. However, percutaneous cholecystostomy stands as an alternative therapeutic approach among the elderly or patients with several comorbidities.

**Methods:** In December 2016, a systematic database search on PubMed, Scopus and Web of Science was conducted to identify articles on percutaneous cholecystostomy published from January 2013 to November 2016, using the query “(acute cholecystitis OR severe cholecystitis) AND (cholecystostomy OR percutaneous cholecystostomy OR cholecystostomy tube)”. In total, 290 articles were found and submitted to inclusion and exclusion criteria.

**Results:** A total of 13 records involving 1130 patients from 10 different countries met all inclusion criteria and were therefore included in this systematic review. All studies found eligible concluded percutaneous cholecystostomy is a potentially safe and effective therapeutic approach among high-risk surgical patients in the setting of acute cholecystitis. Percentage of patients undergoing percutaneous cholecystostomy followed by cholecystectomy varied between 7.2% and a maximum of 66.7%, with a conversion rate fluctuating between 0.0% and 66.7%. Complication and mortality rates ranged from 2.2% to 41.7% and 0.0% and 43.2%, respectively.

**Conclusion:** Percutaneous cholecystostomy is generally considered safe and effective among high-risk surgical patients diagnosed with acute cholecystitis.<sup>1,2</sup>

## References

1. Popowicz A, Lundell L, Gerber P, et al. Cholecystostomy as bridge to surgery and as definitive treatment or acute cholecystectomy in patients with acute cholecystitis. *Gastroenterol Res Pract.* 2016;2016.
2. Jung W, Park D. Timing of cholecystectomy after percutaneous cholecystostomy for acute cholecystitis. *Korean J Gastroenterol.* 2015;66:209–14.

<http://dx.doi.org/10.1016/j.pbj.2017.07.152>

## PS092

### The influence of maternal age and parity on perinatal outcomes – A preliminary study



B. Adrianowicz

Perinatology Students' Scientific Group, Department of Obstetrics and Perinatology, Jagiellonian University Medical College, Poland

E-mail address: [beaadrr@gmail.com](mailto:beaadrr@gmail.com).

**Aim:** The aim of the study was to compare maternal, perinatal and neonatal outcomes depending on maternal age and parity.

**Introduction:** Advanced maternal age at childbirth has been associated with adverse perinatal and neonatal outcomes. As mean maternal age in developed countries is increasing decade by decade, the issue of perinatal outcomes among older patients seems to be of utmost importance.

**Methods:** It is a preliminary study that enrolled 243 women who gave birth in the Department of Obstetrics and Perinatology of the University Hospital in Kraków, Poland, during a one-month period (in May 2017). The patients were divided into 2 groups: >30 and ≤30 years old. The two groups were subsequently subdivided into 4 subgroups. Maternal, perinatal and neonatal outcomes were compared between all the subgroups.

**Results:** Comparison of women at age >30 and ≤30 revealed that advanced maternal age may constitute a predisposing factor for stillbirth, preterm delivery and congenital disorders. At the same time, the patients in the first group were at lower risk of SGA (small for gestational age) and LGA (large for gestational age)

as well as lack of progress in labour. Nevertheless, none of these parameters reached a statistical significance. Primiparas both  $\leq 30$  and  $>30$  year-old had a much greater risk of lack of progress in labour (OR = 11.3;  $p = 0.0015$  and OR = 19.5;  $p = 0.00027$ ) as well as emergency caesarean section (OR = 5.6;  $p = 0.00056$  and OR = 2.5;  $p = 0.027$ ) and a lower risk of elective C-section (OR = 0.4;  $p = 0.0027$  among women  $\leq 30$  years old) when compared to multiparas. These results met statistical significance criteria.

**Conclusion:** Advanced maternal age can undoubtedly be associated with several adverse perinatal outcomes. However, the results of the study do not seem to be unequivocal. Further investigation of the influence of maternal age on the course of pregnancy and delivery should be conducted.

<http://dx.doi.org/10.1016/j.pbj.2017.07.153>

## PS118

### Varicose veins surgical treatment: Endovenous laser ablation versus open surgery



Tomas Liaudginas\*, Vaida Kazlauskaitė

Faculty of Medicine, Vilnius University, Vilnius, Lithuania

E-mail address: [tliudginas@gmail.com](mailto:tliudginas@gmail.com)

(T. Liaudginas).

**Aim:** Compare long-term outcomes of varicose veins treatment between endovenous laser ablation and open surgery.

**Introduction:** 25–33% of the Western population adults is suffering from varicose veins and surgical treatment is characterized by high recurrence rate of 60% after 5 years follow-up observation. That leads to look for the most effective treatment option.

**Methods:** A retrospective study of 182 patients who had varicose veins endovascular or open surgery during 2015-06–2015-12 were made. Patients' demographic and clinical data were registered. Patients were contacted and asked to evaluate surgery outcomes subjectively. Patients were divided in two groups: I – ones who had endovenous laser ablation and II – those who had open surgery.

**Results:** 136 (74.7%) patients agreed to participate in the study: 26 (19.1%) men and 110 (80.9%) women; average age – 48.36 years ( $\pm 1.087$ ). 65 (47.8%) – group I, (41 (63.1%) were operated because of the symptoms, 24 (36.9%) were operated because of the esthetical reasons), 71 (52.2%) – group II, (58 (81.7%) were operated because of the symptoms, 13 (18.3%) were operated because of the esthetical reasons.) The symptoms renewed: group I – 9 (22%) patients, group II – 25 (43.1%). Esthetical relapse: group I – 3 (12.5%), group II – 5 (38.5%). Patients assessed esthetical view: group I – 48 (73.8%) were completely satisfied with the results, 15 (23.1%) were partly satisfied and 2 (3.1%) were unsatisfied, respectively in group II the results were 44 (62%), 26 (36.6%) and 1 (1.4%) ( $p = 0.202$ ). Patients also evaluated their symptoms after treatment. Group I: 53 (81.5%) patients were completely satisfied, 9 (13.8%) – partly satisfied, 3 (4.6%) – unsatisfied, respectively in group II – 52 (73.2%), 14 (19.7%) and 5 (7%) ( $p = 0.513$ ). According to the EQ-5D-3L questionnaire patients treated in group I evaluated their health – 80.29 ( $\pm 13.199$ ), II – 74.56 ( $\pm 16.615$ ).

**Conclusion:** Chronic venous insufficiency symptoms more often renewed to patients who had open surgery. Patients' were more satisfied after endovenous laser ablation. Also, those patients who were treated with endovascular procedure evaluated their health better.

<http://dx.doi.org/10.1016/j.pbj.2017.07.154>

## PS147

### Patients' quality of life after surgical treatment of oral cancer



A. Petrović

Faculty of Medicine, Novi Sad, Serbia

E-mail address: [andjus007@gmail.com](mailto:andjus007@gmail.com).

**Aim:** To evaluate the quality of life of patients after surgical treatment of oral cancer.

**Introduction:** Oral cancer surgery may have resulted in damage to the act of chewing, swallowing, speech and communication, lead to social incompetence and decline in the quality of life of patients.

**Methods:** The study included 30 surgically treated patients previously diagnosed with oral cancer at the Department of Maxillofacial Surgery, Clinical Center in Novi Sad. The survey and assessment of Karnofsky index of patients was conducted within the regular control visits to a specialist clinic for maxillofacial surgery. The questionnaire is in addition to general information included questions related to the primary tumor localization, harmful habits, stomatognathic system function and psychosocial condition after surgical treatment of oral cancer.

**Results:** Mean age of patients was 63 years, of which 73.3% were male and 26.7% female. The most common localization of oral cancer was under the mouth of 9 (30%) patients. Alcohol and cigarettes were consumed by 70% of patients. In 50–60% of patients, stomatognathic system function was reduced to varying degrees of severity. Chewing function was statistically significantly worse in patients after marginal resection of the mandible ( $p < 0.05$ ). Depression was observed in 18.2% of men and 12.5% women, and the problem of appearance in public places had 40% of patients. Karnofsky index of 25 (83.3%) patients was 100% while the remaining was lower as a result of comorbidity.

**Conclusion:** The quality of life in terms of stomatognathic system function and psychosocial status was reduced to varying degrees of expression in many patients after surgical treatment of oral cancer. Therefore, the detection of diseases in an early stage is crucial to the survival and quality of life for patient.

<http://dx.doi.org/10.1016/j.pbj.2017.07.155>

## PS221

### Long-term outcomes in simultaneous pancreas-kidney transplant recipients: Retrospective single centre study



Gniewkiewicz Michał<sup>1,2</sup>,  
Czerwińska Magdalena<sup>1,2,\*</sup>

<sup>1</sup> Student Scientific Group of Transplantation Medicine and Nephrology, Poland

<sup>2</sup> Department of Transplantation Medicine, Nephrology and Internal Medicine, Medical University of Warsaw, Poland

E-mail address: [mczerwinskam@gmail.com](mailto:mczerwinskam@gmail.com)  
(C. Magdalena).

**Aim:** The aim of this study is to present long-term outcomes of SPKT.

**Introduction:** Simultaneous pancreas-kidney transplantation (SPKT) is the treatment of choice for patients with end-stage renal failure due to type 1 diabetes mellitus (DM1).

Since the 1980s, pancreas transplant has become the most effective strategy to restore normoglycemia in patients with DM1.

**Methods:** We performed a retrospective analysis of 73 SPKT recipients who underwent transplantation between 1988 and 2015.

**Results:** 50.68% of patients were male. During the time of surgery the mean age was  $37.8 \pm 7.44$  years. DM1 was diagnosed average  $25 \pm 6.08$  years before SPKT. For 21.9% it was pre-emptive transplant. 60.9% and 17.19% were on haemodialysis and CADO respectively (the mean dialysis time was 29.05 months). Reoperation due to pancreatic and kidney complications amounted respectively 23.3% vs 8.3%. DGF was observed in 9.6% of kidney graft recipients. Mean HLA - A, - B, - DR mismatches were: 1.42, 1.58, 1.27. All patient received induction of immunosuppression (polyclonal immunoglobulins: ATG/Thymoglobulin – 64% or monoclonal: daclizumab/basiliximab – 36%). Kidney graft survival at 1, 5, 10, 15 years 100%, 97%, 85% and 67%; and pancreas survival is 95%, 92%, 87% and 67% respectively. There was noticed tendency to increase creatinine level (from 1.18 at 1 year to 1.78 at 15 years) and decrease of haemoglobin level (from 13.84 at 1 year to 12.65 at 15 years). Patients with longer time of dialysis were more commonly infected by HCV ( $p=0.004$ ), more often hospitalized due to cardiovascular complications ( $p=0.004$ ) and had shorter survival time ( $p=0.03$ ). HBV infection correlated with longer time of hospitalization during transplantation procedure ( $p=0.006$ ), more often delay grant function of pancreas ( $p=0.008$ ), higher serum level of CRP ( $p=0.04$ ) and more frequent hospitalizations in subsequent years ( $p=0.003$ ).

**Conclusion:** Shorter dialysis time improves patient prognosis after SPKTx. HBV and HCV infection is associated with more frequent complications and worse prognosis. Cardiovascular complications are more likely to affect dialysis patients.

<http://dx.doi.org/10.1016/j.pbj.2017.07.156>

### PS031

#### The impact of suturing hemostasis on ovarian reserve during conserving surgeries on the ovaries



A.A. Solomatina, E.A. Tumasyan\*, I.Z. Khamzin

*N.I. Pirogov Russian National Research Medical University, Russia*

*E-mail address: elizavetatumasyan@gmail.com (E.A. Tumasyan).*

**Aim:** To determine the effects of using suturing hemostasis in cases of cystectomy of unilateral endometriomas and mature teratomas (MT).

**Introduction:** Nowadays there is a noticeably growing rate of benign ovarian tumors requiring urgent treatment. It is known that ovarian tumors lead to diminished ovarian reserve (OR). Conserving surgeries bring to a further reduction of OR.

**Methods:** The study involved 66 patients with endometriomas and 69 with MT. The mean age was  $28.07 \pm 5.3$ . All patients underwent laparoscopic cystectomy. The methods to stop bleeding were: ligature hemostasis with absorbable polyglycolic suture, USP 2-0 (I group) and bipolar coagulation (BPC) – Autcon II 350, current power 35 W – (II group). Before and 6–12 months after surgery serum levels of Antimüllerian hormone (AMH) were evaluated; by ultrasound (Toshiba Aplio 500, 3.6–8.8 MHz) we measured the volume of healthy ovarian tissue ( $V_{cm^3}$ ), antral follicle count (AFC), their site and diameter.

**Results:** At the pre-surgical stage patients with endometriomas had reasonably lower ultrasound and biochemical markers than patients with MT. 6 months after suturing hemostasis patients with endometriomas had a 1.8 times higher AFC, its diameter and, as a

result, the volume of ovarian tissue of the operated gonad compared to the group after BPC. Studying similar indices by the patients with MT showed the difference of 1.3 times respectively. Deformed follicles with small diameter (3–4 mm) ousted to the periphery were located on the echograms. Six months after laparoscopy the AMH level of all patients decreased, the biggest reduction (1.7 and 1.9 times correspondingly to groups) was noted by the patients with endometriomas.

**Conclusion:** To preserve women's reproductive potential after conserving surgeries on the ovaries, intracorporeal suturing is a preferred hemostatic method over bipolar energy. Enucleation of endometriomas and MT leads to diminished OR regardless of the energy type used as a hemostasis.

<http://dx.doi.org/10.1016/j.pbj.2017.07.157>

### PS102

#### Complications after open surgery for the abdominal aorta and its branches depending on patients' age



Kaszuba Aleksandra\*, Gajdosz Anna, Iwańska Anna, Kacorzyc Radosław

*Student Scientific Society at the Second Department of Internal Diseases UJ CM, Poland*

*E-mail address: aleksandra.kaszuba.94@gmail.com (K. Aleksandra).*

**Aim:** Age is one of the risk factors for postoperative complications in open surgery of the aorta. The awareness of their frequency may lead to earlier diagnosing and referral for procedure in order to avoid negative results of surgery and further therapy.

**Introduction:** The aim of our study was to assess the dependent of age frequency of postoperative complications among patients undergoing primary open surgical procedure within abdominal aorta due to aneurysm (AAA) or/and peripheral arterial disease (PAD).

**Methods:** The study group consisted of 249 patients (84.7% men), aged  $69.1 \pm 8.2$  with AAA or/and PAD who underwent open abdominal aorta surgery between August 2015 and January 2017. Patients were divided into three groups depending on age group I < 65 years (61 patients, aged  $58.8 \pm 5$ ), group II 65–74 yrs (118 patients, aged  $68.5 \pm 2.8$ ) and group III > 74 yrs (70 patients, aged  $79.1 \pm 3.4$ ). We considered coexistent diseases, smoking habit, Revised Cardiac Risk Index for Pre-Operative Risk (Lee index), some laboratory tests, type and mode of surgery (elective vs urgent/emergent). Statistical analysis was performed with Kruskal Wallis and Chi2 tests.

**Results:** Frequency of some complications such as myocardial infarction, pneumonia, sepsis, stroke or bleeding was similar in compared groups. Nevertheless, we observed a statistically significant difference in the frequency of acute kidney injury demanding dialysis (respectively, for groups I–III: 3.28% vs. 17.80% vs. 20.00%;  $p=0.013$ ), multi-organ failure (1.64% vs. 10.17% vs. 14.29%;  $p=0.039$ ) and intrahospital mortality (1.64% vs. 11.86% vs. 18.57%;  $p=0.009$ ). The groups were comparable regarding the coexistent diseases – the only differentiative feature was hypertension that occurred less in group I (62.30%) than in other groups (82.20% and 81.43%,  $p=0.01$ ). There was no significant difference between groups in preoperative risk determined by Lee index and mode of surgery procedure. However, older patients had higher mean creatine level on admission than younger patients ( $72.2 \pm 21.5$  vs.  $91.3 \pm 34.2$  vs.  $94.4 \pm 37.7$   $\mu\text{mol/l}$ ,  $p < 0.005$ ), lower eGFR ( $109 \pm 31$  vs.  $84.5 \pm 29.1$  vs.  $73.8 \pm 24.7$   $\text{ml/min/1.73 m}^2$ ,  $p < 0.001$ ) and more

often underwent procedure due to ruptured aneurysm (4.92% vs. 9.32% vs. 11.43%,  $p < 0.001$ ).

**Conclusion:** Age is a significant feature which increases the frequency of acute kidney injury, multi-organ failure and intra-hospital mortality in compared group despite similar coexistent diseases, mode of surgery and preoperative risk determined by Lee index.<sup>1,2</sup>

## References

1. Thomas S, Huber SAL. Experience in the United States with intact abdominal aortic aneurysm repair. *J Vasc Surg.* 2001;33.
2. Jennifer A, Heller MDSAL. Two decades of abdominal aortic aneurysm repair: have we made any progress? *J Vasc Surg.* 2000;32.

<http://dx.doi.org/10.1016/j.pbj.2017.07.158>

## PS076

### Modern treatment and diagnostics for submucosal tumors of the upper third of the esophagus. Analysis of preoperational and postoperational data

A.N. Burakov\*, A.A. Smirnov

First Pavlov State Medical University of St. Petersburg, Russia

E-mail address: [aleksandr.medox@yandex.ru](mailto:aleksandr.medox@yandex.ru)

(A.N. Burakov).



**Aim:** This study was to evaluate diagnostic data and outcomes of endoscopic treatment for submucosal tumors (SMTs) of the upper third of the esophagus.

**Introduction:** Esophageal submucosal tumors (SMTs) are very rare, with prevalence of 0.5% in autopsy series. Among them leiomyomas are the most common, they originate from the muscularis propria (4th EUS layer) or muscularis mucosa (2th EUS layer) of the esophageal wall.<sup>1</sup> Submucosal lesions of the upper third is very rare and occur in 4% of cases.<sup>2</sup> Submucosal tunneling endoscopic resection (STER) and endoscopic submucosal dissection (ESD) are modern techniques for treating SMTs. The choice between them depends on layer of origin of the tumor.

**Methods:** In this study were included 2 patients with SMT of the upper third of the esophagus. For diagnostics we used esophageal symptoms questionnaire, endoscopic ultrasonography (EUS) and/or computed tomography (CT) to determine layer of origine, size and relation of lesions to the surrounding structures and organs. Esophageal manometry were used to identify problems with movement and pressure in the esophagus. Immunohistochemistry and histological analysis were performed postoperatively.

**Results:** Both patients were asymptomatic, tumors were found accidentally during routine esophagoscopy. Although in both cases manometry of the esophagus revealed increased distal latency (DL) comparing with mean value in patient without esophageal SMT. In the first case tumor arised from 4th EUS layer, hence we used STER, subcutaneous emphysema of the neck occured during operation. in the second case lesion originated from 2th EUS layer, therefore ESD was performed. En bloc resection was achieved in both cases, histological diagnoses were leiomyomas.

**Conclusion:** Upper third of the esophagus is the most difficult location for performing endoscopic techniques. Determination of the layer of origin is crucial, as on that depends the choice of treatment tactics. Increased DL in such patients requires further study.

**Acknowledgements:** We would like to show our gratitude to the Rector of our university, S.F. Bagnenko, for the support of this study.

## References

1. Rice TW. Benign esophageal tumors: esophagoscopy and endoscopic esophageal ultrasound. *Semin Thorac Cardiovasc Surg.* 2003;15:20–6.
2. Mutrie CJ, et al. Esophageal leiomyoma: a 40-year experience. *Ann Thorac Surg.* 2005;79:1122–5.

<http://dx.doi.org/10.1016/j.pbj.2017.07.159>

## PS035

### The role of the state of uterine-placental-foetal circulation on the clinical course of gestational process and its impact on perinatal outcome



Burra Mithilesh<sup>1,\*</sup>, Varahabhatla Vamsi<sup>1</sup>, Katnam Sahithi<sup>1</sup>, Nataliya Gaidai<sup>2</sup>

<sup>1</sup> Students of 4th Year Medicine, Department of Obstetrics and Gynaecology, Zaporozhye State Medical University, Ukraine

<sup>2</sup> Scientific Tutor, Associate Professor, Department of Obstetrics and Gynaecology, Zaporozhye State Medical University, Ukraine

E-mail address: [mithil.bharadwaj@gmail.com](mailto:mithil.bharadwaj@gmail.com)

(B. Mithilesh).

**Aim:** To identify the relationship between the condition of utero-placento-foetal circulation with the clinical course of gestational process and its impact on perinatal outcome.

**Introduction:** The period of foetal development before birth is so extensive however, only a small part of the duration of this period, which largely determines the quality of his later life. It is a proven fact that the events during the prenatal period effects the outcomes of pregnancy which are favourable in childbirth, later leading to diseases in adulthood.

**Methods:** We analysed the course of pregnancy, delivery, the condition of the foetus and newborn from 72 pregnant women (24–41 weeks of gestation) with placental dysfunction at the 3rd Maternity Hospital, Zaporozhye.

**Results:** According to CT, distress of the foetus were confirmed in 22.7% of pregnant women with impaired hemodynamics I-A degree, 24.8% with impaired hemodynamics, at 30.6% with circulatory disorders of the II degree. On analysis of the hemodynamics in the system of maternal-placento-foetal revealed violations of IPC(I-A) in 46% of cases, ACC(I-B) at 28.7%, IPC and SPC(II) at 12.7%, critical blood vessels PPK(III) and 3.4% of cases. The frequency of caesarean section in pregnant women with dysfunction of placenta was 28.2%, of which the foetal distress was 22.4%, vacuum extraction of the foetus were used in 3.2%. The analysis of the development of newborn from mothers with placental dysfunction, identified the violation of their status at birth and Apgar score 7–5 points received at birth 11.2% of newborn.

**Conclusion:** Analysis of indicators of physical development of newborns in the early neonatal period were distinguished by the presence of signs of functional immaturity. Clinical and statistical analysis conducted revealed a high frequency of complications of pregnancy and childbirth in women with dysfunction of the placenta.

**Acknowledgements:** I would like to express my special thanks of gratitude to my associate professor Nataliya Victorovna Gaidai as well as my co-authors Varahabhatla Vamsi and Katnam Sahithi who gave me this golden opportunity to do this wonderful project on the topic. The role of the state of uterine placental foetal circulation

on the clinical course of gestational processes and its impact on perinatal outcome. Secondly, I would also like to thank my parents and friends who helped me a lot in finalizing this project within the limited time frame.

<http://dx.doi.org/10.1016/j.pbj.2017.07.160>

#### PS014

##### The possibility of optimization of hemodynamics in the fetoplacental pool as a factor of influence on perinatal outcome



Vinisha Tekwani<sup>1,\*</sup>, Varahabhatla Vamsi<sup>1</sup>,  
Katnam Sahithi<sup>1</sup>, Nataliya Gaidai<sup>2,\*</sup>

<sup>1</sup> Students of 4th year Medicine, Ukraine

<sup>2</sup> Scientific Supervisor, Associate Professor, Ukraine

E-mail address: [vinishatekwani33@gmail.com](mailto:vinishatekwani33@gmail.com)  
(V. Tekwani).

**Aim:** To study the possibility of optimization of hemodynamics in the fetoplacental pool as a factor of influence on perinatal outcome.

**Introduction:** Endothelial dysfunction in uteroplacental pool is a universal response of placenta to adverse effects of hypoxia, which leads to a high percentage of obstetric complications. Recreation is a way of optimization of hemodynamics in fetoplacental complex in the interests of antenatal protection of the foetus.

**Methods:** The study was conducted at the 3rd Maternity hospital, Zaporozhye including 40 pregnant women with VD with age group of 21–36 years (Primipara – 52.5%, multipara is 47.5%). 40 pregnant women with chronic venous insufficiency to restore homeostasis used the IR thermo-camera, designed and implemented by the Department of Clinical Pathophysiology, Institute of physiology. Pregnant women of the main group underwent 3 sessions of IR sonotherapy (1 time per week), lasting 30 min at temperature of 35 °C.

**Results:** Pregnant women with VD after using sonotherapy in the infrared heat chamber in the complex sanatorium treatment, on comparison with the control group, a more pronounced therapeutic effect of lowering body weight by  $22.3 \pm 1.2\%$ , and decrease of systolic  $14.6 \pm 0.2$  mmHg and diastolic  $15.1 \pm 1.1$  mmHg pressure. Ended pregnancy in a core group of women, the birth of full-term newborns with no signs of distress, with an Apgar score of 7–9 points, body mass 2980–4000 g. 1 in the case of birth by caesarean section for obstetric indications.

**Conclusion:** The research conducted in the sanatorium “Velikii Lug”, confirms the effectiveness of the use of the IR sonotherapy in optimizing antenatal protection of the fetus against the background of endothelial dysfunction.

**Acknowledgements:** I wish to express my sincere gratitude to my scientific supervisor Mrs. Nataliya Viktorovna Gaidai for providing me an golden opportunity to be a part in this project. I also sincerely thank my co-authors Mr. Vamsi Varahabhatla and Katnam Sahithi for their guidance and encouragement in carrying out this project work. Finally, I would like to thank my family for supporting me financially and morally.

<http://dx.doi.org/10.1016/j.pbj.2017.07.161>

#### PS090

##### Bile duct injuries after cholecystectomy: A retrospective tertiary centre study comparing outcomes of different types of surgical treatment



R. Zulpaitė<sup>1,\*</sup>, A. Sileikyte<sup>1</sup>, A. Sileikis<sup>2</sup>

<sup>1</sup> Faculty of Medicine, Vilnius University, Vilnius, Lithuania

<sup>2</sup> Center of Abdominal Surgery, Vilnius University Hospital, Santaros Klinikos Santariskiu str. 2, 08661 Vilnius, Lithuania

E-mail address: [ruta.zulp@gmail.com](mailto:ruta.zulp@gmail.com)  
(R. Zulpaitė).

**Aim:** Evaluation of long-term outcomes after different types of surgical management of postcholecystectomy bile duct injuries (BDI).

**Introduction:** Cholecystectomy is one of the most routinely performed procedures in abdominal surgery. Despite the growing experience of surgeons and benefits of minimal invasive approach, BDIs still occur. The treatment of this complication is challenging.

**Methods:** This was a single-center retrospective study. The outcomes of 64 consecutive adult patients, surgically treated after postcholecystectomy BDI 2002–2016, were reviewed. The newest EAES ATOM classification was used to describe injuries. The anatomic characteristics of the injury and long-term treatment outcomes were evaluated.

**Results:** 48 (75%) BDI followed laparoscopic cholecystectomy. 26% of injuries were detected intraoperatively, 58% detected <7 days, 16% >7 days after the procedure. The injury of non-main bile duct was diagnosed in 10 (16%) cases. The injuries of main bile duct: choledochal duct 22 (34%), hepatic duct 22 (34%), bifurcation with right-left communication preserved 5 (8%), bifurcation with right-left interrupted 1 (2%), right/left hepatic duct 4 (6%). 26 (41%) patients with a cystic stump leak or partial division of duct were managed endoscopically. This treatment was successful for 7 (88%) cystic stump leaks and 8 (58%) partial divisions. 13 (20%) partial divisions of duct were closed by suture. 8 (73%) patients had complications which later required endoscopic management or hepaticojejunostomy. End-to-end anastomosis (6 (10%)) or hepaticojejunostomy (16 (25%)) was initially performed after the complete division with or without loss of substance was detected. End-to-end strategy was successful in 4 (67%) cases, others finally required hepaticojejunostomy. The complication rate after initial hepaticojejunostomy - 25%.

**Conclusion:** Endoscopic treatment is optimal for cystic stump leaks and partial divisions of ducts. Complete divisions with or without loss of substance may be treated by hepaticojejunostomy and end-to-end anastomosis with similar long-term outcomes. While end-to-end anastomosis is more physiological, this strategy should be considered when possible.

<http://dx.doi.org/10.1016/j.pbj.2017.07.162>

#### PS041

##### Perinatal loss in multiple pregnancies



L.G. Sichinava, A.O. Dulaeva\*, D.S. Spiridonov

Pirogov Russian National Research Medical University (RNRMU), Russia

E-mail address: [littleinwonderland@gmail.com](mailto:littleinwonderland@gmail.com)  
(A.O. Dulaeva).

**Aim:** The aim of the study was to analyze causes of perinatal loss in multiple pregnancies.

**Introduction:** In population rate of multiple pregnancies varies from 0.7 to 1.5%. Multiple pregnancies are complicated by perinatal loss 4–9 times more frequently than singleton pregnancies.

**Methods:** Retrospective study of medical histories was carried out. Thirty patients with twin pregnancy and perinatal loss of one or both fetuses were included. Thirteen (43.3%) twins were mono-chorionic (MC), 17 (56.7%) – dichorionic (DC). At 11–14 week of gestation chorionicity was diagnosed by ultrasound; transvaginal measurement of cervix was performed at 19–21 week; biometry was done to identify degree of fetus' discordance.

**Results:** Complications of DC pregnancy: discordant fetal growth – 17 (100%), fetal growth restriction – 7 (41.2%), cervical insufficiency – 4 (23.5%). Discordant fetal growth was diagnosed in 17 DC twins: 8 (47.1%) –  $\leq 20\%$ , 9 (52.9%) –  $> 20\%$ . 8 (47.1%) patients with discordance  $> 25\%$  had highest degree of fetal growth restriction (estimated fetal weight  $< 5\%$ ).

Perinatal loss in patients with DC twins was 61.8% (21 of 34 children). Highest mortality [10 of 21 (47.7%)] was among newborns at 22–27 week of gestation with DC type of placentation: 7 – intrauterine death, 3 died postnatally. Seventeen cases of intrauterine death were diagnosed: 7 (41.2%) – 22–27 weeks, 3 (17.6%) – 28–31 weeks, 5 (29.4%) – 35–36 weeks, 2 (11.8%) – at term.

Complications of MC pregnancy: discordant fetal growth – 13 (100%), twin-to-twin transfusion syndrome (TTTS) – 11 (84.6%), fetal growth restriction – 9 (69.2%), cervical insufficiency – 4 (30.8%). Discordant fetal growth was diagnosed in 13 MC twins: 7 (53.8%) –  $\leq 20\%$ , 6 (46.2%) –  $> 20\%$ . Four (30.8%) patients with discordance  $> 25\%$  had selective fetal growth restriction.

Perinatal loss in patients with MC twins was 80.8% (21 of 26 children). Highest mortality [13 of 21 (61.9%)] was among newborns at 22–27 week of gestation: all of them died antenatally. Nineteen cases of intrauterine death were diagnosed: 13 (68.4%) – 22–27 weeks, 4 (21.0%) – 28–31 weeks, 1 (5.3%) – 35–36 weeks, 1 (5.3%) – at term.

**Conclusion:** There were 1.3 times more perinatal losses in MC twins than in DC twins (80.8% vs. 61.8%). Regardless of chorionicity, perinatal losses were observed more frequently at 22–27 weeks of gestation: DC (47.7%) and MC (61.9%) twins. Causes of perinatal loss in DC twins: prematurity – 52.9%, discordant fetal growth ( $> 20\%$ ) – 52.9%, fetal growth restriction – 41.2%. Causes of perinatal loss in MC twins: TTTS – 84.6%, prematurity – 76.9%, fetal growth restriction – 69.2%, discordant fetal growth ( $> 20\%$ ) – 46.2%.

<http://dx.doi.org/10.1016/j.pbj.2017.07.163>

PS230

### Hyaluronic acid solution as a treatment of adhesive intestinal obstruction in children – A positive effect



M.A. Isa\*, O.B. Bodnar

*Bukovinian State Medical University, Department of Paediatric Surgery and Otolaryngology, Ukraine*  
E-mail address: [mashforreal@yahoo.com](mailto:mashforreal@yahoo.com)  
(M.A. Isa).

**Aim:** To explore the possibility of using hyaluronic acid solution (HAS) for the treatment of intraperitoneal adhesions in children.

**Introduction:** Adhesive intestinal obstruction (AIO) has been found to be a challenging problem of abdominal surgery with increased occurrence in children worldwide. Intraperitoneal adhesions occur commonly after abdominal surgery and frequently cause intestinal obstruction. Current means of adhesion prevention includes good surgical technique and anti-adhesion barriers. This study is hence directed towards the effect of hyaluronic acid solution (HAS) in reducing the incidence and recurrence of adhesions.

**Methods:** 84 children were operated on for AIO. 21 children (25%) were operated on for early adhesive intestinal obstruction (EAIO), 63 (75%) – on late adhesive intestinal obstruction (LAIO) and 12 (14.29%) for recurrent AIO. Following surgery, these children were divided into two groups; group I (56 patients) and group II (28 patients). The Hyaluronic Acid Solution; Defensal was used. Follow-up on the children took place from 1 to 4 years.

**Results:** 13 children (23.21%) in group I were found to have adhesion syndrome in the first year after surgery. This increased to 20 (35.71%) patients over the 4 year period. Children in the II group who had undergone treatment for adhesion syndrome (cured conservatively using HAS) over a 2.5 year postoperative period were not found to have adhesive syndrome at the end of the follow-up period with the exception of 2 (7.14%) patients. When compared to group I patients who had no treatment by the HAS, group II patients showed a higher degree of recovery with minimal recurrence.

**Conclusion:** Although accompanied by a minimal recurrence rate, HAS shows effectiveness as a treatment for adhesive intestinal obstruction in children. This serves as a step further towards a complete prevention of postoperative adhesion common in children.<sup>1–11</sup>

### References

1. Alwan MH, van Rij AM, Greig SF. Postoperative adhesive small bowel obstruction: the resources impacts. *N Z Med J.* 1999;12:421–3.
2. Wilkins BM, Spitz L. Incidence of postoperative adhesion obstruction following neonatal laparotomy. *Br J Surg.* 1986;73:762–4.
3. Festen C. Postoperative small bowel obstruction in infants and children. *Ann Surg.* 1982;196:580–3.
4. Janik JS, Ein SH, Filler RM, et al. An assessment of the surgical management of adhesive small bowel obstruction in infants and children. *J Pediatr Surg.* 1981;16:225–9.
5. Vijay K, Anindya C, Bhanu P, Mohan M, Rao PL. Adhesive small bowel obstruction (ASBO) in children—role of conservative management. *Med J Malaysia.* 2005;60:81–4.
6. Akgur FM, Tanyel FC, Buyukpamukcu N, Hicsonmez A. Adhesive small bowel obstruction in children: the place and predictors of success for conservative treatment. *J Pediatr Surg.* 1991;26:37–41.
7. ten Broek RP, Issa Y, van Santbrink EJ, Bouvy ND, Kruiwagen RF, Jeekel J, Bakkm EA, Rovers MM, van Goor H. Burden of adhesions in abdominal and pelvic surgery: systematic review and met-analysis. *BMJ.* 2013;347:f5588.
8. Loftus T, Moore F, VanZant E, Bala T, Brakenridge S, Croft C, Lottenberg L, Richards W, Mazingo D, Atteberry L, et al. A protocol for the management of adhesive small bowel obstruction. *J Trauma Acute Care Surg.* 2015;78:13–9 [discussion 19–21].
9. Di Saverio S, Catena F, Kelly MD, Tugnoli G, Ansaloni L. Severe adhesive small bowel obstruction. *Front Med.* 2012;6:436–9.
10. Okabayashi K, Ashrafian H, Zacharakis E, Hasegawa H, Kitagawa Y, Athanasiou T, Darzi A. Adhesions after abdominal surgery: a systematic review of the incidence, distribution and severity. *Surg Today.* 2014;44:405–20.
11. Catena F, Di Saverio S, Cocolini F, Ansaloni L, De Simone B, Sartelli M, Van Goor H. Adhesive small bowel adhesions obstruction: Evolutions in diagnosis, management and prevention. *World J Gastrointest Surg.* 2016;8:222–31.

<http://dx.doi.org/10.1016/j.pbj.2017.07.164>