

## PS096

### The influence of smoking cessation-related weight gain on cardiovascular risk in patients treated with drug-eluting stent after acute coronary syndrome

Aleksandra Buczyńska\*, Karol Kasprzycki, Aleksandra Pizun, Marta Tomica

Students' Scientific Group, The Department of Coronary Heart Disease, The John Paul II Hospital in Cracow Faculty of Medicine, Jagiellonian University Medical College, Poland

E-mail address: [abuczynska93@gmail.com](mailto:abuczynska93@gmail.com) (A. Buczyńska).

**Aim:** Aim of this study was to analyse association of smoking cessation influence on long-term clinical outcomes in patients with an acute coronary syndrome (ACS).

**Introduction:** Smoking and obesity are important cardiovascular risk factors. Patients often put on weight after quitting smoking.

**Methods:** 137 consecutive ACS patients of the Department of Coronary Heart Disease John Paul II Hospital in Cracow admitted between 2011 and 2013 were enrolled in the study. They had no previous history of Coronary Heart Disease and underwent Percutaneous Coronary Intervention with implantation of at least one Drug Eluting Stent. Telephone follow-up was carried out after a minimum time of 3 years. Patients were divided into 2 groups: patients who stopped smoking (91) and non-smokers (46) which were compared according to weight gain, increased of the BMI, morbidity of diabetes mellitus (DM), reasons of admission to hospital, occurrence of another ACS, stroke and neoplasm.

**Results:** The population consisted of 66% males, 34% females mean age 67 SD 11.29. Patients who used to smoke were significantly younger than non-smokers (64.99 vs. 71.37;  $P=0.048$ ). Both groups did not statistically differ in terms of gender and frequency of DM. The ex-smokers were admitted more frequently due to STEMI while in the non-smokers NSTEMI and Unstable Angina predominated. The weight and BMI in both groups did not differ on the date of ACS. However after 3 years there was a statistically significant difference: ex-smokers put on weight on average 1.3 kg while non-smokers lost 2.17 kg ( $p=0.01$ ). There was no association between the patients' history of smoking and occurrence of stroke, malignancy or another ACS.

**Conclusion:** Smoking cessation does not appear to influence long-term clinical outcomes after ACS. However it associates with weight gain which obviously increases cardiovascular risk. Our findings need further investigation and follow-up in a larger cohort of ACS patients.

<http://dx.doi.org/10.1016/j.pbj.2017.07.049>

## PS059

### Neonatal abstinence syndrome – Retrospective review

G. Knezović\*, N. Marić, V. Mijatović, A. Vejnović, V. Pavlović

Faculty of Medicine, University of Novi Sad, Department of Pharmacology, Toxicology and Clinical Pharmacology, Serbia

E-mail address: [goranknezovic@hotmail.com](mailto:goranknezovic@hotmail.com) (G. Knezović).

**Aim:** To evaluate the characteristics of newborns diagnosed with neonatal abstinence syndrome (NAS) and the characteristics

of their mothers in Vojvodina from 2012 to 2016., as well as the interrelationship of certain features.

**Introduction:** NAS is a collection of symptoms and signs that occur as a result of the sudden interruption of fetal exposure to certain substances (methadone, heroin, buprenorphine, etc.) that were used or abused by the mother during pregnancy.<sup>1,2</sup> It is manifested in a multitude of symptoms including central nervous system irritability, over-activity of the vegetative nervous system and dysfunction of the gastrointestinal tract.<sup>3,4</sup> The occurrence of NAS is closely related to the maintenance therapy of pregnant opioid addicts.<sup>1</sup>

**Methods:** This study analyzed medical records of women who gave birth at the Clinic of Gynaecology and Obstetrics in Novi Sad, whose children were diagnosed with NAS after birth, as well as the medical records of newborns treated at the Neonatology Department of the Institute for Child and Youth Health Care of Vojvodina diagnosed with NAS. Medical records included data from the medical history of the newborn and personal and gynaecological medical history of their mothers.

**Results:** A total of 41 cases of NAS were registered. An increase in incidence was noticed during the five-year period of about 15%. Mothers were mostly unemployed (80.49%). Slightly more than half of respondents (57.5%) during pregnancy were on one of substitution treatment modalities. The majority of newborns with NAS (75.61%) were male. The clinical picture was significantly more expressed in children whose mothers consumed methadone, compared to mothers who consumed heroin during pregnancy ( $p=0.0002$ ).

**Conclusion:** The incidence of diagnosed NAS cases is growing. Representation of male newborns with NAS is three times higher than female newborns. Methadone cause more NAS symptoms than heroin.

**Acknowledgements:** The Ministry of Education and Science of Republic Serbia (grant number 41012) supported this research work.

## References

- Kocherlakota P. Neonatal abstinence syndrome. *Pediatrics*. 2014;134:e547–61.
- Cramton REM, Gruchala NE. Babies breaking bad: neonatal and iatrogenic withdrawal syndromes. *Curr Opin Pediatr*. 2013;25:532–42.
- Ko JY, Patrick SW, Tong VT, Patel R, Lind JN, Barfield WD. Incidence of neonatal abstinence syndrome – 28 States, 1999–2013. *Morb Mortal Wkly Rep*. 2016;65:799–802.
- Bersani I, Corsello M, Mastandrea M, Patacchiola V, Foligno S, Garofalo V, et al. Neonatal abstinence syndrome. *Early Hum Dev*. 2013;89:s85–7.

<http://dx.doi.org/10.1016/j.pbj.2017.07.050>

## PS176

### Associations of epicardial adipose tissue thickness and cardiometabolic risk factors in STEMI patients treated with percutaneous coronary intervention

A. Gadeikytė<sup>1,\*</sup>, A. Varoniukaitė<sup>1</sup>, O. Gustienė<sup>2</sup>

<sup>1</sup> Lithuanian University of Health Sciences, Kaunas, Lithuania

<sup>2</sup> Department of Cardiology, Medical Academy, Lithuanian University of Health Sciences, Kaunas, Lithuania

E-mail address: [arvilegadeikyte@gmail.com](mailto:arvilegadeikyte@gmail.com) (A. Gadeikytė).

**Aim:** To evaluate EAT thickness in STEMI patients treated with percutaneous coronary intervention (PCI) and its associations with body mass index (BMI), blood lipids and acute left ventricular dysfunction.



**Introduction:** Epicardial adipose tissue (EAT), located between the myocardium and visceral layer of pericardium is an emerging risk factor for cardiometabolic diseases.

**Methods:** The retrospective study consisted of patients hospitalised for STEMI treated with PCI from 2014 to 2016. EAT thickness was measured from the parasternal long-axis view at end-diastole. Cholesterol levels were determined in a blood sample. According to median patients were divided in two groups: thin EAT group (<2.27 mm,  $n = 270$ ) and thick EAT group ( $\geq 2.27$  mm,  $n = 223$ ). Statistical analysis was performed with SPSS using Mann–Whitney test, *T*-test, logistic regression analysis. Values of cholesterol levels were evaluated by ROC curves.  $p < 0.05$  was significant.

**Results:** Total 492 patients (332 males,  $66.62 \pm 12.24$  year-old) were enrolled. Groups did not differ by age, gender, morbidity of diabetes mellitus and triglyceride levels. Patients had higher BMI ( $29.41 \pm 4.97$  vs.  $28.13 \pm 4.67$  kg/m<sup>2</sup>,  $p = 0.009$ ), total cholesterol ( $>4.82$  mmol/l: 35.2 vs. 26.4%,  $p = 0.024$ ), low density lipoprotein cholesterol ( $>2.5$  mmol/l: 45.8 vs. 33.3%,  $p = 0.004$ ) and reduced high density lipoprotein cholesterol (HDL-C) levels ( $\leq 1$  mmol/l: 24.4 vs. 10.4%,  $p = 0.009$ ) in thick EAT group. Logistic regression analysis revealed that higher BMI (OR = 1.532, 95% CI 1.008–2.328,  $p = 0.002$ ) and HDL-C  $\leq 1$  mmol/l (OR = 1.777, 95% CI 1.159–2.724,  $p = 0.008$ ) were associated with thicker EAT. Killip class  $\geq III$  was more frequent (17.6 vs. 10.3%,  $p = 0.02$ ) in thick than thin EAT group.

**Conclusion:** Increased EAT thickness was associated with obesity, cardiometabolic risk factors and influenced severity of left ventricular dysfunction.

<http://dx.doi.org/10.1016/j.pbj.2017.07.051>

#### PS123

##### Evaluation of spleen volume: Practical diagnostic role of linear measurements, 2D and 3D coefficients in computed tomography



Justyna Tęczar\*, Iwona Kucybała, Anna Gajdosz, Kamil Krupa, Jakub Wnuk, Maria Widomska

Students' Scientific Group at the Department of Diagnostic Imaging, Chair of Radiology, JU CM, Poland

E-mail address: [justyna.teczar@tlen.pl](mailto:justyna.teczar@tlen.pl)

(J. Tęczar).

**Aim:** The aim of the study was to find which linear measurements, field and volume coefficients correlate best with the real volume of the spleen and can be further used for determination of splenomegaly.

**Introduction:** Spleen is involved in a wide spectrum of abnormalities, which might lead to an increase in organ size. Splenic enlargement on CT is diagnosed basing on rather subjective criteria. The product of the length, estimated height and thickness of the spleen ("splenic index", cut-off  $\geq 480$ ) has also been proposed as an indicator for evaluating splenic size on CT.

**Methods:** Abdominal CT examinations of 153 patients' (77 females, 76 males) were retrospectively analysed in terms of maximal length, thickness, hilum thickness (axial plane), height (longest measurement in coronal plane), 90° height (maximum vertical height at coronal section), estimated height (number of axial scans where spleen was visible multiplied by the thickness of CT scans) (Impax Software) and real spleen volume (Vitrea software). Two-dimensional and three-dimensional coefficients were acquired through proper mathematical formulas. Splenomegaly cut-off: 314.5 ml. Pearson's correlation coefficient was calculated for the relationship between single, field, volume measurements and real volume (Statistica software).

**Results:** There was a statistically significant correlation between all single, field and volume measurements and real volume ( $p < 0.05$ ). For single measurements, the correlation is the strongest for height ( $r = 0.813$ , sensitivity 65%, specificity 91.7%, PPV 71.4%, NPV 95.6%). For two-dimensional, it is the coefficient calculated from length and 90° height ( $r = 0.918$ , 85%, 94.7%, 70.8%, 97.7%). For three-dimensional, it is the coefficient calculated from length, 90° height and hilum thickness ( $r = 0.919$ , 75%, 96.2%, 75%, 96.2%). Cut-off for splenic index from our calculations was  $\geq 1148$ .

**Conclusion:** Coefficient from length, 90° height and hilum thickness correlate best with the real volume of the spleen. Splenic index in our study is far from the perfection for clinical practice.

<http://dx.doi.org/10.1016/j.pbj.2017.07.052>

#### PS128

##### Influence of blood inflammatory parameters to erythropoietin resistance in haemodialysis patients



Skirmantė Rėkutė

Vilnius University Medicine Faculty, Lithuania

E-mail address: [skirmante.rekute@gmail.com](mailto:skirmante.rekute@gmail.com).

**Aim:** To evaluate the correlation between the inflammatory blood parameters and the resistance to EPO among the hemodialysis patients.

**Introduction:** Erythropoietin therapy is considered to be the standard treatment of anemia in chronic kidney disease patients, yet some patients do not respond well to this therapy. This is called EPO resistance and could be generally associated with the chronic inflammation.

**Methods:** A retrospective one single centre study, which analysed medical records of 30 HD patients who had advanced CKD and received EPO treatment in Vilnius University Hospital Santaros Clinics from 2016–2009 to 2016–2011. Data analysed – concentrations of C-reactive protein, neutrophils, lymphocytes, platelets, as well as EPO dose per kilo and hemoglobin concentration (measured at the beginning of the EPO therapy and one month after the treatment).

**Results:** Patients were grouped into 2 categories: 1 group ( $n = 14$ ) – concentration of hemoglobin increased, 2 group ( $n = 16$ ) – concentration decreased after treatment. In 1 group average concentration of platelets were statistically significantly ( $p = 0.039$ ) higher ( $230.2 \pm 73.70$ ), compared to 2 group ( $174.1 \pm 66.96$ ).

Furthermore, platelets concentration among patients with hemoglobin level of  $>100$  g/l ( $n = 17$ ) after one month of treatment were statistically significantly ( $p = 0.012$ ) higher ( $231.06 \pm 56.41$ ) compared to those patients with hemoglobin level of  $<100$  g/l ( $n = 13$ ) ( $160.08 \pm 78.17$ ) after treatment.

Additionally, patients with hemoglobin levels after one month of treatment  $100$ – $125$  g/l ( $n = 15$ ) were separated into two groups based on C-reactive protein level:  $>5$  (1 group) and  $<5$  (2 group). Average concentration of erythropoietin was statistically significantly higher in 1 group ( $n = 9$ ) ( $223.82 \pm 69.15$  VV/kg) than in 2 group ( $n = 6$ ) ( $116.68 \pm 59.68$  VV/kg).

Correlation analysis revealed that among patients with hemoglobin levels of  $<110$  g/l after treatment there is a statistically significant positive correlation (+0.428) between change of hemoglobin levels before and after treatment and erythropoietin dose and statistically significant ( $p = 0.023$ ) negative correlation (–0.481) with lymphocytes concentration in blood.

**Conclusion:** HD patients with a higher concentration of platelets respond to EPO therapy better than those with a lower concentration. Increased EPO dose results in higher Hgb