



ORIGINAL ARTICLE

## Cardiac rehabilitation after acute coronary syndrome: Do all patients derive the same benefit?



Sílvia Aguiar Rosa<sup>a,\*</sup>, Ana Abreu<sup>a</sup>, Rui Marques Soares<sup>a</sup>, Pedro Rio<sup>a</sup>,  
Custódia Filipe<sup>b</sup>, Inês Rodrigues<sup>a</sup>, André Monteiro<sup>a</sup>, Cristina Soares<sup>a</sup>,  
Vítor Ferreira<sup>a</sup>, Sofia Silva<sup>a</sup>, Sandra Alves<sup>a</sup>, Rui Cruz Ferreira<sup>a</sup>

<sup>a</sup> Department of Cardiology, Santa Marta Hospital, Lisbon, Portugal

<sup>b</sup> Department of Physiatry, Santa Marta Hospital, Lisbon, Portugal

Received 9 February 2016; accepted 12 September 2016

Available online 24 February 2017

### KEYWORDS

Acute coronary syndrome;  
Cardiac rehabilitation;  
Cardiopulmonary exercise testing

### Abstract

**Introduction:** Cardiac rehabilitation (CR) has been demonstrated to improve exercise capacity in acute coronary syndrome (ACS), but not all patients derive the same benefit. Careful patient selection is crucial to maximize resources.

**Objective:** To identify in a heterogeneous ACS population which patients would benefit the most with CR, in terms of functional capacity (FC), by using cardiopulmonary exercise testing (CPET).

**Methods:** A retrospective analysis of consecutive ACS patients who underwent CR and CPET was undertaken. CPET was performed at baseline and after 36 sessions of exercise. Peak oxygen uptake ( $pVO_2$ ), percentage of predicted  $pVO_2$ , minute ventilation/ $CO_2$  production ( $VE/VCO_2$ ) slope,  $VE/VCO_2$  slope/ $pVO_2$  and peak circulatory power (PCP) ( $pVO_2$  times peak systolic blood pressure) were assessed in two moments. The differences in  $pVO_2$  ( $\Delta pVO_2$ ), % $pVO_2$ , PCP and exercise test duration were calculated. Patients were classified according to baseline  $pVO_2$  (group 1,  $<20$  ml/kg/min vs. group 2,  $\geq 20$  ml/kg/min) and left ventricular ejection fraction (group A,  $<50\%$  vs. group B,  $\geq 50\%$ ).

**Results:** We analyzed 129 patients, 86% male, mean age  $56.3 \pm 9.8$  years. Both group 1 ( $n=31$ ) and group 2 ( $n=98$ ) showed significant improvement in FC after CR, with a more significant increase in  $pVO_2$ , in group 1 ( $\Delta pVO_2$   $4.4 \pm 7.3$  vs.  $1.6 \pm 5.4$ ;  $p=0.018$ ). Significant improvement was observed in CPET parameters in group A ( $n=34$ ) and group B ( $n=95$ ), particularly in  $pVO_2$  and test duration.

**Conclusion:** Patients with lower baseline  $pVO_2$  ( $<20$  ml/kg/min) presented more significant improvement in FC after CR. CPET which is not routinely used in assessment before CR in context of ACS, could be a valuable tool to identify patients who will benefit the most.

© 2017 Sociedade Portuguesa de Cardiologia. Published by Elsevier España, S.L.U. All rights reserved.

\* Corresponding author.

E-mail address: [silviaguiarosa@gmail.com](mailto:silviaguiarosa@gmail.com) (S. Aguiar Rosa).

**PALAVRAS-CHAVE**

Síndrome coronária aguda;  
Reabilitação cardíaca;  
Prova de esforço cardiorrespiratória

**Reabilitação cardíaca após síndrome coronária aguda – terão todos os doentes o mesmo benefício?****Resumo**

**Introdução:** A reabilitação cardíaca (RC) melhora a capacidade funcional (CF) de exercício na síndrome coronária aguda (SCA), contudo nem todos os doentes apresentam benefício idêntico. A criteriosa seleção de doentes é crucial para a rentabilização de recursos.

**Objetivo:** Identificar na população heterogénea de SCA quais os doentes que mais poderiam beneficiar em termos de capacidade funcional (CF) após programa de RC, através de prova de esforço cardiorrespiratória (PECR).

**Métodos:** Realizámos uma análise retrospectiva de doentes consecutivos com SCA que foram submetidos a RC e PECR. A PECR foi realizada previamente e após 36 sessões de treino de exercício. Parâmetros avaliados: consumo de oxigénio no pico (pVO<sub>2</sub>), percentagem do pVO<sub>2</sub> previsto (%pVO<sub>2</sub>), declive ventilação minuto/CO<sub>2</sub> produzido (VE/VCO<sub>2</sub>), declive VE/VCO<sub>2</sub>/pVO<sub>2</sub>, potência circulatória no pico (PCP) (pVO<sub>2</sub> x pressão arterial sistólica no pico). Foram calculadas as variações de pVO<sub>2</sub> ( $\Delta$ pVO<sub>2</sub>). Os doentes foram analisados de acordo com o pVO<sub>2</sub> basal (G1 < 20 versus G2  $\geq$  20 ml/kg/min) e fração de ejeção do ventrículo esquerdo (FEVE) (GA < 50% versus GB  $\geq$  50%).

**Resultados:** Analisámos 129 doentes, 86% homens, idade média 56,3 $\pm$ 9,8 anos. Ambos, G1 (n=31) e G2 (n=98) apresentaram melhoria significativa na CF após RC, com maior aumento significativo de pVO<sub>2</sub> no G1 ( $\Delta$ pVO<sub>2</sub> 4,4 $\pm$ 7,3 versus 1,6 $\pm$ 5,4; p=0,018). Verificou-se melhoria dos parâmetros da PECR em GA (n=34) e GB (n=95), particularmente no pVO<sub>2</sub> e duração da PECR.

**Conclusão:** Doentes com menores pVO<sub>2</sub> (<20 ml/kg/min) apresentaram, após programa de RC, melhoria mais significativa da CF. A PECR, que não é rotina na avaliação de SCA pré RC, poderá constituir um instrumento valioso na identificação dos doentes que mais beneficiam de RC.

© 2017 Sociedade Portuguesa de Cardiologia. Publicado por Elsevier España, S.L.U. Todos os direitos reservados.

%pVO <sub>2</sub>	percentage of predicted pVO <sub>2</sub>
ACS	acute coronary syndrome
CPET	cardiopulmonary exercise testing
CR	cardiac rehabilitation
ET	exercise training
LV	left ventricular
LVEF	left ventricular ejection fraction
PCP	peak circulatory power
pVO <sub>2</sub>	peak oxygen uptake
VE/VCO <sub>2</sub>	minute ventilation/CO <sub>2</sub> production

**Introduction**

Cardiac rehabilitation (CR) has long been considered a cornerstone in secondary prevention after acute coronary syndrome (ACS). It is a coordinated multidisciplinary intervention that aims to improve the cardiac patient's physical, psychological and social functioning. CR programs include exercise training and strategies to reduce modifiable cardiovascular risk factors such as hypertension, dyslipidemia, diabetes and smoking, and to improve adherence to pharmacologic and non-pharmacologic therapy.<sup>1,2</sup>

CR has been shown to reduce reinfarction, cardiovascular and non-cardiovascular readmission and death<sup>3,4,2,5</sup> and

to improve health-related quality of life, and to be cost-effective.<sup>1,6</sup>

Exercise training programs increase patients' cardiorespiratory fitness, reducing symptoms and improving physiologic responses to physical effort.<sup>7</sup> As a consequence, heart rate and blood pressure decrease during physical activity, reducing myocardial oxygen demand.<sup>1</sup>

The positive effects of CR in cardiovascular disease include stabilization of the progression of atherosclerosis and improvement in endothelial function, resulting in increased coronary compliance, elasticity and flow.<sup>1,8-11</sup>

Exercise testing is essential before CR to assess exercise capacity, to prescribe exercise training programs, to stratify cardiovascular risk and to estimate prognosis. After CR, it provides an objective measure of the program's effect on functional capacity.

Achievement of higher functional capacity is clinically significant because it not only impacts on health-related quality of life but it also is prognostically valuable, strongly predicting the risk of death. Assessing fitness at baseline and after CR is important since an improvement in functional capacity lowers the risk of death.<sup>12</sup>

Cardiopulmonary exercise testing (CPET) is the best tool for this purpose, providing a detailed assessment of functional capacity through the analysis of respiratory gas exchange during exercise. Among important CPET parameters is peak oxygen uptake (pVO<sub>2</sub>), which is an accurate measure of exercise capacity and an important predictor of

outcome in patients with coronary artery disease (CAD).<sup>13-15</sup> Peak  $\text{VO}_2$  reflects tissue oxygen uptake in exercise, and depends on cardiac output, arterial oxygen content and oxygen extraction ability of muscle tissue. A supervised exercise training program can increase  $\text{pVO}_2$  by 11-36%, with the greatest improvement in the most deconditioned individuals.<sup>3</sup> Peak  $\text{VO}_2$  varies according to age and gender, so it is also adjusted in relation to the predicted value for age and gender ( $\%\text{pVO}_2$ ).<sup>15</sup>

Another useful parameter, considered by some authors to be a better prognostic parameter than  $\text{pVO}_2$ , is the slope of the ventilation/ $\text{CO}_2$  ratio ( $\text{VE}/\text{VCO}_2$  slope), which expresses ventilatory efficiency and has been shown to be an independent predictor of outcome.<sup>14,16,17</sup> The ratio ( $\text{VE}/\text{VCO}_2$  slope)/ $\text{pVO}_2$  integrates the two above parameters and also has prognostic value.<sup>18</sup>

Another form of assessment is calculation of peak circulatory power (PCP), defined as  $\text{pVO}_2$  times systolic pressure at peak exercise.

Despite its advantages, CPET is not routinely used after ACS.

The aim of the present study was, using CPET parameters, to identify which patients would benefit most in terms of functional capacity after a CR program. Patients were divided according to  $\text{pVO}_2$  and left ventricular ejection fraction (LVEF) at baseline, and the benefit of CR was evaluated by comparing CPET performance before and after the program.

## Methods

### Study design

This was a retrospective observational study of consecutive patients who suffered an ACS and participated in CR after hospital discharge, in whom CPET was performed before and after the program. All patients admitted with ACS were invited to take part in a CR program.

The data were collected between January 2004 and December 2013, in a single center.

The study population was divided into two groups according to baseline functional capacity based on the Weber-Janicki classification<sup>19</sup>: group 1 (G1),  $\text{pVO}_2 < 20$  ml/kg/min; and group 2 (G2),  $\text{pVO}_2 \geq 20$  ml/kg/min. A second analysis was performed, dividing patients according to left ventricular (LV) systolic function before CR, using the cut-point of 50% LVEF: group A (GA), LVEF  $< 50\%$ ; and group B (GB), LVEF  $\geq 50\%$ .

### Cardiac rehabilitation

Early ambulatory phase (phase 2) of CR program was performed by a multidisciplinary team, led by a cardiologist.

Before starting the program, patients were stratified according to CAD risk, exercise capacity and safety concerns. All underwent clinical and echocardiographic assessment and exercise testing.

The exercise program was prescribed on the basis of CPET results and was individualized for each patient based on  $\text{pVO}_2$ , in accordance with current guidelines.<sup>20</sup>

Continuous moderate-intensity exercise training was performed in 36 twice-weekly sessions of 60 min each. Each session included warm-up (10 min), aerobic activity (30 min), muscle-strengthening activity and flexibility exercises (10 min) and cool-down (10 min).

The program was begun at lower intensity (65-70% of maximum training heart rate), and was progressively elevated to a target of 85-95% of maximum training heart rate, without reaching the anaerobic threshold.

Besides the exercise training component, the phase 2 CR program also included:

- identification and control of cardiovascular risk factors, including diabetes, hypertension, dyslipidemia, obesity, smoking, and sedentary lifestyle;
- lifestyle modification with individualized dietary counseling by a dietician, psychologic counseling for selected patients, smoking cessation sessions for some patients, social support in selected cases and promotion of adherence to drug therapy;
- group educational sessions by the multidisciplinary CR team, with the participation of family members, to explain atherosclerotic disease and its consequences and the importance of risk factors and secondary prevention measures.

Immediately after the end of the program, clinical and echocardiographic assessment and exercise testing were repeated, and patients were scheduled for consultations to quantify the benefits obtained the program. Follow-up by telephone was performed to check adherence to healthy behaviors, particularly exercise and medication, control of cardiovascular risk factors and the occurrence of cardiac events.

### Cardiopulmonary exercise testing

All patients underwent CPET at baseline and after 36 exercise training sessions.

Maximal symptom-limited treadmill CPET was performed using the modified Bruce protocol. CPET was monitored with continuous 12-lead electrocardiogram, blood pressure cuff, saturation probe and face mask to measure respiratory gases. Blood pressure was measured at rest, at each stage, at peak exercise and at the first, third and sixth minute of the recovery phase. Respiratory gases were analyzed using an Innocor<sup>®</sup> gas analyzer and  $\text{VO}_2$ ,  $\text{CO}_2$  production ( $\text{VCO}_2$ ) and ventilation (VE) were measured on a breath-by-breath basis.

Patients were encouraged to continue exercising until the  $\text{VCO}_2/\text{VO}_2$  ratio was  $\geq 1.15$ .

Peak  $\text{VO}_2$  adjusted for body mass, or for fat-free mass in obese patients (body mass index  $> 30$  kg/m<sup>2</sup>), was analyzed as  $\%\text{pVO}_2$  according to age and gender. The  $\text{VE}/\text{VCO}_2$  slope was calculated by automatic linear regression with values obtained during the exercise test. The  $\text{VE}/\text{VCO}_2$  slope/ $\text{pVO}_2$  ratio was calculated using the obtained  $\text{VE}/\text{VCO}_2$  slope and  $\text{pVO}_2$ . PCP was calculated as  $\text{pVO}_2$  times peak systolic blood pressure. The differences in  $\text{pVO}_2$ ,  $\%\text{pVO}_2$ , PCP and exercise test duration before and after CR were also calculated.

## Echocardiography

All patients underwent a comprehensive Doppler echocardiographic study using commercially available ultrasound systems (Vivid 9<sup>®</sup>, General Electric) before and after the CR program.

Echocardiographic parameters of LV systolic and diastolic function were assessed, as well as right ventricular function parameters, pulmonary artery systolic pressure and the presence of cardiac valve disease. LV end-diastolic and end-systolic volume and LVEF were determined by Simpson's biplane method.

## Statistical analysis

The statistical analysis was performed using SPSS Statistics (version 22; IBM SPSS, Chicago, IL). Continuous variables were expressed as mean  $\pm$  standard deviation. CPET parameters at baseline and after CR were compared using the two-tailed Student's *t* test or the non-parametric Wilcoxon test, as appropriate. Characteristics of the study groups were compared using the Student's *t* test or the Wilcoxon-Mann-Whitney test for continuous variables and Pearson's chi-square or Fisher's exact test for categorical variables.

## Results

During the study period 4724 patients were admitted with ACS diagnosis. Of these, around 50% were transferred from non-tertiary centers without a catheterization laboratory and were followed in those hospitals after discharge. Of the 2363 (50%) followed in our center, only 12% agreed to be included in the hospital's CR program and completed the program.

The study included the 129 patients who underwent CPET before and after CR, 111 (86.0%) male, mean age 56.3  $\pm$  9.8 years, 88.1% of whom had suffered STEMI. The clinical characteristics of the overall population are presented in [Table 1](#).

Compared to G1, G2 patients (with lower baseline functional capacity) were older, included fewer males and had a higher prevalence of diabetes and dyslipidemia ([Table 1](#)).

CR improved functional capacity in both groups. In G1, pVO<sub>2</sub> increased from 17.7  $\pm$  1.8 to 22.1  $\pm$  7.2 ml/kg/min (*p* < 0.001). PCP and exercise test duration also increased significantly (*p* < 0.001 for both). VE/VCO<sub>2</sub> slope/pVO<sub>2</sub> decreased from 1.7  $\pm$  0.4 to 1.4  $\pm$  0.6 (*p* < 0.001). There was no significant change in VE/VCO<sub>2</sub> slope in this group ([Table 2](#)).

In G2 all CPET parameters improved after CR, pVO<sub>2</sub> rising from 28.1  $\pm$  5.6 to 29.7  $\pm$  6.2 ml/kg/min (*p* < 0.001) and %pVO<sub>2</sub> from 96.3% to 100.3% (*p* = 0.001). PCP and exercise test duration also increased significantly. VE/VCO<sub>2</sub> slope and VE/VCO<sub>2</sub> slope/pVO<sub>2</sub> decreased significantly (*p* = 0.032 and *p* < 0.001, respectively) ([Table 2](#)).

As expected, CPET parameters were substantially different between groups before and after CR, with a better performance in G2. However, when range of improvement is compared, G1 achieved a greater benefit after CR. The improvement in pVO<sub>2</sub> was greater in G1, in absolute value ( $\Delta$ pVO<sub>2</sub> 4.4  $\pm$  7.3 vs. 1.6  $\pm$  5.4; *p* = 0.018) and in percentage of predicted value ( $\Delta$ %pVO<sub>2</sub> 17.9  $\pm$  35.4% vs. 4.0  $\pm$  19.9%; *p* = 0.009). In G1 improvement in pVO<sub>2</sub> was more than 10% in 71% of patients, contrasting with only 40.8% in G2. The increase in PCP was also significantly greater in G1 (883.3  $\pm$  1217.3 vs. 238.5  $\pm$  1250.5 mmHg.ml/kg/min; *p* = 0.015). Exercise test duration increased more in G1, although not significantly ([Table 3](#)).

When patients were grouped according to LV systolic function, both groups presented improved functional capacity after exercise training. In GA, pVO<sub>2</sub> increased from 24.7  $\pm$  6.1 to 27.7  $\pm$  6.9 ml/kg/min (*p* = 0.002), %pVO<sub>2</sub> from 84.9  $\pm$  19.4 to 95.4  $\pm$  24.0% (*p* = 0.001) and exercise test duration from 13.4  $\pm$  2.6 to 15.0  $\pm$  3.3 min (*p* < 0.001) ([Table 2](#)). GB patients also achieved a significant improvement, as demonstrated by increased pVO<sub>2</sub>, %pVO<sub>2</sub>, PCP and exercise test duration and decreased VE/VCO<sub>2</sub> slope and (VE/VCO<sub>2</sub> slope)/pVO<sub>2</sub> ratio ([Table 2](#)).

**Table 1** Clinical characteristics of the overall population and comparison between study groups.

	Overall (n=129)	G1 (n=31)	G2 (n=98)	p	GA (n=34)	GB (n=95)	p
Age (years)	56.3 $\pm$ 9.8	61.3 $\pm$ 7.8	54.8 $\pm$ 9.9	0.001	57.3 $\pm$ 10.5	56.0 $\pm$ 9.6	0.497
Male	86.00%	74.20%	89.80%	0.039	94.10%	83.20%	0.153
Hypertension	52.00%	53.30%	51.50%	0.864	52.90%	51.60%	0.894
Diabetes	14.20%	25.80%	10.40%	0.042	29.40%	8.60%	0.007
Dyslipidemia	59.10%	80.00%	52.60%	0.008	55.90%	60.20%	0.66
Smoking	46.10%	48.40%	45.40%	0.769	35.30%	50.00%	0.14
BMI	26.8 $\pm$ 5.7	26.5 $\pm$ 6.5	26.9 $\pm$ 5.5	0.837	25.9 $\pm$ 7.9	27.2 $\pm$ 4.8	0.898
Family history of CAD	14.80%	9.70%	16.50%	0.562	11.80%	16.00%	0.556
STEMI	88.10%	90.00%	87.50%	0.864	88.20%	88.00%	0.497
No. of vessels with CAD	1.5 $\pm$ 0.7	1.6 $\pm$ 0.8	1.5 $\pm$ 0.6	0.626	1.6 $\pm$ 0.7	1.5 $\pm$ 0.6	0.578
Untreated CAD	26.40%	26.70%	26.30%	0.97	26.50%	26.40%	0.991
LVEF (%)	54.1 $\pm$ 9.9	53.4 $\pm$ 11.9	54.3 $\pm$ 9.2	0.662	41.9 $\pm$ 7.3	58.5 $\pm$ 6.4	<0.001
Sinus rhythm	90.60%	93.50%	89.70%	0.729	85.30%	92.60%	0.3

BMI: body mass index; CAD: coronary artery disease; G1: group 1 (pVO<sub>2</sub> < 20 ml/kg/min); G2: group 2 (pVO<sub>2</sub>  $\geq$  20 ml/kg/min); GA: group A (LVEF < 50%); GB: group B (LVEF  $\geq$  50%); LVEF: left ventricular ejection fraction; STEMI: ST-segment myocardial infarction.

**Table 2** Comparison of cardiopulmonary exercise testing parameters before and after cardiac rehabilitation in the study groups.

CPET parameters	G1 (n=31)			G2 (n=98)			GA (n=34)			GB (n=95)		
	Before CR	After CR	p	Before CR	After CR	p	Before CR	After CR	p	Before CR	After CR	p
Peak VO <sub>2</sub> (ml/kg/min)	17.7 ± 1.8	22.1 ± 7.2	<0.001	28.1 ± 5.6	29.7 ± 6.2	<0.001	24.7 ± 6.1	27.7 ± 6.9	0.002	26.0 ± 6.9	28.0 ± 7.3	0.001
% predicted peak VO <sub>2</sub>	68.6 ± 12.3	86.5 ± 33.0	<0.001	96.3 ± 21.7	100.3 ± 21.8	0.001	84.9 ± 19.4	95.4 ± 24.0	0.001	91.4 ± 24.1	97.6 ± 26.1	0.003
VE/VCO <sub>2</sub> slope	29.2 ± 6.1	28.6 ± 5.8	0.213	25.7 ± 5.5	24.9 ± 4.6	0.031	25.9 ± 5.4	25.5 ± 4.9	0.241	26.8 ± 5.9	26.0 ± 5.2	0.028
PCP (mmHg·ml/kg/min)	2863.7 ± 581.6	4866.4 ± 1213.0	<0.001	4866.4 ± 1213.0	5104.9 ± 1196.3	<0.001	4206.7 ± 1372.4	4583.8 ± 1417.4	0.034	4449.0 ± 1398.0	4848.3 ± 1367.2	<0.001
(VE/VCO <sub>2</sub> slope)/pVO <sub>2</sub>	1.7 ± 0.4	1.4 ± 0.6	<0.001	1.0 ± 0.3	0.9 ± 0.3	0.031	1.1 ± 0.5	1.0 ± 0.6	0.003	1.1 ± 0.4	1.0 ± 0.4	<0.001
Duration (min)	10.8 ± 2.6	13.2 ± 2.4	<0.001	14.2 ± 2.8	15.9 ± 2.5	<0.001	13.4 ± 2.6	15.0 ± 3.3	<0.001	13.4 ± 3.3	15.4 ± 2.5	<0.001

CPET: cardiopulmonary exercise testing; G1: group 1 (pVO<sub>2</sub> <20 ml/kg/min); G2: group 2 (pVO<sub>2</sub> ≥20 ml/kg/min); GA: group A (LVEF <50%); GB: group B (LVEF ≥50%); PCP: peak circulatory power.

There were no significant differences in CPET parameters between these two groups before or after CR. GA presented greater increases in pVO<sub>2</sub> (3.1±5.6 vs. 2.0±6.1 ml/kg/min; p=0.382), %pVO<sub>2</sub> (10.6±19.0 vs. 6.2±26.9%; p=0.291) and exercise test duration (1.6±2.5 vs. 2.0±2.6; p=0.920), although they were not significant.

In patients with initial VE/VCO<sub>2</sub> slope >30, a significant improvement in this parameter was achieved after CR (34.4±5.8 vs. 31.1±5.1; p=0.02).

In a follow-up of 5.17±2.04 years, there were no significant differences in death, cardiac events, lifestyle or adherence to therapy between the study groups (Table 4). The lower mortality and ACS recurrence in all groups is noteworthy. Regarding lifestyle, more than 90% of the overall population maintained therapeutic adherence and more than half of patients performed regular physical exercise and kept to a healthy diet.

## Discussion

The main result of our study was that the CPET parameter pVO<sub>2</sub> is useful in the selection of ACS patients who will have a better functional response to exercise training.

Our data indicate that the inclusion of ACS patients in a supervised 36-session exercise training program improved functional capacity independently of baseline performance, and that patients with pVO<sub>2</sub> <20 ml/kg/min before CR derived greater benefit. The improvement in pVO<sub>2</sub> was seen in absolute terms, as well as adjusted to predicted value for age and gender. As demonstrated in previous papers,<sup>1,21</sup> our data showed that the most deconditioned individuals had the greatest improvement in functional capacity after the program, with 71% of patients in G1 achieving an improvement of >10% in pVO<sub>2</sub> compared to 40.8% in G2.

This greater benefit was also reflected in VE/VCO<sub>2</sub> slope/pVO<sub>2</sub>, PCP and exercise test duration, which have important prognostic implications. Kavanagh et al. demonstrated that the main predictor of outcome in CAD patients is exercise tolerance, as objectively measured by pVO<sub>2</sub>. A 10% decrease in mortality was observed for each increase of 1 ml/kg/min in pVO<sub>2</sub>.<sup>13</sup>

Comparing patients according to baseline LVEF, our research demonstrates that CR is beneficial regardless of LV systolic function. More than half of the patients (55.9%) in GA saw an increase in pVO<sub>2</sub> of >10%, compared to 45.3% in GB. This finding, although not statistically significant, suggests that patients with LV dysfunction may derive more benefit from an exercise training program. Previous studies have shown that patients with LV systolic dysfunction benefit more from CR.<sup>22</sup>

The number of ACS patients undergoing CR was low for various reasons, as analyzed in previous studies, including financial constraints, difficulties in transportation, conflicts with work, low motivation and geographic distance. There are multiple barriers to CR, three of which have been identified as patient-related, physician-related and health system-related factors.<sup>23</sup>

Cardiac rehabilitation, though widely recommended in ACS guidelines, is poorly implemented in several countries, including Portugal, where previous studies have shown that only 3% of ACS patients discharged from hospital are admitted to a CR program.<sup>24-26</sup> Although an increasing number of

**Table 3** Comparison between cardiopulmonary exercise testing parameters before and after cardiac rehabilitation in groups 1 and 2.

CPET parameters	G1 (n=31)	G2 (n=98)	p
Peak VO <sub>2</sub> (ml/kg/min) before CR	17.7±1.8	28.1±5.6	<0.001
Peak VO <sub>2</sub> (ml/kg/min) after CR	22.1±7.2	29.7±6.2	<0.001
% predicted peak VO <sub>2</sub> before CR	68.6±12.3	96.3±21.7	<0.001
% predicted peak VO <sub>2</sub> after CR	86.5±33.0	100.3±21.8	<0.001
VE/VCO <sub>2</sub> slope before CR	29.2±6.1	25.7±5.5	0.003
VE/VCO <sub>2</sub> slope after CR	28.6±5.8	24.9±4.6	0.002
PCP (mmHg.ml/kg/min) before CR	2863.7±581.6	4866.4±1213.0	<0.001
PCP (mmHg.ml/kg/min) after CR	3746.9±1432.0	5104.9±1196.3	<0.001
(VE/VCO <sub>2</sub> slope)/pVO <sub>2</sub> before CR	1.7±0.4	1.0±0.3	<0.001
(VE/VCO <sub>2</sub> slope)/pVO <sub>2</sub> after CR	1.4±0.6	0.9±0.39	<0.001
Duration (min) before CR	10.8±2.6	14.2±2.8	<0.001
Duration (min) after CR	13.2±2.4	15.9±2.5	<0.001
ΔpVO <sub>2</sub>	4.4±7.3	1.6±5.4	0.018
Δ%pVO <sub>2</sub>	17.9±35.4	4.0±19.9	0.009
Increase in pVO <sub>2</sub> >10%	71.0%	40.8%	0.003
Δ CPET duration (min)	2.5±2.8	1.7±2.5	0.101
Δ PCP (mmHg.ml/kg/min)	883.3±1217.3	238.5±1250.5	0.015

Δ: difference; CPET: cardiopulmonary exercise test; CR: cardiac rehabilitation; G1: group 1 (pVO<sub>2</sub> <20 ml/kg/min); G2: group 2 (pVO<sub>2</sub> ≥20 ml/kg/min); PCP: peak circulatory power.

**Table 4** Comparison of follow-up parameters in the study groups.

	Overall (n=129)	G1 (n=31)	G2 (n=98)	p	GA (n=34)	GB (n=95)	p
Death (%)	4.7	3.2	5.1	0.555	8.0	3.8	0.329
All-cause hospitalization (%)	47.9	46.7	48.3	0.876	47.8	47.9	0.994
Cardiac hospitalization (%)	31.8	25.8	33.7	0.412	32.0	31.7	0.979
ACS (%)	6.2	0.0	8.2	0.197	4.0	6.7	0.999
PCI (%)	13.2	6.5	15.3	0.359	4.0	15.4	0.192
CABG (%)	1.6	0.0	2.0	0.999	4.0	1.0	0.351
Adherence to therapy (%)	95.6	92.9	96.5	0.595	100.0	94.6	0.581
Regular exercise (%)	53.5	40.9	57.8	0.170	73.3	49.3	0.090
Healthy diet (%)	54.4	47.6	54.1	0.608	57.1	51.5	0.699
Risk factor control (%)	75.0	68.0	77.2	0.354	75.0	75.0	1.000

ACS: acute coronary syndrome; CABG: coronary artery bypass grafting; G1: group 1 (pVO<sub>2</sub> <20 ml/kg/min); G2: group 2 (pVO<sub>2</sub> ≥20 ml/kg/min); GA: group A (LVEF <50%); GB: group B (LVEF ≥50%); PCI: percutaneous coronary intervention.

ACS patients have been included in CR programs in Portugal in recent years, recruitment in the country remains far below the European average of 30%.<sup>27</sup>

Including more patients in CR programs will require modifications in the health system, including increases in the number of CR centers and improvements in specialized training of health professionals. Failure by cardiologists and surgeons to refer patients for CR is particularly evident in some cases, including the elderly, women, obese patients, smokers and those of low socioeconomic status or suffering from depression.<sup>28</sup> Only with a more suitable health structure and better trained professionals will be possible to recruit and motivate more ACS patients for CR.

Although in our center there is presently no limitation on the number of ACS patients who can be referred for CR, it should be borne in mind that many hospitals do not have

facilities available to provide CR for all patients. Knowing who will derive the greatest benefit will help in the selection of the best responders to CR.

### Study limitations

This is a single-center retrospective study and thus suffers from inherent limitations, including the small number of patients. The size of our sample was due to the fact that of the patients followed after discharge in our center, only a minority agreed to be included in CR at the hospital and completed the program, for the reasons described.

Furthermore, the present study only included patients who performed CPET before and after CR, and since CPET is not a routine procedure, this excluded patients who underwent standard exercise testing, which does not include gas exchange analysis.

## Conclusion

In conclusion, of the heterogeneous group of patients with ACS, those with the worst functional capacity identified by CPET, and those with lower LVEF, obtained the greatest benefit from CR. These individuals should be the main target for referral for comprehensive CR, and CPET may be in fact a valuable tool for their identification.

## Ethical disclosures

**Protection of human and animal subjects.** The authors declare that no experiments were performed on humans or animals for this study.

**Confidentiality of data.** The authors declare that they have followed the protocols of their work center on the publication of patient data.

**Right to privacy and informed consent.** The authors have obtained the written informed consent of the patients or subjects mentioned in the article. The corresponding author is in possession of this document.

## Conflicts of interest

The authors have no conflicts of interest to declare.

## References

- Anderson L, Taylor RS. Cardiac rehabilitation for people with heart disease: an overview of Cochrane systematic reviews. *Cochrane Database Syst Rev.* 2014;12. CD011273.
- Heran BS, Chen JM, Ebrahim S, et al. Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database Syst Rev.* 2011. CD001800.
- Leon AS, Franklin BA, Costa F, et al. Cardiac rehabilitation and secondary prevention of coronary heart disease: an American Heart Association scientific statement from the Council on Clinical Cardiology (Subcommittee on Exercise, Cardiac Rehabilitation, and Prevention) and the Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity), in collaboration with the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation.* 2005;111:369–76.
- Dunlay SM, Pack QR, Thomas RJ, et al. Participation in cardiac rehabilitation, readmissions, and death after acute myocardial infarction. *Am J Med.* 2014;127:538–46.
- Taylor RS, Brown A, Ebrahim S, et al. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *Am J Med.* 2004;116:682–92.
- Clark AM, Hartling L, Vandermeer B, et al. Meta-analysis: secondary prevention programs for patients with coronary artery disease. *Ann Intern Med.* 2005;143:659–72.
- Balady GJ, Williams MA, Ades PA, et al., American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology; American Heart Association Council on Cardiovascular Nursing; American Heart Association Council on Epidemiology and Prevention; American Heart Association Council on Nutrition, Physical Activity, and Metabolism; American Association of Cardiovascular and Pulmonary Rehabilitation. Core components of cardiac rehabilitation/secondary prevention programs: 2007 update: a scientific statement from the American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology; the Councils on Cardiovascular Nursing, Epidemiology and Prevention, and Nutrition, Physical Activity, and Metabolism; and the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation.* 2007;115:2675–82.
- Oliveira NL, Ribeiro F, Alves AJ, et al. The effects of exercise training on arterial stiffness in coronary artery disease patients: a state-of-the-art review. *Clin Physiol Funct Imaging.* 2014;34:254–62.
- Niebauer J, Hambrecht R, Velich T, et al. Attenuated progression of coronary artery disease after 6 years of multifactorial risk intervention: role of physical exercise. *Circulation.* 1997;96:2534–41.
- Hambrecht R, Adams V, Erbs S, et al. Regular physical activity improves endothelial function in patients with coronary artery disease by increasing phosphorylation of endothelial nitric oxide synthase. *Circulation.* 2003;107:3152–8.
- Tanaka H, Dinunno FA, Monahan KD, et al. Aging, habitual exercise, and dynamic arterial compliance. *Circulation.* 2000;102:1270–5.
- Myers J, Prakash M, Froelicher V, et al. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med.* 2002;346:793–801.
- Kavanagh T, Mertens DJ, Hamm LF, et al. Prediction of long-term prognosis in 12 169 men referred for cardiac rehabilitation. *Circulation.* 2002;106:666–71.
- Kavanagh T, Mertens DJ, Hamm LF, et al. Peak oxygen intake and cardiac mortality in women referred for cardiac rehabilitation. *J Am Coll Cardiol.* 2003;42:2139–43.
- Wasserman K, Hansen JE, Sue DY, et al. Principles of exercise testing and interpretation. 3rd ed. Baltimore: Lippincott Williams & Wilkins; 1999.
- Ferreira AM, Tabet JY, Frankenstein L, et al. Ventilatory efficiency and the selection of patients for heart transplantation. *Circ Heart Fail.* 2010;3:378–86.
- Arena R, Myers J, Aslam SS, et al. Peak  $\dot{V}O_2$  and  $\dot{V}E/\dot{V}CO_2$  slope in patients with heart failure: a prognostic comparison. *Am Heart J.* 2004;147:354–60.
- Guazzi M, De Vita S, Cardano P, et al. Normalization for peak oxygen uptake increases the prognostic power of the ventilatory response to exercise in patients with chronic heart failure. *Am Heart J.* 2003;146:542–8.
- Weber KT, Kinasevitz GT, Janicki JS, et al. Oxygen utilization and ventilation during exercise in patients with chronic cardiac failure. *Circulation.* 1982;65:1213–23.
- Pescatello LS, Arena R, Riebe D, et al. Exercise prescription for patients with cardiovascular and cerebrovascular disease. American College of Sports Medicine. ACSM's Guidelines for Exercise Testing and Prescription. 9th ed. Philadelphia: Lippincott Williams & Wilkins; 2014. p. 240–52.
- Ades PA. Cardiac rehabilitation and secondary prevention of coronary heart disease. *N Engl J Med.* 2001;345:892–902 [review].
- Zhang YM, Lu Y, Tang Y, et al. The effects of different initiation time of exercise training on left ventricular remodeling and cardiopulmonary rehabilitation in patients with left ventricular dysfunction after myocardial infarction. *Disabil Rehabil.* 2016;38:268–76.
- Conraads VM, Deaton C, Piotrowicz E, et al. Adherence of heart failure patients to exercise: barriers and possible solutions: a position statement of the Study Group on Exercise Training in Heart Failure of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail.* 2012;14:451–8.
- Abreu A, Bettencourt N, Fontes P. Overview of cardiac rehabilitation in Portugal 2007–2009. *Rev Port Cardiol.* 2010;29:545–58.

25. Mendes M. Reabilitação cardíaca em Portugal: a intervenção que falta! *Saúde Tecnol.* 2009;3:5–9.
26. Reabilitação Cardíaca: Realidade Nacional e Recomendações Clínicas. Coordenação Nacional para as Doenças Cardiovasculares, Ministério da Saúde; 2009.
27. Bjarnason-Wehrens B. Results from the European Cardiac Rehabilitation Inventory Survey (ECRIS). In: *Europrevent 2008*. Brussels: European Society of Cardiology; 2008.
28. Mendes M. Barriers to participation in cardiac rehabilitation programs. *Rev Port Cardiol.* 2011;30:509–14.